
Score vs selected-PSM stage features (position)
* BH q < 0.05 within outcome/scenario/method; all three predictors and stages
Between participants
Within participant
Within participant
adjusted for task order + attempt
A: Reach Ring (selected PSM)
A: Grab Ring (selected PSM)
A: Transfer Ring (selected PSM)
A: Place Ring (selected PSM)
A: Drop Ring (selected PSM)
C: Reach String (selected PSM)
C: Threading (selected PSM)
C: Wrap around PSM (selected PSM)
C: Pull/cinch (selected PSM)
-0.22 +0.39 +0.12
+0.03 +0.22 +0.25
+0.08 +0.03 +0.12
-0.11 +0.08 +0.50
+0.28 +0.13 +0.13
-0.01 +0.32 +0.28
-0.77 -0.79 +0.37
+0.48 +0.38 +0.09
+0.09 -0.01 +0.10
+0.23 +0.13 +0.11
-0.09 -0.05 -0.07
+0.16 -0.05 -0.06
excluded excluded excluded
excluded excluded excluded
excluded excluded excluded
-0.16 +0.17 -0.25
+0.10 -0.08 -0.28
+0.10 -0.04 -0.18
no score variation
no score variation
no score variation
no score variation
no score variation
no score variation
no score variation
no score variation
no score variation
+0.14 -0.61 +0.63
+0.05 +0.23 +0.10
-0.15 +0.09 -0.15
excluded excluded excluded
excluded excluded excluded
excluded excluded excluded
Signed lag
Absolute lag
Peak r
Rank association
1.00
0.75
0.50
0.25
0.00
−0.25
−0.50
−0.75
−1.00
One PSM chosen per recording/stage by largest signed xcorr peak, before score matching.
Positive signed lag = gaze leads. Scores describe whole attempts. Missing outcomes are excluded.
Selected PSM indicates stronger correlation, not necessarily the operated instrument.