
John Clouse IMD HW 4 Problem 3

Table of Contents

Initialize	1
Planet positions and velocities, V_{∞} in and out	1
Flyby params	1
Energy	1
Results	2

Initialize

```
clearvars -except hw_pub function_list

CelestialConstants;

JD_launch = 2447807.5;
JD_Venus = 2447932.5;
JD_Earth1 = 2448235.5;
```

Planet positions and velocities, V_{∞} in and out

```
[r_earth_L,~] = MeeusEphemeris(Earth, JD_launch,Sun);
[r_venus, v_venus] = MeeusEphemeris(Venus,JD_Venus,Sun);
[r_earth_1,~] = MeeusEphemeris(Earth, JD_Earth1,Sun);

[~,v_in] = lambert(r_earth_L, r_venus,(JD_Venus-JD_launch)*day2sec,1,Sun);
[v_out,~] = lambert(r_venus, r_earth_1,(JD_Earth1-JD_Venus)*day2sec,-1,Sun);

v_inf_in = v_in - v_venus;
v_inf_out = v_out - v_venus;
```

Flyby params

```
psi = acos(dot(v_inf_in,v_inf_out)/norm(v_inf_in)/norm(v_inf_out));
rp = Earth.mu/(norm(v_inf_in)^2)*(1/cos((pi-psi)/2)-1);
hp = rp - Venus.R; %km
```

Energy

```
energy_pre_flyby = norm(v_in)^2/2 - Sun.mu/norm(r_venus);
energy_post_flyby = norm(v_out)^2/2 - Sun.mu/norm(r_venus);
percent_change = (energy_post_flyby - energy_pre_flyby)...
    /abs(energy_pre_flyby)*100;
```

Results

The v-infinities do not match exactly because the events aren't precisely targeted. True event JDs can be found to make them line up even better. The altitude of closes approach to Venus is shown below. The energy changed by 15%, making a trajectory with a higher aphelion.

```
fprintf('Closest approach altitude: %.3f km\n',hp);  
fprintf('Heliocentric energy before: %.3f km^2/s^2\n',energy_pre_flyby);  
fprintf('Heliocentric energy after: %.3f km^2/s^2\n',energy_post_flyby);
```

```
Closest approach altitude: 17105.222 km  
Heliocentric energy before: -528.482 km^2/s^2  
Heliocentric energy after: -448.317 km^2/s^2
```

Published with MATLAB® R2013b