
HW8 Problem 2

Table of Contents

Initialize	1
CKF State	1
Joseph State	2
Potter State	2
Batch State	3

John Clouse

Initialize

```
clear
close all
logx_fig = figure;
loglog_fig = figure;
% Epsilon values
len = 10;
eps = ones(len,1)*1e-5; % start at 1e-6 after the loop iterates
for ii = 1:len
    eps(ii:end) = eps(ii:end)/10;
end
std_dev = 1./eps;
R = 1;

X_exact = [3;1];
X_ap0 = [4;2]; % a priori state
```

CKF State

```
X_ckf = zeros(2,len);
diff_ckf = zeros(2,len);
for ii = 1:len
    H1 = [1 2*eps(ii)];
    P1_ap = eye(2)*std_dev(ii)*std_dev(ii);
    K1 = P1_ap*H1'*inv(H1*P1_ap*H1' + R);
    X_est1 = X_ap0 + K1*(H1*X_exact - H1*X_ap0);
    P1 = (eye(2) - K1*H1)*P1_ap;
    X_ap1 = X_est1;

    %P1 is now P2_ap
    P2_ap = P1;
    H2 = [1 3*eps(ii)];
    K2 = P2_ap*H2'*inv(H2*P2_ap*H2' + R);
    X_est2 = X_ap1 + K2*(H2*X_exact - H2*X_ap1);
    P2 = (eye(2) - K2*H2)*P2_ap;
    X_ckf(:,ii) = X_est2;
```

```
diff_ckf(:,ii) = X_exact - X_ckf(:,ii);
end

row = 2;
HW8_P2_plot(eps, -diff_ckf, row, 'Kalman', logx_fig, loglog_fig)
```

Joseph State

```
I = eye(2);
X_joseph = zeros(2,len);
diff_joseph = zeros(2,len);
for ii = 1:len
    H = [1 2*eps(ii); 1 3*eps(ii)];
    P_ap = I*std_dev(ii)*std_dev(ii);
    X_ap = X_ap0;
    for jj = 1:2
        K = P_ap*H(jj,:)'*inv(H(jj,:)*P_ap*H(jj,:) + R);
        P = (I-K*H(jj,:))*P_ap*(I-K*H(jj,:))' + K*R*K';
        X_est = X_ap + K*(H(jj,:)*X_exact - H(jj,:)*X_ap);
        P_ap = P; % a priori for next measurement
        X_ap = X_est;
    end
    X_joseph(:,ii) = X_est;
    diff_joseph(:,ii) = X_exact - X_joseph(:,ii);
end
row = 3;
HW8_P2_plot(eps, -diff_joseph, row, 'Joseph', logx_fig, loglog_fig)
```

Potter State

```
X_potter = zeros(2,len);
diff_potter = zeros(2,len);
for ii = 1:len
    H = [1 2*eps(ii); 1 3*eps(ii)];
    P_ap = I*std_dev(ii)*std_dev(ii);
    X_ap = X_ap0;
    W_bar = sqrt(P_ap);
    for jj = 1:2
        F = W_bar*H(jj,:)';
        alpha = inv(F'*F + R);
        gamma = 1/(1+sqrt(R*alpha));
        K = alpha*W_bar*F;
        W = W_bar-gamma*K*F';
        X_est = X_ap + K*(H(jj,:)*X_exact - H(jj,:)*X_ap);
        W_bar = W; % sequential update
        X_ap = X_est;
    end
    X_potter(:,ii) = X_est;
    diff_potter(:,ii) = X_exact - X_potter(:,ii);
end
row = 4;
HW8_P2_plot(eps, -diff_potter, row, 'Potter', logx_fig, loglog_fig)
```

Batch State

```
X_batch = zeros(2,len);
diff_batch = zeros(2,len);
for ii = 1:len
    H = [1 2*eps(ii); 1 3*eps(ii)];
    info_mat = inv(I*std_dev(ii)*std_dev(ii));
    info_mat = info_mat + H(1,:)'*inv(R)*H(1,:);
    info_mat = info_mat + H(2,:)'*inv(R)*H(2,:);

    N = inv(I*std_dev(ii)*std_dev(ii))*X_ap0;
    N = N + H(1,:)'*inv(R)*H(1,:)*X_exact;
    N = N + H(2,:)'*inv(R)*H(2,:)*X_exact;
    X_est = inv(info_mat)*N;
    X_batch(:,ii) = X_est;
    diff_batch(:,ii) = X_exact - X_batch(:,ii);
end
row = 1;
HW8_P2_plot(eps, -diff_batch, row, 'Batch', logx_fig, loglog_fig)
```

Published with MATLAB® R2013b