

Matgeo Presentation

Pushkar Gudla
AI24BTECH11012
IIT Hyderabad.

November 5, 2024

Contents

1 Problem

2 Solution

- Setup and Variable Definitions
- Converting to Matrix form
- Parametric Form of the Line
- Finding Points of Intersection
- Finding Area through Integration

3 Codes

- C code to verify the Area
- Generating Parabola and Plotting the Figure

4 Figure

Problem Statement

The area of the region bounded by the curve $x^2 = 4y$ and the straight line $x = 4y - 2$ is

Setup and Variable Definitions

Variable	Description
$g(x)$	Equation of Conic
L	Equation of line
h	A point on line L
m	Direction vector of line L
x_1 and x_2	Points of intersection of L and $g(x)$

Table: Variables and given data

Converting the equations to Matrix form

To simplify the calculation of points of intersection, the curve $x^2 = 4y$ is represented in standard conic matrix form:

$$g(x) = x^T \mathbf{V}x + 2\mathbf{u}^T x + f = 0 \quad (3.1)$$

$$\mathbf{V} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad (3.2)$$

$$\mathbf{u} = \begin{pmatrix} 0 \\ 2 \end{pmatrix} \quad (3.3)$$

$$f = 0 \quad (3.4)$$

Parametric Form of the Line

$$L : \mathbf{x} = \mathbf{h} + k\mathbf{m} \quad (3.5)$$

where k is a scalar constant

$$\mathbf{h} = \begin{pmatrix} -2 \\ 0 \end{pmatrix} \quad (3.6)$$

$$\mathbf{m} = \begin{pmatrix} 1 \\ \frac{1}{4} \end{pmatrix} \quad (3.7)$$

$$\mathbf{x}_i = \mathbf{h} + k_i\mathbf{m} \quad (3.8)$$

Finding Points of Intersection

On substituting L in $g(x)$ we get a quadratic equation in k . We find that the discriminant is not equal to zero and thus we get two values k_1 and k_2 . The point obtained by substituting k_1 in L is \mathbf{x}_1 and similarly on substituting k_2 in L we get \mathbf{x}_2 .

$$k_1 = \frac{1}{\mathbf{m}^\top \mathbf{V} \mathbf{m}} \left(-\mathbf{m}^\top (\mathbf{V} \mathbf{h} + \mathbf{u}) + \sqrt{[\mathbf{m}^\top (\mathbf{V} \mathbf{h} + \mathbf{u})]^2 - g(\mathbf{h}) (\mathbf{m}^\top \mathbf{V} \mathbf{m})} \right) \quad (3.9)$$

$$k_2 = \frac{1}{\mathbf{m}^\top \mathbf{V} \mathbf{m}} \left(-\mathbf{m}^\top (\mathbf{V} \mathbf{h} + \mathbf{u}) - \sqrt{[\mathbf{m}^\top (\mathbf{V} \mathbf{h} + \mathbf{u})]^2 - g(\mathbf{h}) (\mathbf{m}^\top \mathbf{V} \mathbf{m})} \right) \quad (3.10)$$

On solving for k_1 and k_2 we get $k_1 = 4$ and $k_2 = 1$. Thus

$$\mathbf{x}_1 = \begin{pmatrix} 2 \\ 1 \end{pmatrix} \quad (3.11)$$

$$\mathbf{x}_2 = \begin{pmatrix} -1 \\ \frac{1}{4} \end{pmatrix} \quad (3.12)$$

Finding Area through Integration

The area between the curve and the line from $x = -1$ to $x = 2$ is calculated using the following integration:

$$\text{Area} = \int_{-1}^2 \left(\frac{x}{4} + \frac{1}{2} - \frac{x^2}{4} \right) dx$$

Solving this integral, we get $\text{Area} = \frac{9}{8}$.

Hence, the area bound between the curve $x^2 = 4y$ and the line $x = 4y - 2$ is $\frac{9}{8}$.

C code to verify the Area

We can use the following C code to verify that the area we found is correct.

```
https://github.com/GPushkar16/EE1030/blob/main/Matgeo/question-4/codes/area\_between\_curves.c
```

Generating Parabola and Plotting the Figure

We can use the following C code to generate points that lie on the parabola:

```
https://github.com/GPushkar16/EE1030/blob/main/Matgeo/question-4/codes/generate\_parabola.c
```

We can then use the following Python code to generate the figure:

```
https://github.com/GPushkar16/EE1030/blob/main/Matgeo/question-4/codes/plot.py
```

Figure

