The evidenced agriculture problem

Season A 2026, NISR Seasonal Agricultural Survey. National maize is about 2.0 tonnes per hectare. Parliament's target was 2.26. The miss is not in every district, and it is not a storage problem in this report.

2.0 vs 4.3

Maize tonnes per hectare, all farms vs large farms, Season A 2026

Table 13. Smallholders 1.90, all farms 1.99, large farms 4.34. Beans are 0.70 nationally and 0.54 on large farms.

Two district facts, both printed

District	Maize t/ha	Against 2.26	Smallholders irrigating	Large farms irrigating
Nyagatare	2.89	Above target	6.4%	50%
Kirehe	2.52	Above target	14.9%	78.6%
Kamonyi	1.86	Below target	22.4%	93.3%

Yield from Table 13. Kamonyi's large-farm yield is Table 14, 6.47 t/ha. Irrigation from Table 31. Nyagatare and Kirehe are the only two districts at or above 2.26.

What the same report says about the other levers

Lever	What is published	What a district card may say
Irrigation	12.6% of smallholders and 60.3% of large farms, Table 31. The gap is inside districts.	Where the smallholder rate is low and the large-farm rate is high, irrigation is the measured gap.
Improved seed	Table 20 is all crops: 35.2% of smallholders, 82.6% of large farms. Table 21 is maize, 64.3% improved, with no district and no farm-size split.	An all-crop gap. Not a maize-seed finding, and not a variety.
Inorganic fertilizer	Table 25, by district and farm type. In Gakenke, smallholders are ahead of large farms.	Name it only where smallholders lag in that district.
Storage	Table 16: maize post-harvest loss is 0.3% of use, one national row.	No district store ranking.

The screen that follows

For each district, show maize yield against 2.26, the irrigation pair, the all-crop seed pair, and the fertilizer pair, each with its table. The action is irrigation only when that gap is the one the cells support. Otherwise the action is the input gap, labeled all crops. A dry-season rainfall layer can gate the irrigation action. It cannot replace a NISR cell.