

Synthesis of (P[^]N[^]C)Gold(III) Complexes via Tandem Oxidative Addition/C–H Auration

Michał Biedrzycki, Jaime Martín, Alexandre Genoux, Elia Boschi, and Cristina Nevado*

Cite This: *ACS Org. Inorg. Au* 2025, 5, 322–327

Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

ABSTRACT: A new method for the synthesis of cyclometalated gold(III) complexes featuring (P[^]N[^]C) ligands is reported. This protocol employs a tandem oxidative addition/cycloauration process, enabling the formation of (P[^]N[^]C)gold(III)-aryl, -alkenyl and -alkynyl derivatives by creating two Au–C bonds in one pot from a structurally diverse range of C(sp²)- and C(sp)-iodide substrates. Additionally, a complementary two-step method was developed to access (P[^]N[^]C)gold(III)-alkynyl complexes, which exhibit emission in the blue and green light regions.



- ✦ Tandem oxidative addition / cycloauration
- ✦ One pot formation of two Au–C bonds
- ✦ Synthesis of gold(III)-Csp²/Csp complexes
- ✦ Photoluminescent properties

KEYWORDS: gold, ligands, oxidative addition, cyclometalation, phosphorescence

Gold(III) complexes have emerged as versatile tools in catalysis,^{1–4} medicine^{5–8} and material science.^{9–12} One of the major challenges to advance these fields is to stabilize gold in high oxidation states, given its high redox potentials ($E_{\text{red}}^{\text{Au(III)/Au(I)}} = +1.41$ V and $E_{\text{red}}^{\text{Au(III)/Au(0)}} = +1.50$ V in aqueous solution)¹³ and its subsequent propensity to undergo reductive elimination. Bi- and tridentate ligands, predominantly featuring carbon and nitrogen donor groups, present an effective strategy to modify the redox properties of the metal and stabilize gold(III) via cyclometalation.¹⁴ Interestingly, these complexes exhibit unique catalytic performances,^{15–19} promising antitumoral activities,^{20–24} and versatile photochemical properties.^{25–28}

Seminal work by Constable on the synthesis of cyclometalated (N[^]C)gold(III) compounds via transmetalation from organomercury precursors²⁹ inspired the widespread incorporation of (N[^]C) ligands into gold by innovative synthetic methods. Some of these involve oxidative addition of aryl diazonium salts and aryl iodides to gold(I) precursors, and direct cycloauration under microwave and high temperature conditions (Figure 1, top left).^{30–34} The introduction of an additional phenyl ring to access biscyclometalated (N[^]C[^]C)gold(III) derivatives^{35–42} was attained by our group using microwave irradiation (Figure 1, top right).³⁵

Bidentate (P[^]N) ligands, featuring both soft and hard coordination sites, represent an appealing alternative to conventional (N[^]C) templates for the stabilization of gold species^{43–47} and have proved fruitful in the implementation of gold(I)/gold(III) redox catalytic cycles.^{48–54} (P[^]N)gold(III) complexes are typically synthesized by coordination of a gold(I) precursor to the phosphine group followed by oxidative addition of aryl diazonium salts upon blue light

irradiation^{55,56} or of aryl iodides in the presence of silver salts (Figure 1, bottom left).⁵⁷

Recently, our group disclosed the extension of these systems with a straightforward synthesis of (P[^]N[^]C) pincer ligands.^{58,59} Coordination of the phosphine group to gold(I), followed by oxidation with Selectfluor enabled the cycloauration step delivering the corresponding (P[^]N[^]C)gold(III) chlorides (Figure 1, bottom right). These species underwent facile exchange of the chloro ligand, enabling the structural characterization of elusive gold(III) hydride, hydroxo and formate species.

We hypothesized that the κ^1 -P coordination of (P[^]N[^]C) ligands to a soft gold(I) center would stabilize a cationic gold intermediate which, upon oxidative addition of aryl-halides, would facilitate the coordination of the N quinoline group and trigger the concomitant Csp²-H bond activation of the phenyl ring delivering the cyclometalated species.


Herein, we report the successful implementation of this concept showcasing the synthesis of (P[^]N[^]C)gold(III)-aryl, -alkenyl and -alkynyl derivatives via oxidative addition of Csp²-I and Csp-I bonds onto (P[^]N[^]C)gold(I) precursors. This sequence results in the formation of tridentate (P[^]N[^]C)gold(III) complexes in the absence of external oxidants, following a tandem oxidative addition/cycloauration


Received: June 4, 2025

Revised: July 10, 2025

Accepted: July 11, 2025

Published: August 23, 2025





less sterically congested regioisomers were exclusively obtained (**1g**, 77%; **1h**, 75%).

This methodology was also compatible with other Csp^2 -based electrophiles (Figure 3). When 1 equiv of vinyl iodide was used in combination with AgBF_4 , the corresponding ($\text{P}^{\wedge}\text{N}^{\wedge}\text{C}$)gold(III)-vinyl complex **2a** was obtained in 99% yield. Furthermore, oxidative addition of 1,2-disubstituted alkenyl iodides proceeded with retention of the configuration, as demonstrated by the reactions of (*E*)- β -iodostyrene and (*Z*)- β -iodostyrene, which produced the corresponding ($\text{P}^{\wedge}\text{N}^{\wedge}\text{C}$)gold(III)-alkenyl complexes **2b** and **2c** in 82 and 96% yields, respectively. The ^1H NMR spectrum of **2b** in dichloromethane- d_2 at 298 K exhibited a doublet of doublets at 7.16 ppm ($J_{\text{HH}} = 16.2$ Hz, $J_{\text{HP}} = 1.7$ Hz) corresponding to a vinylic proton, consistent with a *trans* configuration of the double bond. In contrast, the vinylic signals of **2c** overlapped with the aromatic region of the ($\text{P}^{\wedge}\text{N}^{\wedge}\text{C}$) ligand, while the two adamantyl groups became inequivalent, unlike in compound **2b**. This observation suggests that the alkenyl group in **2c** lies perpendicular to the ($\text{P}^{\wedge}\text{N}^{\wedge}\text{C}$)Au moiety due to steric constraints imposed by the double bond configuration, in line with previous hydroauration products reported for this ligand system.⁵⁹

The excellent stability of these new ($\text{P}^{\wedge}\text{N}^{\wedge}\text{C}$)gold(III) complexes⁶⁰ prompted us to investigate their photophysical behavior. Tridentate ($\text{C}^{\wedge}\text{N}^{\wedge}\text{C}$) and ($\text{N}^{\wedge}\text{C}^{\wedge}\text{C}$)gold(III)-alkynyl complexes are known for their remarkable emissive properties and some have been used as dopants in OLED devices.^{26,61} Interestingly, the tandem oxidative addition/cycloauration protocol with phenylethynyl iodide and silver tetrafluoroborate allowed us to prepare the corresponding ($\text{P}^{\wedge}\text{N}^{\wedge}\text{C}$)gold(III)-alkynyl compound **4a**, demonstrating versatility of this method to incorporate not only $\text{C}(\text{sp}^2)$ -but also $\text{C}(\text{sp})$ -I based reagents (Figure 4, top).⁶⁰ However, the isolation of gold(III) complexes with more complex alkynyl iodides using this approach proved challenging, prompting the development of an alternative route via a gold(III)-halide precursor.^{25,26,61} Thus, using ($\text{P}^{\wedge}\text{N}^{\wedge}\text{C}$)gold(III)-chloride precursor **3**,⁵⁸ a copper-mediated transmetalation process with various terminal alkynes delivered the corresponding alkynyl derivatives **4a**, **4b**, **4c** and **4d** in 90, 95, 74, and 75% yields, respectively (Figure 4, top).

Complex **4a** emitted green light with maxima at 506, 545, and 588 nm when excited at $\lambda \leq 375$ nm in degassed dichloromethane at room temperature (quantum yield




Figure 3. Synthesis of cyclometalated ($\text{P}^{\wedge}\text{N}^{\wedge}\text{C}$)Au(III)alkenyl complexes **2a–c** by oxidative addition of alkenyl iodides.




Figure 4. Synthesis of ($\text{P}^{\wedge}\text{N}^{\wedge}\text{C}$)Au(III)-alkynyl complexes **4a–d**. Absorption (black) and emission (red) spectra in dichloromethane.

$\phi(\text{CH}_2\text{Cl}_2) = 2.4\%$). In contrast, compounds **4b** and **4c** exhibited blue light emissions with maxima at 445 and 407 nm when excited at $\lambda \leq 345$ nm, with quantum yields of 1.2% and 0.8%, respectively. Complex **4d** displayed a weak emission with a maximum at 510 nm and quantum yield below the detection threshold (Figure 4, bottom).^{26,61} The quantum yields of these complexes are comparable to other luminescent cationic gold(III) complexes reported by the groups of Venkatesan (0.3–0.9%)⁶² and Yam (0.1–0.6%)⁶³ in solution at room temperature. The different emission maxima in the green or blue region of our complexes showcase the versatility of the ligand scaffold to tune the emission properties of gold(III) species.

In conclusion, this study presents a novel approach to cyclometalated gold(III) complexes featuring ($\text{P}^{\wedge}\text{N}^{\wedge}\text{C}$) ligands. The tandem oxidative addition/cycloauration protocol accommodates both electron-withdrawing and electron-donating groups on $\text{C}(\text{sp}^2)$ -iodides and ($\text{P}^{\wedge}\text{N}^{\wedge}\text{C}$) ligands, enabling the synthesis of a wide range of derivatives including gold(III)-aryl, -alkenyl and -alkynyl complexes. Additionally, an alternative two-step method facilitated the preparation of ($\text{P}^{\wedge}\text{N}^{\wedge}\text{C}$)gold(III)-alkynyl complexes, which exhibited emissive properties in the blue and green light regions.

■ ASSOCIATED CONTENT

Data Availability Statement

The data underlying this study are available in the published article and its [Supporting Information](#).

■ Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsorginorgau.5c00057>.

Experimental procedures, synthesis and characterization details of the new compounds, NMR spectra, photo-physical properties and X-ray diffraction analyses ([PDF](#))

Accession Codes

Deposition Numbers [2451717](#) and [2451722](#) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge via the joint Cambridge Crystallographic Data Centre (CCDC) and Fachinformationszentrum Karlsruhe [Access Structures service](#).

■ AUTHOR INFORMATION

Corresponding Author

Cristina Nevado — Department of Chemistry, University of Zurich, Zurich CH 8057, Switzerland; orcid.org/0000-0002-3297-581X; Email: cristina.nevado@chem.uzh.ch

Authors

Michał Biedrzycki — Department of Chemistry, University of Zurich, Zurich CH 8057, Switzerland

Jaime Martín — Department of Chemistry, University of Zurich, Zurich CH 8057, Switzerland; orcid.org/0000-0003-0909-3509

Alexandre Genoux — Department of Chemistry, University of Zurich, Zurich CH 8057, Switzerland

Elia Boschi — Department of Chemistry, University of Zurich, Zurich CH 8057, Switzerland; orcid.org/0000-0002-1429-4949

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acsorginorgau.5c00057>

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We thank Prof. Dr. Anthony Linden for the X-ray structure analysis of compounds **1a** (CCDC 2451717) and **3** (CCDC 2451722).

■ REFERENCES

- (1) Huang, B. R.; Hu, M. Y.; Toste, F. D. Homogeneous Gold Redox Chemistry: Organometallics, Catalysis, and Beyond. *Trends Chem.* **2020**, *2*, 707–720.
- (2) Rocchigiani, L.; Bochmann, M. Recent Advances in Gold(III) Chemistry: Structure, Bonding, Reactivity, and Role in Homogeneous Catalysis. *Chem. Rev.* **2021**, *121*, 8364–8451.
- (3) Praveen, C.; Dupeux, A.; Michelet, V. Catalytic Gold Chemistry: From Simple Salts to Complexes for Regioselective C–H Bond Functionalization. *Chem. A Eur. J.* **2021**, *27*, 10495–10532.
- (4) Font, P.; Ribas, X. Fundamental Basis for Implementing Oxidant-Free Au(I)/Au(III) Catalysis. *Eur. J. Inorg. Chem.* **2021**, *2021*, 2556–2569.
- (5) Tong, K.-C.; Lok, C.-N.; Wan, P.-K.; Hu, D.; Fung, Y. M. E.; Chang, X.-Y.; Huang, S.; Jiang, H.; Che, C.-M. An anticancer

gold(III)-activated porphyrin scaffold that covalently modifies protein cysteine thiols. *Proc. Natl. Acad. Sci. U.S.A.* **2020**, *117*, 1321–1329.

(6) Lu, Y. L.; Ma, X. Y.; Chang, X. Y.; Liang, Z. L.; Lv, L.; Shan, M.; Lu, Q. Y.; Wen, Z. F.; Gust, R.; Liu, W. K. Recent development of gold(I) and gold(III) complexes as therapeutic agents for cancer diseases. *Chem. Soc. Rev.* **2022**, *51*, 5518–5556.

(7) Moreno-Alcántar, G.; Picchetti, P.; Casini, A. Gold Complexes in Anticancer Therapy: From New Design Principles to Particle-Based Delivery Systems. *Angew. Chem., Int. Ed.* **2023**, *62*, No. e202218000.

(8) Mertens, R. T.; Gukathasan, S.; Arojoye, A. S.; Olewe, C.; Awuah, S. G. Next Generation Gold Drugs and Probes: Chemistry and Biomedical Applications. *Chem. Rev.* **2023**, *123*, 6612–6667.

(9) Yam, V. W.-W.; Law, A. S.-Y. Luminescent d8 metal complexes of platinum(II) and gold(III): From photophysics to photofunctional materials and probes. *Coord. Chem. Rev.* **2020**, *414*, No. 213298.

(10) Tang, M.-C.; Chan, M.-Y.; Yam, V. W.-W. Molecular Design of Luminescent Gold(III) Emitters as Thermally Evaporable and Solution-Processable Organic Light-Emitting Device (OLED) Materials. *Chem. Rev.* **2021**, *121*, 7249–7279.

(11) Malmberg, R.; Venkatesan, K. Conceptual advances in the preparation and excited-state properties of neutral luminescent (C⁺N) and (C⁺C*) monocyclometalated gold(III) complexes. *Coord. Chem. Rev.* **2021**, *449*, No. 214182.

(12) Yam, V. W.-W.; Chan, A. K.-W.; Hong, E. Y.-H. Charge-transfer processes in metal complexes enable luminescence and memory functions. *Nat. Chem. Rev.* **2020**, *4*, 528–541.

(13) Bratsch, S. G. Standard Electrode Potentials and Temperature Coefficients in Water at 298.15 K. *J. Phys. Chem. Ref. Data* **1989**, *18*, 1–21.

(14) Kumar, R.; Linden, A.; Nevado, C. Luminescent (N⁺C⁺C) Gold(III) Complexes: Stabilized Gold(III) Fluorides. *Angew. Chem., Int. Ed.* **2015**, *54*, 14287–14290.

(15) Roşca, D.-A.; Fernandez-Cestau, J.; Morris, J.; Wright, J. A.; Bochmann, M. Gold(III)-CO and gold(III)-CO₂ complexes and their role in the water-gas shift reaction. *Sci. Adv.* **2015**, *1*, No. e1500761.

(16) Kumar, R.; Krieger, J. P.; Gómez-Bengo, E.; Fox, T.; Linden, A.; Nevado, C. The First Gold(III) Formate: Evidence for β -Hydride Elimination. *Angew. Chem., Int. Ed.* **2017**, *56*, 12862–12865.

(17) Cui, J.-F.; Ko, H.-M.; Shing, K.-P.; Deng, J.-R.; Lai, N. C.-H.; Wong, M.-K. C,O-Chelated BINOL/Gold(III) Complexes: Synthesis and Catalysis with Tunable Product Profiles. *Angew. Chem., Int. Ed.* **2017**, *56*, 3074–3079.

(18) Holmsen, M. S. M.; Nova, A.; Balcells, D.; Langseth, E.; Oien-Odegaard, S.; Heyn, R. H.; Tilset, M.; Laurenczy, G. Mutation at Gold(III): A Mechanistic Study of a Catalytic Acetylene Functionalization via a Double Insertion Pathway. *ACS Catal.* **2017**, *7*, 5023–5034.

(19) Font, P.; Valdés, H.; Guisado-Barrios, G.; Ribas, X. Hemilabile MIC⁺N ligands allow oxidant-free Au(I)/Au(III) arylation-lactonization of γ -alkenoic acids. *Chem. Sci.* **2022**, *13*, 9351–9360.

(20) Gukathasan, S.; Parkin, S.; Awuah, S. G. Cyclometalated Gold(III) Complexes Bearing DACH Ligands. *Inorg. Chem.* **2019**, *58*, 9326–9340.

(21) Mertens, R. T.; Parkin, S.; Awuah, S. G. Cancer cell-selective modulation of mitochondrial respiration and metabolism by potent organogold(III) dithiocarbamates. *Chem. Sci.* **2020**, *11*, 10465–10482.

(22) Abás, E.; Bellés, A.; Rodríguez-Diéguez, A.; Laguna, M.; Grasa, L. Selective cytotoxicity of cyclometalated gold(III) complexes on Caco-2 cells is mediated by G2/M cell cycle arrest. *Metallomics* **2021**, *13* (7), mfab034.

(23) van der Westhuizen, D.; Slabber, C. A.; Fernandes, M. A.; Joubert, D. F.; Kleinhans, G.; van der Westhuizen, C. J.; Stander, A.

Munro, O. Q.; Bezuidenhout, D. I. A Cytotoxic Bis(1,2,3-triazol-5-ylidene)carbazolidine Gold(III) Complex Targets DNA by Partial Intercalation. *Chem. A Eur. J.* **2021**, *27*, 8295–8307.

(24) Luo, Y.; Cao, B.; Zhong, M.; Liu, M.; Xiong, X.; Zou, T. Organogold(III) Complexes Display Conditional Photoactivities: Evolving From Photodynamic into Photoactivated Chemotherapy in Response to O₂ Consumption for Robust Cancer Therapy. *Angew. Chem., Int. Ed.* **2022**, *61*, No. e202212689.

(25) Bachmann, M.; Fessler, R.; Blacque, O.; Venkatesan, K. Towards blue emitting monocyclometalated gold(III) complexes – synthesis, characterization and photophysical investigations. *Dalton Trans.* **2019**, *48*, 7320–7330.

(26) Beucher, H.; Kumar, S.; Merino, E.; Hu, W.-H.; Stemmler, G.; Cuesta-Galisteo, S.; González, J. A.; Jagielski, J.; Shih, C.-J.; Nevado, C. Highly Efficient Green Solution Processable Organic Light-Emitting Diodes Based on a Phosphorescent κ^3 -(N^{^C}^C^{^A})-Gold(III)-Alkynyl Complex. *Chem. Mater.* **2020**, *32*, 1605–1611.

(27) Wong, C.-Y.; Lai, S.-L.; Leung, M.-Y.; Tang, M.-C.; Li, L.-K.; Chan, M.-Y.; Yam, V. W.-W. Realization of Long Operational Lifetimes in Vacuum-Deposited Organic Light-Emitting Devices Based on para-Substituted Pyridine Carbazolylgold(III) C^{^C}^N Complexes. *J. Am. Chem. Soc.* **2023**, *145*, 2638–2646.

(28) Yang, J.; Giuso, V.; Hou, M.-C.; Remadna, E.; Forté, J.; Su, H.-C.; Gourlaouen, C.; Mauro, M.; Bertrand, B. Biphenyl Au(III) Complexes with Phosphine Ancillary Ligands: Synthesis, Optical Properties, and Electroluminescence in Light-Emitting Electrochemical Cells. *Inorg. Chem.* **2023**, *62*, 4903–4921.

(29) Constable, E. C.; Leese, T. A. Cycloaurated derivatives of 2-phenylpyridine. *J. Organomet. Chem.* **1989**, *363*, 419–424.

(30) Henderson, W. The Chemistry of Cyclometallated Gold(III) Complexes with C,N-Donor Ligands. In *Advances in Organometallic Chemistry*, West, R., Hill, A. F., Eds.; Academic Press, 2006; Vol. 54, pp 207–265.

(31) Shaw, A. P.; Tilset, M.; Heyn, R. H.; Jakobsen, S. Microwave methods for the synthesis of gold(III) complexes. *J. Coord. Chem.* **2011**, *64*, 38–47.

(32) Tlahuext-Aca, A.; Hopkinson, M. N.; Daniliuc, C. G.; Glorius, F. Oxidative Addition to Gold(I) by Photoredox Catalysis: Straightforward Access to Diverse (C,N)-Cyclometallated Gold(III) Complexes. *Chem. A Eur. J.* **2016**, *22*, 11587–11592.

(33) Serra, J.; Parella, T.; Ribas, X. Au(III)-aryl intermediates in oxidant-free C–N and C–O cross-coupling catalysis. *Chem. Sci.* **2017**, *8*, 946–952.

(34) Martín, J.; Gómez-Bengoa, E.; Genoux, A.; Nevado, C. Synthesis of Cyclometallated Gold(III) Complexes via Catalytic Rhodium to Gold(III) Transmetalation. *Angew. Chem., Int. Ed.* **2022**, *61*, No. e202116755.

(35) Kumar, R.; Linden, A.; Nevado, C. Luminescent (N^{^C}^C^{^A}) Gold(III) Complexes: Stabilized Gold(III) Fluorides. *Angew. Chem., Int. Ed.* **2015**, *54*, 14287–14290.

(36) Feuerstein, W.; Holzer, C.; Gui, X.; Neumeier, L.; Kloppe, W.; Breher, F. Synthesis of New Donor-Substituted Biphenyls: Pre-ligands for Highly Luminescent (C^{^C}^C^{^D}) Gold(III) Pincer Complexes. *Chem. A Eur. J.* **2020**, *26*, 17156–17164.

(37) Beucher, H.; Schörgenhumer, J.; Merino, E.; Nevado, C. Chelation-assisted C–C bond activation of biphenylene by gold(I) halides. *Chem. Sci.* **2021**, *12*, 15084–15089.

(38) Hylland, K. T.; Schmidtke, I. L.; Wragg, D. S.; Nova, A.; Tilset, M. Synthesis of substituted (N,C) and (N,C,C) Au(III) complexes: the influence of sterics and electronics on cyclometalation reactions. *Dalton Trans.* **2022**, *51*, 5082–5097.

(39) For the synthesis of (C^{^N}^C^{^A})gold(III) complexes, see refs 39–42. Wong, K.-H.; Cheung, K.-K.; Chan, M. C.-W.; Che, C.-M. Application of 2,6-Diphenylpyridine as a Tridentate [C^{^N}^C^{^A}] Dianionic Ligand in Organogold(III) Chemistry. Structural and Spectroscopic Properties of Mono- and Binuclear Transmetalated Gold(III) Complexes. *Organometallics* **1998**, *17*, 3505–3511.

(40) Fernandez-Cestau, J.; Bertrand, B.; Blaya, M.; Jones, G. A.; Penfold, T. J.; Bochmann, M. Synthesis and luminescence

modulation of pyrazine-based gold(III) pincer complexes. *Chem. Commun.* **2015**, *51*, 16629–16632.

(41) Eppel, D.; Rudolph, M.; Rominger, F.; Hashmi, A. S. K. Mercury-Free Synthesis of Pincer [C^{^N}^C^{^A}]Au^{III} Complexes by an Oxidative Addition/CH Activation Cascade. *ChemSusChem* **2020**, *13*, 1986–1990.

(42) McEllin, A. J.; Gault, C. A.; Whitwood, A. C.; Lynam, J. M.; Bruce, D. W. On the mercuration, palladation, transmetalation and direct auration of a C^{^N}^C pincer ligand. *Dalton Trans.* **2023**, *52*, 872–876.

(43) Vreeken, V.; Broere, D. L. J.; Jans, A. C. H.; Lankelma, M.; Reek, J. N. H.; Siegler, M. A.; van der Vlugt, J. I. Well-Defined Dinuclear Gold Complexes for Preorganization-Induced Selective Dual Gold Catalysis. *Angew. Chem., Int. Ed.* **2016**, *55*, 10042–10046.

(44) Navarro, M.; Toledo, A.; Joost, M.; Amgoune, A.; Mallet-Ladeira, S.; Bourissou, D. π Complexes of P^{^A}P and P^{^N}N chelated gold(I). *Chem. Commun.* **2019**, *55*, 7974–7977.

(45) Genoux, A.; Biedrzycki, M.; Merino, E.; Rivera-Chao, E.; Linden, A.; Nevado, C. Synthesis and Characterization of Bidentate (P^{^N})Gold(III) Fluoride Complexes: Reactivity Platforms for Reductive Elimination Studies. *Angew. Chem., Int. Ed.* **2021**, *60*, 4164–4168.

(46) Navarro, M.; Tabey, A.; Szalóki, G.; Mallet-Ladeira, S.; Bourissou, D. Stable Au(III) Complexes Bearing Hemilabile P^{^N} and C^{^N} Ligands: Coordination of the Pendant Nitrogen upon Oxidation of Gold. *Organometallics* **2021**, *40*, 1571–1576.

(47) Vesseur, D.; Li, S.; Mallet-Ladeira, S.; Miqueu, K.; Bourissou, D. Ligand-Enabled Oxidative Fluorination of Gold(I) and Light-Induced Aryl-F Coupling at Gold(III). *J. Am. Chem. Soc.* **2024**, *146*, 11352–11363.

(48) Messina, M. S.; Stauber, J. M.; Waddington, M. A.; Rheingold, A. L.; Maynard, H. D.; Spokoyny, A. M. Organometallic Gold(III) Reagents for Cysteine Arylation. *J. Am. Chem. Soc.* **2018**, *140*, 7065–7069.

(49) Rodriguez, J.; Zeineddine, A.; Sosa Carrizo, E. D.; Miqueu, K.; Saffon-Merceron, N.; Amgoune, A.; Bourissou, D. Catalytic Au(I)/Au(III) arylation with the hemilabile MeDalpos ligand: unusual selectivity for electron-rich iodoarenes and efficient application to indoles. *Chem. Sci.* **2019**, *10*, 7183–7192.

(50) Chintawar, C. C.; Yadav, A. K.; Patil, N. T. Gold-Catalyzed 1,2-Diarylation of Alkenes. *Angew. Chem., Int. Ed.* **2020**, *59*, 11808–11813.

(51) Rodriguez, J.; Tabey, A.; Mallet-Ladeira, S.; Bourissou, D. Oxidative additions of alkynyl/vinyl iodides to gold and gold-catalyzed vinylation reactions triggered by the MeDalpos ligand. *Chem. Sci.* **2021**, *12*, 7706–7712.

(52) Cadge, J. A.; Gates, P. J.; Bower, J. F.; Russell, C. A. Migratory Insertion of CO into a Au–C Bond. *J. Am. Chem. Soc.* **2022**, *144*, 19719–19725.

(53) Das, A.; Patil, N. T. Ligand-Enabled Gold-Catalyzed C(sp²)-O Cross-Coupling Reactions. *ACS Catal.* **2023**, *13*, 3847–3853.

(54) McCallum, T. Heart of gold: enabling ligands for oxidative addition of haloorganics in Au(I)/Au(III) catalyzed cross-coupling reactions. *Org. Biomol. Chem.* **2023**, *21*, 1629–1646.

(55) Huang, L.; Rudolph, M.; Rominger, F.; Hashmi, A. S. K. Photosensitizer-Free Visible-Light-Mediated Gold-Catalyzed 1,2-Difunctionalization of Alkynes. *Angew. Chem., Int. Ed.* **2016**, *55*, 4808–4813.

(56) Huang, L.; Rominger, F.; Rudolph, M.; Hashmi, A. S. K. A general access to organogold(III) complexes by oxidative addition of diazonium salts. *Chem. Commun.* **2016**, *52*, 6435–6438.

(57) Zeineddine, A.; Estévez, L.; Mallet-Ladeira, S.; Miqueu, K.; Amgoune, A.; Bourissou, D. Rational development of catalytic Au(I)/Au(III) arylation involving mild oxidative addition of aryl halides. *Nat. Commun.* **2017**, *8*, 565.

(58) Martín, J.; Schörgenhumer, J.; Biedrzycki, M.; Nevado, C. (P^{^N}^N^{^A}) Ligands to Stabilize Gold(III): A Straightforward Access to Hydroxo, Formate, and Hydride Complexes. *Inorg. Chem.* **2024**, *63*, 8390–8396.

(59) Martin, J.; Schörghamer, J.; Nevado, C. Interrogating the *anti*-Insertion of Alkynes into Gold(III). *JACS Au*. **2025**, *5*, 1439–1447.

(60) For more details, see the [Supporting Information](#).

(61) Beucher, H.; Kumar, S.; Kumar, R.; Merino, E.; Hu, W.-H.; Stemmler, G.; Cuesta-Galisteo, S.; González, J. A.; Bezinge, L.; Jagielski, J.; Shih, C.-J.; Nevado, C. Phosphorescent κ^3 -(N⁺C⁺C)-Gold(III) Complexes: Synthesis, Photophysics, Computational Studies and Application to Solution-Processable OLEDs. *Chem. A Eur. J.* **2020**, *26*, 17604–17612.

(62) Zehnder, T. N.; Blacque, O.; Venkatesan, K. Luminescent monocyclometalated cationic gold(III) complexes: synthesis, photophysical characterization and catalytic investigations. *Dalton Trans.* **2014**, *43*, 11959–11972.

(63) Leung, M.-Y.; Leung, S. Y.-L.; Yim, K.-C.; Chan, A. K.-W.; Ng, M.; Yam, V. W.-W. Multiresponsive Luminescent Cationic Cyclometalated Gold(III) Amphiphiles and Their Supramolecular Assembly. *J. Am. Chem. Soc.* **2019**, *141*, 19466–19478.