Q

Main page Contents Featured content Current events Random article Donate to Wikipedia Wikipedia store

The Free Encyclopedia

Interaction Help

About Wikipedia

Community portal Recent changes Contact page Tools What links here Related changes

Upload file Special pages Permanent link Page information Wikidata item Cite this page Print/export

Create a book Download as PDF Printable version

Languages Afrikaans

O

Alemannisch العربية Aragonés Asturianu Башкортса Беларуская Беларуская (тарашкевіца)

Български Bosanski Català Čeština ChiShona Cymraeg Dansk Deutsch Eesti Ελληνικά Español Esperanto Euskara فارسى Français Gàidhlig

Galego 赣語 한국어 Հայերեն हिन्दी Hrvatski Íslenska Italiano עברית Қазақша Kiswahili Kreyòl ayisyen Latina Latviešu Lëtzebuergesch Lietuvių Magyar മലയാളം Bahasa Melayu Nederlands 日本語 Norsk bokmål Norsk nynorsk Occitan

ਪੰਜਾਬੀ

ينجابى

Polski

Piemontèis

Português

Română Русский

Scots

Shqip

Simple English

Slovenčina Slovenščina

كوردبي ناوهندى

Suomi

Svenska

Српски / srpski Srpskohrvatski /

српскохрватски

Tagalog தமிழ் ใทย Türkçe Українська Tiếng Việt Winaray ייִדיש 粵語 中文 Edit links

Coordinate system From Wikipedia, the free encyclopedia

1.1 Number line

3 Transformations

5 Coordinate maps

7 See also

8 References

9 External links

Number line

1.2 Cartesian coordinate system

1.6 Other commonly used systems

7.1 Relativistic Coordinate Systems

Common coordinate systems

1.3 Polar coordinate system

2 Coordinates of geometric objects

4 Coordinate curves and surfaces

6 Orientation-based coordinates

Article Talk


This is an old revision of this page, as edited by KasparBot (talk | contribs) at 18:01, 16 May 2015 (embed authority control with wikidata information). The present address (URL) is a permanent link to this revision, which may differ significantly from the current revision.

(diff) ← Previous revision | Latest revision (diff) | Newer revision → (diff)

"Coordinate" redirects here. For other uses, see Coordinate (disambiguation).

In geometry, a coordinate system is a system which uses one or more numbers, or coordinates, to uniquely determine the position of a point or other geometric element on a manifold such as Euclidean space.[1][2] The order of the coordinates is significant and they are sometimes identified by their position in an ordered tuple and sometimes by a letter, as in "the x-coordinate". The coordinates are taken to be real numbers in elementary mathematics, but may be complex numbers or elements of a more abstract system such as a commutative ring.

The use of a coordinate system allows problems in geometry to be translated into problems about numbers and vice versa; this is the basis of analytic geometry.[3] Contents [hide] 1 Common coordinate systems



Search

Read Edit View history

1.4 Cylindrical and spherical coordinate systems 1.5 Homogeneous coordinate system

### The simplest example of a coordinate system is the identification of points on a line with real numbers using the number line. In this system, an arbitrary point O (the

Main article: Number line

### origin) is chosen on a given line. The coordinate of a point P is defined as the signed distance from O to P, where the signed distance is the distance taken as positive or

negative depending on which side of the line P lies. Each point is given a unique coordinate and each real number is the coordinate of a unique point.[4]


-7 -6 -5 -4 -3 -2 -1 0 9 Cartesian coordinate system

In three dimensions, three perpendicular planes are chosen and the three

coordinates of a point are the signed distances to each of the planes. [5] This

Main article: Cartesian coordinate system The prototypical example of a coordinate system is the Cartesian coordinate system. In the plane, two perpendicular lines are chosen and the coordinates of a point are taken to be the signed distances to the lines.

can be generalized to create n coordinates for any point in n-dimensional Euclidean space. Depending on the direction and order of the coordinate axis the system may




be a right-hand or a left-hand system.

a given angle θ, there is a single line through the pole whose angle with the polar axis is θ (measured counterclockwise from the axis to the line). Then there is a unique

point on this line whose signed distance from the origin is r for given number r. For a given pair of coordinates (r, θ) there is a single point, but any point is represented

by many pairs of coordinates. For example  $(r, \theta)$ ,  $(r, \theta+2\pi)$  and  $(-r, \theta+\pi)$  are all polar coordinates for the same point. The pole is represented by  $(0, \theta)$  for any value of  $\theta$ .



(2,3)

#### Cylindrical and spherical coordinate systems Main articles: Cylindrical coordinate system and Spherical coordinate system

coordinates (r, z) to polar coordinates  $(\rho, \phi)$  giving a triple  $(\rho, \theta, \phi)$ . [8] Cylindrical coordinates system have the following coordinates,  $\rho, \phi, z$ Homogeneous coordinate system

Main article: Homogeneous coordinates A point in the plane may be represented in homogeneous coordinates by a triple (x, y, z) where x/z and y/z are the Cartesian coordinates of the point. [9] This introduces an "extra" coordinate since only two are needed to specify a point on the plane, but this system is useful in that it represents any point on the projective plane without the

use of infinity. In general, a homogeneous coordinate system is one where only the ratios of the coordinates are significant and not the actual values.

There are two common methods for extending the polar coordinate system to three dimensions. In the cylindrical coordinate system, a z-coordinate with the same

meaning as in Cartesian coordinates is added to the r and  $\theta$  polar coordinates. [7] Spherical coordinates take this a step further by converting the pair of cylindrical

# Orthogonal coordinates: coordinate surfaces meet at right angles

orthogonal coordinates

Other commonly used systems

The log-polar coordinate system represents a point in the plane by the logarithm of the distance from the origin and an angle measured from a reference line

Some other common coordinate systems are the following:

- intersecting the origin. Plücker coordinates are a way of representing lines in 3D Euclidean space using a six-tuple of numbers as homogeneous coordinates.
- Canonical coordinates are used in the Hamiltonian treatment of mechanics.
- Trilinear coordinates are used in the context of triangles.
- The Whewell equation relates arc length and the tangential angle.

Parallel coordinates visualise a point in n-dimensional space as a polyline connecting points on n vertical lines.

 The Cesàro equation relates arc length and curvature. Coordinates of geometric objects

Barycentric coordinates as used for ternary plots and more generally in the analysis of triangles.

used to distinguish the type of coordinate system, for example the term line coordinates is used for any coordinate system that specifies the position of a line.

## **Transformations** See also: Active and passive transformation

those for the coordinate transformation)

principle of duality.[11]

are described by coordinate transformations which give formulas for the coordinates in one system in terms of the coordinates in another system. For example, in the plane, if Cartesian coordinates (x, y) and polar coordinates  $(r, \theta)$  have the same origin, and the polar axis is the positive x axis, then the coordinate transformation from polar to Cartesian coordinates is given by  $x = r \cos\theta$  and  $y = r \sin\theta$ . With every bijection from the space to itself two coordinate transformations can be associated: such that the new coordinates of the image of each point are the same as the old coordinates of the original point (the formulas for the mapping are the inverse of

Because there are often many different possible coordinate systems for describing geometrical figures, it is important to understand how they are related. Such relations

It may occur that systems of coordinates for two different sets of geometric figures are equivalent in terms of their analysis. An example of this is the systems of

homogeneous coordinates for points and lines in the projective plane. The two systems in a case like this are said to be dualistic. Dualistic systems have the property

that results from one system can be carried over to the other since these results are only different interpretations of the same analytical result; this is known as the


Coordinate curves and surfaces In two dimensions if all but one coordinate in a point coordinate system is held constant and the remaining coordinate is allowed to vary, then the resulting curve is called a **coordinate curve** (some authors use the phrase "coordinate line"). This procedure does not

surface is called a **coordinate surface**. For example the coordinate surfaces obtained by holding ρ constant in the spherical coordinate system are the spheres with center at the origin. In three-dimensional space the intersection of two coordinate surfaces is a coordinate curve. Coordinate hypersurfaces are defined similarly in higher dimensions. [13] Coordinate maps Main article: Manifold The concept of a coordinate map, or chart is central to the theory of manifolds. A coordinate map is essentially a coordinate system for a subset of a given space with

always make sense, for example there are no coordinate curves in a homogeneous coordinate system. In the Cartesian coordinate

system the coordinate curves are, in fact, straight lines. Specifically, they are the lines parallel to one of the coordinate axes. For

other coordinate systems the coordinates curves may be general curves. For example the coordinate curves in polar coordinates



Coordinate surfaces in the Spherical coordinate system

# vectors aligned with those axes.

Absolute angular momentum

See also

 Alpha-numeric grid Analytic geometry

Fractional coordinates

Galilean transformation

Frame of reference

- Nomogram, graphical representations of different coordinate systems Relativistic Coordinate Systems
- Gullstrand

  Painlevé coordinates Isotropic coordinates Kruskal–Szekeres coordinates

Schwarzschild coordinates

- V•T•E Two dimensional
- Bispherical coordinates Bipolar cylindrical coordinates Conical coordinates 6-sphere coordinates Flat-ring cyclide coordinates Flat-disk cyclide coordinates Three dimensional Bi-cyclide coordinates · Cap-cyclide coordinates · Concave bi-sinusoidal single-centered coordinates · Concave bi-sinusoidal double-centered coordinates · Convex

References

Moods p. 1

Weisstein, Eric W., "Coordinate System" , MathWorld. Stewart, James B.; Redlin, Lothar; Watson, Saleem (2008). College Algebra (5th ed.). Brooks Cole. pp. 13–19. ISBN 0-495-56521-0. 5. A Moon P, Spencer DE (1988). "Rectangular Coordinates (x, y, z)". Field Theory Handbook, Including Coordinate Systems, Differential Equations, and Their Solutions (corrected

- ISBN 0-201-55478-X. 7. A Margenau, Henry; Murphy, George M. (1956). The Mathematics of Physics and Chemistry. New York City: D. van Nostrand. p. 178. ISBN 9780882754239. LCCN 55010911 . OCLC 3017486 €.
- A Hodge, W. V. D.; D. Pedoe (1994) [1947]. Methods of Algebraic Geometry, Volume I (Book II). Cambridge University Press. ISBN 978-0-521-46900-5. 11. ^ Woods p. 2
- A Tang, K. T. (2006). Mathematical Methods for Engineers and Scientists 2. Springer. p. 13. ISBN 3-540-30268-9. 13. Liseikin, Vladimir D. (2007). A Computational Differential Geometry Approach to Grid Generation. Springer. p. 38. ISBN 3-540-34235-4. Munkres, James R. (2000) Topology. Prentice Hall. ISBN 0-13-181629-2.

2nd, 3rd print ed.). New York: Springer-Verlag. pp. 9–11 (Table 1.01). ISBN 978-0-387-18430-2.

- 15. A Hanspeter Schaub, John L. Junkins (2003). "Rigid body kinematics". Analytical Mechanics of Space Systems . American Institute of Aeronautics and Astronautics. p. 71. ISBN 1-56347-563-4.
- Shigeyuki Morita, Teruko Nagase, Katsumi Nomizu (2001). Geometry of Differential Forms 

  AMS Bookstore. p. 12. ISBN 0-8218-1045-6.

Voitsekhovskii, M.I.; Ivanov, A.B. (2001), "Coordinates" , in Hazewinkel, Michiel, Encyclopedia of Mathematics, Springer, ISBN 978-1-55608-010-4

Orthogonal coordinate systems

6. Finney, Ross; George Thomas; Franklin Demana; Bert Waits (June 1994). Calculus: Graphical, Numerical, Algebraic (Single Variable Version ed.). Addison-Wesley Publishing Co.

- External links



Wikimedia Commons has media related to Coordinate systems.

Powered By MediaWiki

[hide]

This version of the page has been revised. Besides normal editing, the reason for revision may have been that this version contains factual inaccuracies, vandalism, or material not compatible with the Creative Commons Attribution-ShareAlike License. Privacy policy About Wikipedia Disclaimers Contact Wikipedia Developers Mobile view

Categories: Coordinate systems | Analytic geometry

WIKIMEDIA

 $\Box$ 

Curvilinear coordinates are a generalization of coordinate systems generally; the system is based on the intersection of curves.

Skew coordinates: coordinate surfaces are not orthogonal

- Generalized coordinates are used in the Lagrangian treatment of mechanics.
- There are ways of describing curves without coordinates, using intrinsic equations that use invariant quantities such as curvature and arc length. These include:
- Coordinates systems are often used to specify the position of a point, but they may also be used to specify the position of more complex figures such as lines, planes, circles or spheres. For example Plücker coordinates are used to determine the position of a line in space.[10] When there is a need, the type of figure being described is
  - Main article: List of common coordinate transformations

 such that the old coordinates of the image of each point are the same as the new coordinates of the original point (the formulas for the mapping are the same as those for the coordinate transformation) For example, in 1D, if the mapping is a translation of 3 to the right, the first moves the origin from 0 to 3, so that the coordinate of each point becomes 3 less, while the second moves the origin from 0 to -3, so that the coordinate of each point becomes 3 more.

obtained by holding r constant are the circles with center at the origin. Coordinates systems for Euclidean space other than the Cartesian coordinate system are called curvilinear coordinate systems.[12] In three-dimensional space, if one coordinate is held constant and the remaining coordinates are allowed to vary, then the resulting

the property that each point has exactly one set of coordinates. More precisely, a coordinate map is a homeomorphism from an open subset of a space X to an open subset of R<sup>n</sup>.[14] It is often not possible to provide one consistent coordinate system for an entire space. In this case, a collection of coordinate maps are put together to form an atlas covering the space. A space equipped with such an atlas is called a manifold and additional structure can be defined on a manifold if the structure is consistent where the coordinate maps overlap. For example a differentiable manifold is a manifold where the change of coordinates from one coordinate map to another is always a differentiable function.

#### Astronomical coordinate systems Axes conventions in engineering Coordinate-free

Orientation-based coordinates

 Eddington–Finkelstein coordinates Gaussian polar coordinates

Geographic coordinate system

- Cartesian coordinate system · Polar coordinate system · Parabolic coordinate system · Bipolar coordinates · Elliptic coordinates Cartesian coordinate system · Cylindrical coordinate system · Spherical coordinate system · Parabolic cylindrical coordinates · Paraboloidal coordinates · Oblate spheroidal coordinates · Prolate spheroidal coordinates · Ellipsoidal coordinates · Elliptic cylindrical coordinates · Toroidal coordinates ·
- inverted-sinusoidal spherically aligned coordinates · Quasi-random-intersection cartesian coordinates
- Norse PM, Feshbach H (1953). Methods of Theoretical Physics, Part I. New York: McGraw-Hill. p. 658. ISBN 0-07-043316-X. LCCN 52011515 ₽. 9. A Jones, Alfred Clement (1912). An Introduction to Algebraical Geometry. Clarendon.
- Hexagonal Coordinate System
- Authority control GND: 4165251-4 €