

# Analysis of the COVID-19 Case Rate and Excess Mortality Rate by Pandemic Wave

Yiqiao Zhu<sup>1</sup>, Shuoyuan Gao<sup>1</sup>

<sup>1</sup> Department of Biostatistics at the University of Michigan School of Public Health

---

## Abstract

COVID-19 pandemics are organized into multiple waves, each with temporally and geographically variable transmission and mortality. Understanding these dynamics is critical for assessing public health responses and improving future preparedness. We obtained weekly state-level data on COVID-19 cases, deaths, and excess mortality from January 2020 to April 2025. Population-standardized rates were calculated and used to identify three major waves. Mortality trends were modeled, and the performance of forecasting models was evaluated across waves. Wave 1 showed high excess mortality and the highest case fatality rate (CFR), particularly in the Northeast, reflecting limited early testing. Wave 2 was the deadliest period nationally, with widespread regional impact. Wave 3 presented a mixed pattern, with some states showing improved outcomes, likely due to vaccination and enhanced preparedness. CFR declined across waves, and nonlinear models outperformed linear models in predicting excess mortality. These findings underscore the need for adaptive modeling strategies and proactive public health interventions in managing future pandemics.

**Index Terms:** COVID-19, Excess Mortality, Case Fatality Rate, Mortality Mode

---

## 1 Introduction

This report is organized in a left-to-right reading direction, consistent with standard English-language academic formatting.<sup>1</sup>

The COVID-19 pandemic has had a significant global impact over the last five years. COVID-19 was first discovered in Wuhan, China, in late 2019, and it quickly became a global crisis. By early 2020, cases had been reported on nearly every continent, prompting the World Health Organization to officially declare the pandemic. As infections and deaths exceeded expectations, societies were disrupted on multiple levels, including overburdened health-care systems, severe economic downturns, and widespread psychological stress.

The first confirmed case was reported on January 21, 2020, in the US state of Washington. In just three months, the United States became the epicenter of the global epidemic, with a sharp rise in cumulative cases and deaths. The pandemic's pattern in the United States has been marked by multiple waves of infection, each influenced by regional factors, public health responses, and the emergence of new variants. These waves also represent changes in social behavior, vaccination rates, and health-care capacity. As such, they provide a useful framework for analyzing pandemic progress and the effectiveness of various mitigation strategies.

COVID-19 Pandemics occur in distinct temporal waves, each with different levels of virus transmission, severity, and public health response. Understanding these temporal patterns is critical to evaluating the pandemic's progression and the effectiveness of mitigation strategies. Splitting the pandemic period into well-defined waves creates a framework for analyzing changes in population-level outcomes such as viral virulence, health-care system strain, and excess mortality.

A central challenge in this context lies in the generalizability of predictive models across different waves. Models trained on data from one phase of the pandemic may not perform reliably when applied to other periods, due to shifts in dominant viral variants,

vaccination rates, public health interventions, and healthcare system capacity. These evolving dynamics introduce substantial heterogeneity in observed mortality outcomes and complicate efforts to build robust forecasting models.

The purpose of this study was to analyze Analysis of the COVID-19 Cases Rate and the Excess Mortality Rate by Pandemic Wave. A data-driven approach was utilized to divide the U.S. pandemic timeline (April 2020 to July 2024) into three distinct waves. For the data, a macro-measurement is performed, and response data are presented to support our waves. Analysis relying on these waves summarizes state-to-state differences. Reveal how geographic differences affect outbreak outcomes. Finally, quantitatively assess the generalizability of statistical models trained on one wave to other waves in predicting excess mortality. We used linear regression, localized polynomial regression (LOESS), and spline-based methods to capture temporal variation in COVID-19 case rates and mortality outcomes and to assess the robustness of these models across pandemic phases.

### 1.1 Report Organisation

The remainder of this paper is organized as follows: Section 2 describes the data sources and methods. Section 3 presents the results of mortality trends and model comparisons. Section 4 discusses the findings, implications, and limitations.

## 2 Method

This study integrated data from three primary sources to construct a comprehensive panel of weekly state-level population, COVID-19 case, and death outcomes.

Annual population estimates were obtained from the U.S. Census Bureau's *National and State Population Estimates 2024* dataset [U.S. Census Bureau \(2024\)](#), which provides data for all 50 states and the District of Columbia, and Puerto Rico from April 1, 2020, through July 1, 2024. Weekly counts of COVID-19 confirmed cases were obtained from the State Case Surveillance Dataset, as provided by [Centers for Disease Control and Prevention \(2024b\)](#). Death records, including the percentage of total deaths, COVID-19

---

<sup>1</sup> This document follows a two-column format. Please read each page from top to bottom of the left column, then continue at the top of the right column.

attributable deaths, and expected deaths, were obtained from the CDC ([Centers for Disease Control and Prevention, 2024a](#)). All data were aligned by state and pandemic week based on the MMWR calendar. The case count variable was converted to an integer format and standardized as cases. The year variable in the mortality dataset was also harmonized to accommodate cross-year formats (e.g., “2021/2022”), and all numeric columns were explicitly converted to ensure consistency across datasets.

Two key metrics were derived from mortality records:

$$\text{Excess Deaths} = \text{Total Deaths} - \left( \frac{\text{Total Deaths}}{\frac{\text{Percent of Expected Deaths}}{100}} \right) \quad (1)$$

$$\text{Excess Mortality Rate} = \text{Percent of Expected Deaths} - 100 \quad (2)$$

To construct a unified dataset, we first generated a full grid of weekly dates and U.S. states using a Cartesian join. This allowed us to merge population estimates, COVID-19 cases, and mortality data using consistent identifiers—state, epidemiological week, and year. Observations with missing values in key metrics were imputed with zeros to ensure temporal completeness. The resulting dataset includes one row per state-week combination, with fully aligned demographic, case, and mortality information.

### 2.1: Defining Pandemic Waves Using Data Visualization

To delineate pandemic waves, we calculated weekly COVID-19 case rates, death rates, and excess mortality rates per 100,000 population from January 2020 to April 2025, enabling standardized comparisons across states. These metrics were visualized over time and stratified by region. Wave boundaries were identified based on visual inflection points where all three indicators showed sustained peaks across multiple regions, marked by shaded areas in the time series plots.

### 2.2: State-Level COVID-19 Mortality and Excess Mortality Rate by Wave

**Quantile-Based Classification and Heatmap Visualization.** To analyze geographic patterns in COVID-19 mortality, we calculated the cumulative COVID-19 death rate and excess mortality per 100,000 population for each U.S. state within each pandemic wave. These values were summed across all weeks within each wave. To facilitate interpretation, we categorized states into Low, Medium, and High mortality groups based on the 33rd and 66th percentiles. These groupings were visualized via two heatmaps to highlight temporal and spatial mortality trends.

**Ranking States by Mortality Extremes.** To underscore disparities, we ranked states by total COVID-19 death rate and excess mortality within each wave. The top three and bottom three states were identified and visualized using a faceted bar chart, enabling direct wave-to-wave comparisons.

### 2.3: Virulence Across Waves

To estimate clinical severity across pandemic waves, we calculated the Case Fatality Rate (CFR), defined as:

$$\text{CFR} = \left( \frac{\text{Total Death Rate}}{\text{Total Case Rate}} \right) \times 100$$

Weekly case and death rates (per 100,000) were aggregated across all states within each wave. We excluded observations with missing or zero case rates.

### 2.4: Modeling the Relationship Between Case Rate and Excess Mortality

To evaluate the relationship between COVID-19 case rates and excess mortality rates, we filtered weekly state-level data to include only the relevant metrics and pivoted the dataset to a wide format. We first fitted linear models per wave and visualized trends with scatter plots and regression lines.

We then trained models on one wave and tested on another to evaluate generalizability using RMSE and  $R^2$ . Finally, we compared model fit using linear regression (Equation 3), LOESS smoothing (Equation 4), and cubic splines (Equation 5), particularly when predicting Wave 3 from the Wave 2 model. Additional comparison between models trained on Waves 1 and 3, applied to Wave 2 data, revealed differences in distribution and model behavior across phases.

$$\text{Excess Mortality Rate} = \beta_0 + \beta_1 \cdot \text{Cases Rate} \quad (3)$$

$$\hat{y}_{\text{LOESS}}(x_0) = \sum_{i=1}^n w_i(x_0) \cdot y_i \quad (4)$$

$$\hat{y}_{\text{spline}} = \beta_0 + \sum_{j=1}^5 \beta_j B_j(x) \quad (5)$$

where  $w_i(x_0)$  are kernel-based weights around  $x_0$ , and  $B_j(x)$  are spline basis functions constructed from the case rate using 6 degrees of freedom.

## 3 Results


### 3.1: Defining Pandemic Waves Using Data Visualization

Based on Figure 1 below, in case rates, death rates, and excess mortality rates, we identified three major pandemic waves. These waves were defined using visual inflection points where all three indicators exhibited sustained peaks across multiple U.S. regions. The resulting wave periods are illustrated using shaded grey bands in the time series plots.

- **Wave 1:** March 1 – June 30, 2020
- **Wave 2:** October 1, 2020 – February 28, 2021
- **Wave 3:** July 1 – October 31, 2021

These intervals were chosen based on synchronized national surges across all three health metrics—case rates, death rates, and excess mortality. A detailed summary of each wave is as follows:

- **Wave 1 (March–June 2020):** A sharp rise in death rates occurred in early 2020, particularly in Regions 1–3. Although case rates were relatively low—likely due to limited testing capacity during the early phase—excess mortality spiked, confirming the severity of the initial wave.



**Figure 1.** COVID-19 Case and Death Rates by Region. Shaded areas indicate pandemic waves. Region labels correspond to U.S. Census-defined groupings: 1 = New England, 2 = New York and New Jersey, Puerto Rico, Virgin Islands, 3 = Mid-Atlantic, 4 = Southeast, 5 = Midwest, 6 = South Central, 7 = Central Plains, 8 = Mountain States, 9 = Pacific, 10 = Pacific Northwest.

- **Wave 2 (October 2020–February 2021):** Both case and death rates rose dramatically, peaking around January 2021. Excess mortality reached its highest level during this period, with all three indicators exhibiting synchronized spikes across all regions. This wave represented the most intense national health burden of the pandemic.
- **Wave 3 (July–October 2021):** A resurgence in case rates occurred during summer 2021, peaking between August and September. While death rates increased less sharply than in Wave 2, excess mortality still rose, marking a distinct third wave across most regions.


### 3.2: State-Level COVID-19 Mortality and Excess Mortality by Wave

Figure 2 visualizes state-level COVID-19 death rates and excess mortality rates per 100,000 population across the three defined pandemic waves. States are categorized into low (blue), medium (gold), and high (red) mortality groups based on quantile classification of total death rates within each wave.

During **Wave 1**, mortality levels varied substantially across states, with the majority falling into the low or medium categories.

In **Wave 2**, a notable shift occurred, with many states transitioning into the high mortality group, reflecting the widespread and intense health burden experienced nationwide during this period.

By **Wave 3**, the pattern became more heterogeneous. While several states continued to experience elevated death rates and excess mortality, others demonstrated improvement and shifted into lower mortality categories, indicating regional variation in pandemic dynamics and response.




**Figure 2.** State-level COVID-19 death rates (top) and excess mortality rates (bottom) per 100,000 population across three pandemic waves. States are grouped into low (blue), medium (gold), and high (red) severity based on wave-specific quantiles.

**Wave 1:** Mortality was concentrated in the Northeast, with New Jersey, Connecticut, and Massachusetts reporting the highest death rates. In contrast, Alaska, Hawaii, and Montana reported zero COVID-19 deaths and minimal excess mortality, reflecting the limited early spread in these regions.

**Wave 2:** This was the deadliest period nationally. States such as South Dakota and Arizona exhibited the highest excess mortality, indicating widespread transmission across central regions.

**Wave 3:** Southern and remote states—particularly Florida and Alaska—emerged as mortality hotspots. Meanwhile, states that were heavily affected in Wave 1, such as New Jersey and Massachusetts, experienced among the lowest mortality rates, highlighting regional shifts in pandemic impact.



**Figure 3.** Top and bottom three U.S. states by COVID-19 death rate (top) and excess mortality rate (bottom) per 100,000 population across pandemic waves. States are grouped by wave, with bar color indicating wave membership.

### 3.3: Virulence Across Waves

Figure 4 presents the Case Fatality Rate (CFR) for each of the three pandemic waves.

**Wave 1** exhibited the highest CFR at 4.5%, likely reflecting limited testing availability and clinical uncertainty during the early months of the pandemic.

In **Wave 2**, the CFR dropped significantly to 1.41%, followed by a further decline to 1.22% in **Wave 3**. This downward trend suggests reduced disease severity over time. Contributing factors likely include widespread vaccine roll-out, improved clinical management, and rising population-level immunity.




Figure 4. Case Fatality Rate (CFR) by COVID-19 wave.

### 3.4: Analyzing and Predicting the Relationship Between Case Rate and Excess Mortality Rate

Figure 5 illustrates how different models, trained on Wave 2 data, perform in predicting excess mortality during Wave 3. The goal is to evaluate how well case rates explain variation in excess mortality under different modeling approaches.

- **Observed data (yellow dots):** Actual excess mortality rates plotted against case rates for Wave 3.
- **Linear model (dashed cyan line):** A simple linear regression fitted on Wave 2 data.
- **LOESS model (blue line):** A non-parametric local regression approach capturing smooth trends.
- **Cubic spline model (red line):** A flexible spline regression with six degrees of freedom to model nonlinear patterns.

These models highlight the limitations of purely linear approaches and demonstrate the value of flexible modeling (e.g., LOESS, splines) in capturing pandemic dynamics over time.

#### Observations:

- The linear model tends to over-predict excess mortality when the case rate exceeds 700. This is due to the actual data flattening and trending downward in the upper range of case rates.




Figure 5. Model predictions of excess mortality in Wave 3 using models trained on Wave 2 data.

- Both LOESS and spline models more effectively capture the nonlinear relationship between case rate and excess mortality in Wave 3, particularly in the mid-to-high case rate range.
- Among the three approaches, the spline model produces the smoothest curve and demonstrates the best fit in the upper range of case rates, suggesting its strength in modeling subtle nonlinear patterns.

### 3.5: Model Performance Summary, Table 1

- The model trained on **Wave 2** data performs well in predicting excess mortality in **Wave 3** ( $RMSE = 16.40$ ,  $R^2 = 0.558$ ), indicating that the underlying data structures of these two waves are closely aligned.
- In contrast, the **Wave 1** model performs poorly when applied to either Wave 2 ( $RMSE = 111.21$ ) or Wave 3 ( $RMSE = 78.68$ ), suggesting that the mortality risk structure during the early phase of the pandemic differs substantially from later waves.
- As expected, the **Wave 3** model achieves the best fit on its own data ( $RMSE = 13.70$ ), and can be considered a near-benchmark for within-wave predictive accuracy.

Table 1

Cross-wave prediction performance for linear models. Each entry shows  $RMSE$  and  $R^2$  for a model trained on one wave and tested on another.

Trained on	Tested on	RMSE	$R^2$
Wave 1	Wave 1	18.64	0.479
Wave 1	Wave 2	111.21	0.450
Wave 1	Wave 3	78.68	0.558
Wave 2	Wave 1	26.11	0.479
Wave 2	Wave 2	15.06	0.450
Wave 2	Wave 3	16.40	0.558
Wave 3	Wave 1	24.03	0.479
Wave 3	Wave 2	18.50	0.450
Wave 3	Wave 3	13.70	0.558

## 4 Discussion

This study aimed to characterize the relationship between COVID-19 case rates and excess mortality across distinct

pandemic waves in the United States, and to evaluate the generalizability of predictive models across these waves.

### **Findings**

The first wave highlights the challenges of the early pandemic, where underreporting due to limited testing led to an apparent mismatch between case and death rates. Despite relatively low case counts, excess mortality reveals that the impact was more severe than reported case data alone would suggest.

The second wave reflects the compounded effect of widespread transmission, colder weather, and delayed public health responses. The simultaneous spikes across all three metrics underscore its severity and national reach.

The third wave, while still significant, occurred in a context of greater preparedness. The comparatively lower case fatality rate and somewhat contained mortality may reflect the protective effects of vaccination, improved treatment protocols, and greater public awareness. The persistent rise in excess mortality, however, suggests ongoing vulnerabilities in certain regions or populations.

The variability in Wave 1 likely reflects early differences in exposure, public health response, and testing capacity during the initial stage of the pandemic. The large increase in high-mortality states during Wave 2 suggests this was the deadliest wave, with widespread severe outcomes across multiple regions. The more mixed pattern in Wave 3 may reflect better preparedness, improved treatment protocols, and the growing impact of vaccination efforts.

These patterns underscore the shifting geographic burden of the pandemic over time. The early concentration of mortality in the Northeast gave way to central and southern regions in later waves. The decline in mortality rates in states like New Jersey and Massachusetts by Wave 3 likely reflects improved public health responses, accumulated experience, and broader population-level interventions.

### **Mortality Patterns and Model Performance**

One of the most salient findings is the clear heterogeneity in both mortality burden and model performance across pandemic waves.

Wave 1 exhibited the highest case fatality rate (CFR), suggesting severe clinical outcomes among detected cases. However, this elevated CFR likely reflects the limited testing capacity and healthcare preparedness in early 2020, where mild or asymptomatic cases went undetected, inflating the apparent severity.

In contrast, Waves 2 and 3 showed markedly lower CFRs, consistent with expanded testing, improved treatment protocols, and the initial impact of vaccination campaigns. These trends indicate a reduction in clinical virulence and demonstrate the adaptability of healthcare systems over time.

Geographic disparities in mortality outcomes were also

evident. The Northeast bore the brunt of Wave 1, while central and southern states experienced greater mortality during later waves. Notably, some states that were severely impacted early on, such as New Jersey and Massachusetts, demonstrated significantly improved outcomes in Wave 3. This pattern likely reflects the benefits of accumulated public health experience, stronger mitigation measures, and targeted vaccination efforts.

### **Limitations**

The cross-wave predictive analysis revealed important limitations in the temporal stability of statistical models.

Models trained on Wave 1 data performed poorly when applied to subsequent waves, particularly Wave 2, which had distinct epidemiological characteristics. This suggests that early pandemic conditions were fundamentally different from later phases in terms of virus spread, healthcare response, and population behavior.

Although approaches such as LOESS and cubic splines provided improved fits in scenarios where linear models failed to capture complex trends—particularly in Wave 3—the best-performing predictions were always obtained from models trained and tested on the same wave. This underscores the limited generalizability of pandemic forecasting models across distinct time periods.

Only three variables—COVID-19 cases, deaths, and the excess mortality rate—were considered. Important factors such as vaccination coverage and booster rates, which are likely to influence later trends, were not included. Additionally, time was not explicitly modeled as a variable in the fit.

The excess mortality rate, while valuable for capturing hidden pandemic impacts, may also be influenced by indirect effects such as healthcare disruptions, deferred treatment for non-COVID conditions, or population shifts due to migration.

### **Future Work**

Future work should incorporate a time variable into the prediction model to better capture longitudinal effects.

To better understand excess mortality trends in later waves, future models should introduce vaccine-related variables. Comparing the excess mortality rates between Wave 1 and later waves (Wave 2 and 3) could help quantify the protective effect of vaccination at the state level.

Additionally, incorporating booster coverage, hospitalization data, and indicators of healthcare capacity could further improve model accuracy and interpretability.

### **Acknowledgements**

This project report was conducted as part of the BIOSTAT620: Introduction to Health Data Science course offered by the Department of Biostatistics at the University of Michigan School of Public Health. We would like to thank

Professor Dylan Cable and our Graduate Student Instructor (GSI), Yize Hao, for their instruction and guidance throughout the semester.

## **References**

Centers for Disease Control and Prevention (2024a). *Provisional COVID-19 Death Counts by Week Ending Date and State*. Accessed April 2025. URL: <https://data.cdc.gov/National-Center-for-Health-Statistics/Provisional-COVID-19-Death-Counts-by-Week-Ending-D/r8kw-7aab>.

— (2024b). *Weekly United States COVID-19 Cases and Deaths by State*. Accessed April 2025. URL: <https://data.cdc.gov/Case-Surveillance/Weekly-United-States-COVID-19-Cases-and-Deaths-by-pwn4-m3yp>.

U.S. Census Bureau (2024). *National and State Population Estimates 2024*. Accessed April 2025. URL: <https://www.census.gov/newsroom/press-kits/2024/national-state-population-estimates.html>.