

**A POSTERIORI ERROR ESTIMATES FOR GENERAL
NUMERICAL METHODS FOR SCALAR CONSERVATION LAWS**

By

Bernardo Cockburn

and

Huiing Gau

IMA Preprint Series # 1261

October 1994

**INSTITUTE FOR MATHEMATICS AND ITS APPLICATIONS
UNIVERSITY OF MINNESOTA**

**514 Vincent Hall
206 Church Street S.E.
Minneapolis, Minnesota 55455**

A POSTERIORI ERROR ESTIMATES FOR GENERAL NUMERICAL METHODS FOR SCALAR CONSERVATION LAWS.

BERNARDO COCKBURN* AND HUIING GAU†

Abstract. In this paper, we obtain a simple *a posteriori* error estimate for numerical methods for nonlinear scalar conservation laws. The estimate is totally independent (i) of the dimension of the space, (ii) of the type of nonlinearity f , and (iii) of the numerical method. It can thus be used, regardless of the numerical schemes used to compute the approximate solution, to define mathematically sound adaptivity algorithms for conservation laws. Preliminary numerical results displaying the sharpness of the estimate are presented.

Key words. A posteriori error estimates, scalar conservation laws

AMS(MOS) subject classifications. 65M60, 65N30, 35L65

1. Introduction. In this paper, we show how a straightforward use of Kuznetsov approximation theory [12] leads to a simple *a posteriori* error estimate for numerical methods for the scalar conservation law [11]

$$v_t + \nabla \cdot f(v) = 0 \quad \text{in } (0, T) \times \mathbb{R}^d, \quad (1.1a)$$

$$v(0) = v_0 \quad \text{on } \mathbb{R}^d. \quad (1.1b)$$

The main advantages of the *a posteriori* estimate we obtain are that its form is totally independent (i) of the dimension of the space, (ii) of the type of nonlinearity f , and (iii) of the numerical method used to produce approximations to the entropy solution v of (1.1); this is the single *a posteriori* error estimate with the three above properties in the available literature. Hence, it can be used, regardless of the numerical scheme, to define mathematically sound adaptivity algorithms for conservation laws.

To understand why such a straightforward application of the work of Kuznetsov [12] took almost twenty years to come to light, we briefly display the history of the ideas leading to the estimate presented in this paper; for a more elaborated exposition, see the recent work by Cockburn and Gremaud [9].

Kuznetsov [12] was the first to obtain error estimates for numerical schemes for scalar conservation. He proved that monotone schemes defined in uniform Cartesian grids converge to the entropy solution in $L^\infty(0, T; L^1(\mathbb{R}^d))$ at the optimal rate of $(\Delta x)^{1/2}$. To do so, he first obtained an upper bound for the error that depends on the approximate solution (that is, he obtained an *a posteriori* error estimate) and then used the regularity properties of the approximate solution to obtain a rate of convergence. Since almost every author interested in obtaining error estimates for scalar conservation laws (Sanders [18], Lucier [14, 15, 16], Cockburn [3, 4, 5], Cockburn, LeFloch and Coquel [7], Cockburn and Gremaud [8], and Vila [20]) followed Kuznetsov's original approach, they all obtained an *a posteriori* error estimate and then, in order to get a rate of convergence, they all were bound to obtain regularity properties of the approximate solution – a very hard task indeed for even the simplest schemes in general triangulations.

* School of Mathematics, University of Minnesota, 127 Vincent Hall, Minneapolis, MN 55455. Partially supported by the University of Minnesota Supercomputer Institute.

† School of Mathematics, University of Minnesota, 127 Vincent Hall, Minneapolis, MN 55455.

Only recently, Cockburn and Gremaud [9] found a way to suitably modify Kuznetsov's original approach to obtain *a priori* error estimates for conservation laws, that is, error estimates which do not depend on the regularity properties of the approximate solution – something that was thought to be not possible until now. As a by product, it became more natural to view Kuznetsov's approach as an *a posteriori* error estimates approach that should be used for defining adaptivity strategies. (Let us point out, however, that Lucier [13,16] did use Kuznetsov approximation theory for defining adaptivity methods for conservation laws, and that Cockburn and Chen [1] obtained a result similar to ours (Theorem 2.1) to prove optimal error estimates for the one-dimensional drift-diffusion model for semiconductor device simulation.) In this paper, we fully exploit this point of view and use Kuznetsov's approach to obtain a simple and powerful *a posteriori* error estimate for arbitrary numerical schemes for the general scalar conservation law.

Other *a posteriori* error estimates for the nonlinear case have been obtained by Tadmor [19] and Nessyahu and Tadmor [17] for the one-dimensional case and a strictly convex nonlinearity f . These authors use a duality technique (which leads also to local error estimates) which allows them to obtain an *a posteriori* estimate similar to ours, [17, (2.10a)], the main difference being that we can use the L^1 -norm where they use the Lip' -norm. We want also to mention that recently Johnson and Szepessy [10] obtained *a posteriori* for the much more difficult system of gas dynamics for the Shock-Capturing Streamline Diffusion method.

The paper is organized as follows. In §2, we display the *a posteriori* error estimate, Theorem 2.1, which is a straightforward application of Kuznetsov's approximation theory [12] (for the sake of completeness, we display a proof of the result in the Appendix). In §3, we use Theorem 2.1 to show that the problem of obtaining an *a posteriori* can be rewritten as a minimization problem (involving interpolation operators only), which can be easily used to define adaptivity strategies. This line of work will be further developed elsewhere. In this paper, we restrict ourselves to displaying preliminary numerical results that show that the obtained upper bound for the error is sharp. We end in §4 with some concluding remarks.

2. The *a posteriori* estimate. In what follows, we denote by u any continuous function from $[0, T]$ to $L^1(\mathbb{R}^d)$. Moreover, we assume that u is a smooth function, except on a finite number of $(d-1)$ -dimensional surfaces $\mathcal{C}_i, i = 1, \dots, I$. We denote by $n_{\mathcal{C}_i}$ the unit normal to \mathcal{C}_i such that its component on the t -axis is negative. The jump of $G(u)$ at the point P on surface \mathcal{C}_i , $[G(u)](P)$, is defined as follows:

$$[G(u)](P) = \lim_{\rho \downarrow 0} (G(u(P + \rho n_{\mathcal{C}_i})) - G(u(P - \rho n_{\mathcal{C}_i}))).$$

We set $\mathcal{C}_i(T) = (0, T) \times \mathbb{R}^d \cap \mathcal{C}_i$ and $\mathcal{C}(T) = \cup_{i=1}^I \mathcal{C}_i(T)$. Finally, we set $\Omega(T) = (0, T) \times \mathbb{R}^d \setminus \mathcal{C}(T)$.

With this notation, we have the following approximation result.

THEOREM 2.1 (THE *a posteriori* ESTIMATE). *Let u be as above and let v be the entropy solution of the conservation law (1.1). Then,*

$$\|u(T) - v(T)\|_{L^1(\mathbb{R}^d)} \leq \Phi(v_0, u; T),$$

where

$$\Phi(v_0, u; T) = \|u(0) - v_0\|_{L^1(\mathbb{R}^d)} + \|Residual(u)\|_{L^1(\Omega(T))} + \sum_{i=1}^I \|Residual(u)\|_{L^1(\mathcal{C}_i(T))},$$

where

$$Residual(u) = \begin{cases} u_t + \nabla \cdot f(u), & \text{on } \Omega(T), \\ \max \left\{ 0, \sup_{c \in [a, b]} \{ ([F(u, c)], [U(u - c)]) \cdot n_{C_i} \} \right\} & \text{on } C_i(T), \end{cases}$$

and $U(u - c) = |u - c|$, $F(u, c) = \text{sign}(u - c)(f(u) - f(c))$. The interval $[a, b]$ is the range of the initial data v_0 .

The term $\|Residual(u)\|_{L^1(\Omega(T))}$ is the truncation error on the smooth regions of the approximation defined by u . The term $\|Residual(u)\|_{L^1(C_i(T))}$ measures how close to satisfying the entropy condition are the discontinuities of u on the curve $C_i(T)$. To see this, let us consider the one-dimensional case, $d = 1$, for the sake of simplicity. If we write $C_i(T) = \{(x, t), x = x_i(t), 0 \leq t_{i1} \leq t \leq t_{i2} \leq T\}$, we have the familiar expression

$$\|Residual(u)\|_{L^1(C_i(T))} = \int_{t_{i1}}^{t_{i2}} \max \left\{ 0, \sup_{c \in [a, b]} \{ ([F(u, c)] - \frac{dx_i}{dt} [U(u - c)])(s, x_i(s)) \} \right\} ds.$$

It is well-known that $\|Residual(u)\|_{L^1(C_i)} = 0$ if and only if the discontinuities of u on the curve $C_i(T)$ are discontinuities of an entropy solution of the conservation law (1.1a). Moreover, this only happens if the so-called Rankine-Hugoniot condition, $\frac{dx_i}{dt} = \frac{[f(u)]}{[u]}$, is verified on the curve $C_i(T)$.

In this way, if u is an entropy solution of the conservation law (1.1a), then $Residual(u) \equiv 0$ and the estimate of Theorem 2.1 becomes

$$\|u(T) - v(T)\|_{L^1(\mathbb{R}^d)} \leq \|u(0) - v_0\|_{L^1(\mathbb{R}^d)}.$$

as expected.

Now, suppose that we have a numerical scheme that produces an approximate solution u_h at times t^n , $n = 1, \dots, N$. Let us denote by \mathcal{U} the set of all the *interpolates* u such that $u(t^n) = u_h(t^n)$ for $n = 1, \dots, N$. Then, a simple application of Theorem 2.1, gives the following *a posteriori* error estimate:

$$\|u_h(t^n) - v(t^n)\|_{L^1(\mathbb{R}^d)} \leq \inf_{u \in \mathcal{U}} \Phi(v_0, u; t^n), \quad n = 1, \dots, N. \quad (\star)$$

This estimate is the main contribution of this paper. As we claimed in the introduction, this estimate is totally independent (i) of the dimension of the space, (ii) of the type of nonlinearity f , and (iii) of the numerical method used to produce approximations to the entropy solution v of (1.1).

3. On the sharpness of the *a posteriori* estimate. The main application of the estimate (\star) is the devising of adaptivity strategies. The goal of these strategies is to ensure that the error is smaller than a given tolerance, say ϵ , with the smallest possible computational effort. In order to achieve this goal, it is natural to try to enforce the following inequality:

$$\Phi(v_0, u^*; t^N) \leq \epsilon,$$

where u^* is the member of \mathcal{U} we chose to work with. It is clear that the first task we have to face, before attempting to define any adaptivity strategies, is to study, at least numerically, the behavior of the ratio

$$r(u^*, t^N) = \frac{\Phi(v_0, u^*; t^N)}{\|u_h(t^N) - v(t^N)\|_{L^1(\mathbb{R}^d)}},$$

since we do not want this ratio to be too big. The ideal situation occurs, of course, when we can find a simple interpolation u^* such that $r(u^*, t^N) \equiv 1$. In what follows, we present some preliminary numerical results that show that this ideal situation can be easily achieved.

We consider the approximate solution given by the well-known Engquist-Osher scheme in uniform grids $\{t^n = n \Delta t\}_{n=0}^{N_t} \times \{x_i = i \Delta x\}_{i=0}^{N_x}$ defined on the space domain is the interval $[0, 1)$ with periodic boundary conditions. We use two ways of interpolating the values u_i^n generated by the Engquist-Osher scheme. We denote by u_1^* the standard piecewise-bilinear function such that $u_1^*(t^n, x_i) = u_i^n$. We also define a more sophisticated interpolation function u_2^* which takes into account the nature of the conservation law. We define u_2^* as follows. First, we take $u^*(n \Delta t)$ to be the piecewise-linear function such that $u_2^*(t^n, x_i) = u_i^n$. We extend u_2^* to the strips $(n \Delta t, (n+1) \Delta t) \times [0, 1)$ as follows. We consider all the trapezoids formed by the lines $t = t^n$, $t = t^{n+1}$ and the approximate characteristics $x = x_i + f'(u(t^n, x_i))(t - t^n)$ and $x = x_i + f'(u(t^{n+1}, x_i))(t - t^{n+1})$. We then defined u inside of each of these trapezoids as the standard bilinear isoparametric interpolate of the values of u at the vertices; see Ciarlet [2, pages 229–230]. This construction is well-defined for the examples considered below.

All the integrals are computed by using the mid-point rule.

Example 1. We take $f(v) = v$, $v_0(x) = 1 + .5 \sin(\pi(2x - 1))$, and $T = .1$. The behavior of the ratio $r(u^*, T)$ is displayed in Tables 1 and 2. We can see that with the simple interpolate u_1^* , the ratio $r(u_1^*, T)$ is not bigger than 1.3 and that with the more sophisticated interpolate u_2^* , the ratio $r(u_2^*, T)$ differs very little from 1. These results confirm that the error estimate is sharp and that even with a very simple interpolation procedure very good results can be obtained.

Example 2. We take $f(v) = v^2/2$, $v_0(x) = 1 + .5 \sin(\pi(2x - 1))$, and $T = .1$. The results obtained for the previous example also hold in this case, as we can see in Tables 3 and 4. (Note that ratios less than 1 might be obtained because we are evaluating the integrals in an approximate way.)

Example 3. We take $f(v) = v$, $T = 1/\sqrt{2}$, and $v_0(x) = \begin{cases} 1, & \text{for } x \in (.4, .6), \\ 0, & \text{elsewhere.} \end{cases}$

The results are displayed in Tables 5 and 6. We see, once more, that the ratio $r(u^*, T)$ is close to 1 and that, unlike the previous cases, the results with u_2^* are only slightly better than the results obtained with u_1^* . This might be explained by the fact that (i) the ‘smooth region’ of the exact solution is flat and so the contribution of the *Residual* in that region is negligible, and by the fact that (ii) the approximate solution is very smooth around the location of the exact discontinuity.

4. Concluding remarks. The preliminary results we have presented in §3 show that the error estimate $(*)$ in §2 is sharp indeed. In a forthcoming paper, we study the behavior of the ratio $r(u, T)$ for several numerical schemes in the case of general nonlinearities. It must be pointed out that, in order to treat the general case, we must use interpolates u that have discontinuities. If only continuous functions u are used, there is no way to distinguish entropy-violating discontinuities from entropy-satisfying discontinuities.

Table 1

Example 1: Behavior of $r(u_1^, T)$ with respect to Δx and $\lambda = \Delta t/\Delta x$.*

$1/\Delta x$	$\lambda = .25$	$\lambda = .50$	$\lambda = .75$
10	1.149	1.157	1.146
20	1.184	1.180	1.155
40	1.206	1.202	1.183
80	1.220	1.216	1.204
160	1.227	1.225	1.218
320	1.230	1.229	1.226
640	1.232	1.231	1.230

Table 2

Example 1: Behavior of $r(u_2^, T)$ with respect to Δx and $\lambda = \Delta t/\Delta x$.*

$1/\Delta x$	$\lambda = .25$	$\lambda = .50$	$\lambda = .75$
10	1.02357	1.03430	1.04607
20	1.00642	1.00997	1.00633
40	1.00169	1.00275	1.00225
80	1.00044	1.00073	1.00043
160	1.00011	1.00019	1.00013
320	1.00003	1.00005	1.00004
640	1.00002	1.00002	1.00003

Table 3

Example 2: Behavior of $r(u_1^, T)$ with respect to Δx and $\lambda = \Delta t/\Delta x$.*

$1/\Delta x$	$\lambda = .25$	$\lambda = .50$	$\lambda = .75$
10	1.177	1.161	1.165
20	1.197	1.187	1.175
40	1.232	1.218	1.197
80	1.249	1.238	1.215
160	1.257	1.249	1.225
320	1.262	1.253	1.231
640	1.264	1.257	1.234

Table 4*Example 2: Behavior of $r(u_2^*, T)$ with respect to Δx and $\lambda = \Delta t/\Delta x$.*

$1/\Delta x$	$\lambda = .25$	$\lambda = .50$	$\lambda = .75$
10	1.04012	1.03034	1.0504
20	1.00009	1.00025	1.01097
40	1.00071	0.99700	1.00065
80	1.00055	0.99875	0.99878
160	0.99947	0.99931	0.99895
320	1.00014	0.99968	0.99935
640	1.00019	0.99990	0.99964

Table 5*Example 3: Behavior of $r(u_1^*, T)$ with respect to Δx and $\lambda = \Delta t/\Delta x$.*

$1/\Delta x$	$\lambda = .25$	$\lambda = .50$	$\lambda = .75$
10	1.385	1.347	1.361
20	1.233	1.212	1.248
40	1.155	1.149	1.186
80	1.116	1.112	1.135
160	1.087	1.081	1.096
320	1.064	1.059	1.069
640	1.045	1.042	1.049

Table 6*Example 3: Behavior of $r(u_2^*, T)$ with respect to Δx and $\lambda = \Delta t/\Delta x$.*

$1/\Delta x$	$\lambda = .25$	$\lambda = .50$	$\lambda = .75$
10	1.301	1.265	1.287
20	1.159	1.141	1.184
40	1.088	1.085	1.132
80	1.057	1.059	1.094
160	1.038	1.040	1.066
320	1.028	1.029	1.048
640	1.020	1.021	1.034

Acknowledgments. The first author would like to thank Jonathan Goodman and Monika Geiben for fruitful conversations leading to the writing of this paper.

Appendix. Proof of Theorem 2.1. By Kuznetsov [12, Lemma 2], we have that, for $0 < \epsilon_t < T$ and ϵ_x ,

$$\begin{aligned} \|u(T) - v(T)\|_{L^1(\mathbb{R}^d)} &\leq \|u(0) - v_0\|_{L^1(\mathbb{R}^d)} + E(u, v; T) \\ &\quad + 2\epsilon_x \{ \|v_0\|_{TV(\mathbb{R}^d)} + \|\nabla u\|_{L^\infty(0, T; L^1(\mathbb{R}^d))} \} \\ &\quad + 2\epsilon_t \{ \|f'\| \|v_0\|_{TV(\mathbb{R}^d)} + \|u_t\|_{L^\infty(0, T; L^1(\mathbb{R}^d))} \}, \end{aligned}$$

where $\|f'\| = \sup_{|n|=1} \sup_{v \in [a, b]} |f'(v) \cdot n|$ and

$$E(u, v; \tau) = \int_0^\tau \int_{\mathbb{R}^d} \Theta_\tau(u, v(t, x); t, x) dx dt,$$

where

$$\begin{aligned} \Theta_\tau(u, c; t, x) &= - \int_0^\tau \int_{\mathbb{R}^d} U(u(t', x') - c) \partial_{t'} \varphi(t, x, t', x') dx' dt' \\ &\quad + \int_{\mathbb{R}^d} U(u(\tau, x') - c) \varphi(t, x, \tau, x') dx' - \int_{\mathbb{R}^d} U(u(0, x') - c) \varphi(t, x, 0, x') dx' \\ &\quad - \int_0^\tau \int_{\mathbb{R}^d} F(u(t', x'), c) \cdot \nabla_{x'} \varphi(t, x, t', x') dx' dt'. \end{aligned}$$

The function $\varphi = \varphi(t, x, t', x')$ is taken as follows:

$$\varphi = w_{\epsilon_t}(t - t') \prod_{i=1}^d w_{\epsilon_x}(x_i - x'_i), \quad (x, t), (x', t') \in \mathbb{R}^d \times \mathbb{R}^+,$$

where ϵ_t and ϵ_x are two arbitrary positive numbers and $w_\lambda(s) = \frac{1}{\lambda} w(\frac{s}{\lambda})$, for any $s \in \mathbb{R}$, $\lambda = \epsilon_t, \epsilon_x$. The function w is a nonnegative, even, smooth, compact supported function of unit mass.

It is enough to prove Theorem 2.1 in the case in which the function u has a single discontinuity surface, $\mathcal{C}(T) = \mathcal{C}_1(T)$, which divides $(0, T) \times \mathbb{R}^d$ in two disjoint subsets whose union is $\Omega(T)$. In this case, after performing integration by parts, we obtain

$$\begin{aligned} \Theta(u, v; \tau) &= \int_{\Omega(T)} \text{Residual}(u(t', x')) \text{sign}(u(t', x') - v(t, x)) \varphi(t, x, t', x') dx' dt' \\ &\quad + \int_{\mathcal{C}(T)} ([F(u, c)], [U(u - c)])(t', x') \cdot n_{\mathcal{C}}(t', x') \varphi(t, x, t', x') d\Gamma'. \end{aligned}$$

Since, by the definition of φ , $\int_0^\tau \int_{\mathbb{R}^d} \varphi(t, x, t', x') dx dt \leq 1$, we obtain

$$\begin{aligned} E(u, v; \tau) &\leq \int_{\Omega(T)} |\text{Residual}(u(t', x'))| dx' dt' \\ &\quad + \int_{\mathcal{C}(T)} \max \left\{ 0, \sup_{c \in \mathbb{R}} \{ ([F(u, c)], [U(u - c)])(t', x') \cdot n_{\mathcal{C}}(t', x') \} \right\} d\Gamma' \\ &= \|\text{Residual}(u)\|_{L^1(\Omega(T))} + \|\text{Residual}(u)\|_{L^1(\mathcal{C}(T))}. \end{aligned}$$

We can now let the parameters ϵ_t and ϵ_x go to zero to obtain Theorem 2.1.

REFERENCES

- [1] Z. CHEN AND B. COCKBURN, *Error estimates for a finite element method for the drift-diffusion semiconductor device equations*, SIAM J. Numer. Anal., to appear in 1994.
- [2] P.G. CIARLET, *The finite element method for elliptic problems*, North Holland, 1979.
- [3] B. COCKBURN, *The quasi-monotone schemes for scalar conservation laws, I*, SIAM J. Numer. Anal., 26 (1989), pp. 1325–1341.
- [4] ———, *The quasi-monotone schemes for scalar conservation laws, II*, SIAM J. Numer. Anal., 27 (1990), pp. 247–258.
- [5] ———, *The quasi-monotone schemes for scalar conservation laws, III*, SIAM J. Numer. Anal., 27 (1990), pp. 259–276.
- [6] B. COCKBURN AND C.W. SHU, *TVB Runge-Kutta Local Projection Discontinuous Galerkin Finite Element Method for Scalar Conservation Laws II: General Framework*, Math. Comp., 52 (1989), pp. 411–435.
- [7] B. COCKBURN, F. COQUEL, AND P. LEFLOCH, *An error estimate for finite-volume methods for conservations laws*, Math. Comp., 63, pp. 77–103.
- [8] B. COCKBURN AND P.-A. GREMAUD, *An error estimate for finite element methods for conservations laws*, University of Minnesota Supercomputer Institute Research Report 93-128 (August 1993), SIAM J. Numer. Anal., to appear in 1996.
- [9] ———, *A priori error estimates for scalar conservation laws. Part I: The general approach*, submitted to Math. Comp. (August 1994).
- [10] C. JOHNSON AND A. SZEPESY, *A posteriori error estimate for a finite element method*, Preprint (1993).
- [11] S.N. KRŽKOV, *First order quasilinear equations in several independent variables*, Math. USSR Sbornik, 10 (1970), pp. 217–243.
- [12] N.N. KUZNETSOV, *Accuracy of some approximate methods for computing the weak solutions of a first-order quasi-linear equation*, USSR Comp. Math. and Math. Phys., 16 (1976), pp. 105–119.
- [13] B. J. LUCIER, *A stable adaptive scheme for hyperbolic conservation laws*, SIAM J. Numer. Anal., 22 (1985), pp. 180–203.
- [14] ———, *Error bounds for the methods of Glimm, Godunov and LeVeque*, SIAM J. Numer. Anal., 22 (1985), pp. 1074–1081.
- [15] ———, *On nonlocal monotone difference schemes for scalar conservation laws*, Math. Comp., 47, No. 175 (1986), pp. 19–36.
- [16] ———, *A moving mesh numerical method for hyperbolic conservation laws*, Math. Comp., 46 (1986), pp. 59–69.
- [17] S. NESSYAHU AND E. TADMOR, *The convergence rate of approximate solutions for nonlinear scalar conservation laws*, SIAM J. Numer. Anal., 29 (1992), pp. 1–15.
- [18] R. SANDERS, *On Convergence of monotone finite difference schemes with variable spatial differencing*, Math. Comp., 40 (1983), pp. 91–106.
- [19] E. TADMOR, *Local error estimates for discontinuous solutions of nonlinear hyperbolic equations*, SIAM J. Numer. Anal., 28 (1991), pp. 891–906.
- [20] J.-P. VILA, *Convergence and error estimates in finite volume schemes for general multidimensional scalar conservation laws*, Model. Math. Anal. Numer., 28 (1994), pp. 267–295.

#	Author/s	Title
1173	Zhangxin Chen & Bernardo Cockburn,	Convergence of a finite element method for the drift-diffusion semiconductor device equations: The multidimensional case
1174	Boris Mordukhovich,	Optimization and finite difference approximations of nonconvex differential inclusions with free time
1175	Avner Friedman, David S. Ross, and Jianhua Zhang,	A Stefan problem for reaction-diffusion system
1176	Alex Solomonoff,	Fast algorithms for micromagnetic computations
1177	Nikan B. Firoozye,	Homogenization on lattices: Small parameter limits, H -measures, and discrete Wigner measures
1178	G. Yin,	Adaptive filtering with averaging
1179	Wlodzimierz Byrc and Amir Dembo,	Large deviations for quadratic functionals of Gaussian processes
1180	Ilja Schmelzer,	3D anisotropic grid generation with intersection-based geometry interface
1181	Alex Solomonoff,	Application of multipole methods to two matrix eigenproblems
1182	A.M. Latypov,	Numerical solution of steady euler equations in streamline-aligned orthogonal coordinates
1183	Bei Hu & Hong-Ming Yin,	Semilinear parabolic equations with prescribed energy
1184	Bei Hu & Jianhua Zhang,	Global existence for a class of Non-Fickian polymer-penetrant systems
1185	Rongze Zhao & Thomas A. Posbergh,	Robust stabilization of a uniformly rotating rigid body
1186	Mary Ann Horn & Irena Lasiecka,	Uniform decay of weak solutions to a von Kármán plate with nonlinear boundary dissipation
1187	Mary Ann Horn, Irena Lasiecka & Daniel Tataru,	Well-posedness and uniform decay rates for weak solutions to a von Kármán system with nonlinear dissipative boundary conditions
1188	Mary Ann Horn,	Nonlinear boundary stabilization of a von Kármán plate via bending moments only
1189	Frank H. Shaw & Charles J. Geyer,	Constrained covariance component models
1190	Tomasz Luczaka,	A greedy algorithm estimating the height of random trees
1191	Timo Seppäläinen,	Maximum entropy principles for disordered spins
1192	Yuandan Lin, Eduardo D. Sontag & Yuan Wang,	Recent results on Lyapunov-theoretic techniques for nonlinear stability
1193	Svante Janson,	Random regular graphs: Asymptotic distributions and contiguity
1194	Rachid Ababou,	Random porous media flow on large 3-D grids: Numerics, performance, & application to homogenization
1195	Moshe Fridman,	Hidden Markov model regression
1196	Petr Klouček, Bo Li & Mitchell Luskin,	Analysis of a class of nonconforming finite elements for Crystalline microstructures
1197	Steven P. Lalley,	Random series in inverse Pisot powers
1198	Rudy Yaksick,	Expected optimal exercise time of a perpetual American option: A closed-form solution
1199	Rudy Yaksick,	Valuation of an American put catastrophe insurance futures option: A Martingale approach
1200	János Pach, Farhad Shahrokhi & Mario Szegedy,	Application of the crossing number
1201	Avner Friedman & Chaocheng Huang,	Averaged motion of charged particles under their self-induced electric
1202	Joel Spencer,	The Erdős-Hanani conjecture via Talagrand's inequality
1203	Zhangxin Chen,	Superconvergence results for Galerkin methods for wave propagation in various porous media
1204	Russell Lyons, Robin Pemantle & Yuval Peres,	When does a branching process grow like its mean? Conceptual proofs of $L \log L$ criteria
1205	Robin Pemantle,	Maximum variation of total risk
1206	Robin Pemantle & Yuval Peres,	Galton-Watson trees with the same mean have the same polar sets
1207	Robin Pemantle,	A shuffle that mixes sets of any fixed size much faster than it mixes the whole deck
1208	Itai Benjamini, Robin Pemantle & Yuval Peres,	Martin capacity for Markov chains and random walks in varying dimensions
1209	Wlodzimierz Bryc & Amir Dembo,	On large deviations of empirical measures for stationary Gaussian processes
1210	Martin Hildebrand,	Some random processes related to affine random walks
1211	Alexander E. Mazel & Yurii M. Suhov,	Ground states of a Boson quantum lattice model
1212	Roger L. Fosdick & Darren E. Mason,	Single phase energy minimizers for materials with nonlocal spatial dependence
1213	Bruce Hajek,	Load balancing in infinite networks
1214	Petr Klouček,	The transonic flow problems stability analysis and numerical results
1215	Petr Klouček,	On the existence of the entropic solutions for the transonic flow problem
1216	David A. Schmidt & Chjan C. Lim,	Full sign-invertibility and symplectic matrices
1217	Piermarco Cannarsa & Maria Elisabetta Tessitore,	Infinite dimensional Hamilton-Jacobi equations and Dirichlet boundary control problems of parabolic type
1218	Zhangxin Chen,	Multigrid algorithms for mixed methods for second order elliptic problems
1219	Zhangxin Chen,	Expanded mixed finite element methods for linear second order elliptic problems I
1220	Gang Bao,	A note on the uniqueness for an inverse diffraction problem

- 1221 Moshe Fridman, A two state capital asset pricing model
- 1222 Paolo Baldi, Exact asymptotics for the probability of exit from a domain and applications to simulation
- 1223 Carl Dou & Martin Hildebrand, Enumeration and random random walks on finite groups
- 1224 Jaksa Cvitanic & Ioannis Karatzas, On portfolio optimization under “drawdown” constraints
- 1225 Avner Friedman & Yong Liu, A free boundary problem arising in magnetohydrodynamic system
- 1226 Dominic Welsh, Randomised approximation schemes for Tutte-Gröthendieck invariants
- 1227 Zhangxin Chen, Bernardo Cockburn, Carl L. Gardner, & Joseph W. Jerome, Quantum hydrodynamic simulation of hysteresis in the resonant tunneling diode
- 1228 E.G. Kalnins, G.C. Williams, & Willard Miller, Jr., Intrinsic characterisation of the separation constant for spin one and gravitational perturbations in Kerr geometry
- 1229 Zhangxin Chen, Large-scale averaging analysis of multiphase flow in fractured reservoirs
- 1230 Bruce Hajek & Babu Narayanan, Multigraphs with the most edge covers
- 1231 K.B. Athreya, Entropy maximization
- 1232 F.I. Karpelevich & Yu.M. Suhov, Functional equations in the problem of boundedness of stochastic branching dynamics
- 1233 E. Dibenedetto & V. Vespri, On the singular equation $\beta(u)_t = \Delta u$
- 1234 M.Ya. Kelbert & Yu.M. Suhov, The Markov branching random walk and systems of reaction-diffusion (Kolmogorov-Petrovskii-Piskunov) equations
- 1235 M. Hildebrand, Random walks on random regular simple graphs
- 1236 W.S. Don & A. Solomonoff, Accuracy enhancement for higher derivatives using Chebyshev collocation and a mapping technique
- 1237 D. Gurarie, Symmetries and conservation laws of two-dimensional hydrodynamics
- 1238 Z. Chen, Finite element methods for the black oil model in petroleum reservoirs
- 1239 G. Bao & A. Friedman, Inverse problems for scattering by periodic structure
- 1240 G. Bao, Some inverse problems in partial differential equations
- 1241 G. Bao, Diffractive optics in periodic structures: The TM polarization
- 1242 C.C. Lim & D.A. Schmidt, On noneven digraphs and symplectic pairs
- 1243 H.M. Soner, S.E. Shreve & J. Cvitanić, There is no nontrivial hedging portfolio for option pricing with transaction costs
- 1244 D.L. Russell & B-Yu Zhang, Exact controllability and stabilizability of the Korteweg-de Vries equation
- 1245 B. Morton, D. Enns & B-Yu Zhang, Stability of dynamic inversion control laws applied to nonlinear aircraft pitch-axis models
- 1246 S. Hansen & G. Weiss, New results on the operator Carleson measure criterion
- 1247 V.A. Malyshev & F.M. Spieksma, Intrinsic convergence rate of countable Markov chains
- 1248 G. Bao, D.C. Dobson & J.A. Cox, Mathematical studies in rigorous grating theory
- 1249 G. Bao & W.W. Symes, On the sensitivity of solutions of hyperbolic equations to the coefficients
- 1250 D.A. Huntley & S.H. Davis, Oscillatory and cellular mode coupling in rapid directional solidification
- 1251 M.J. Donahue, L. Gurvits, C. Darken & E. Sontag, Rates of convex approximation in non-Hilbert spaces
- 1252 A. Friedman & B. Hu, A Stefan problem for multi-dimensional reaction diffusion systems
- 1253 J.L. Bona & B-Y. Zhang, The initial-value problem for the forced Korteweg-de Vries equation
- 1254 A. Friedman & R. Gulliver, Organizers, Mathematical modeling for instructors
- 1255 S. Kichenassamy, The prolongation formula for tensor fields
- 1256 S. Kichenassamy, Fuchsian equations in Sobolev spaces and blow-up
- 1257 H.S. Dumas, L. Dumas, & F. Golse, On the mean free path for a periodic array of spherical obstacles
- 1258 C. Liu, Global estimates for solutions of partial differential equations
- 1259 C. Liu, Exponentially growing solutions for inverse problems in PDE
- 1260 Mary Ann Horn & I. Lasiecka, Nonlinear boundary stabilization of parallelly connected Kirchhoff plates
- 1261 B. Cockburn & H. Gau, A posteriori error estimates for general numerical methods for scalar conservation laws
- 1262 B. Cockburn & P-A. Gremaud, A priori error estimates for numerical methods for scalar conservation laws. Part I: The general approach
- 1263 R. Spigler & M. Vianello, Convergence analysis of the semi-implicit euler method for abstract evolution equations
- 1264 R. Spigler & M. Vianello, WKB-type approximation for second-order differential equations in C^* -algebras
- 1265 M. Menshikov & R.J. Williams, Passage-time moments for continuous non-negative stochastic processes and applications
- 1266 C. Mazza, On the storage capacity of nonlinear neural networks
- 1267 Z. Chen, R.E. Ewing & R. Lazarov, Domain decomposition algorithms for mixed methods for second order elliptic problems
- 1268 Z. Chen, M. Espedal & R.E. Ewing, Finite element analysis of multiphase flow in groundwater hydrology
- 1269 Z. Chen, R.E. Ewing, Y.A. Kuznetsov, R.D. Lazarov & S. Maliassov, Multilevel preconditioners for mixed methods for second order elliptic problems