

Soil and Vegetation Survey of the Willow
National Petroleum Reserve-Alaska

Gabriel Benitez

2024-10-15

Contents

1	Soil and Vegetation Survey of the National Petroleum Reserve-Alaska	5
1.1	Acknowledgements	5
1.2	Usage	5
2	Introduction	9
2.1	Survey Purpose	9
3	General Nature of the Area	11
3.1	Climate	11
3.2	Captioned figures and tables	14
4	Field Work	17
5	Footnotes and citations	19
5.1	Footnotes	19
5.2	Citations	19
6	Blocks	21
6.1	Equations	21
6.2	Theorems and proofs	21
6.3	Callout blocks	21

7 Sharing your book	23
7.1 Publishing	23
7.2 404 pages	23
7.3 Metadata for sharing	23
8 useful code	25
9 Calculate the average temperature for each day across all years	27
10 Create the combined line graph	29
11 Create a function to reorder months from Oct to Sep	31
12 Prepare the data for plotting	33
13 Calculate average cumulative precipitation	35
14 Create the plot	37

Chapter 1

Soil and Vegetation Survey of the National Petroleum Reserve-Alaska

1.1 Acknowledgements

This project would have never been possible without the expertise of the scientist, pilots, and the managers of this incredible area. Countless thanks to Tyler Annetts, Jessica Lene-Ashley, Phillip Barber, Luke Breneman, Krista Bryant, Brad Casar, Charolette Crowder, Sara Datson, Noah Hull, Ted Inman, Nic Jelinski, Jamin Johanson, Monica Kopp, Amy Li, Travis Nauman, Nathan Perry, Craig Prink, Nathan Roe, Stephanie Schmidt, Michael Sousa, Michael Singer, Mark Stott, Maddie Tucker.

1.2 Usage

This survey was a cooperative effort of the United States Department of Agriculture, Natural Resources Conservation Service (NRCS) and the United States Department of Interior, Bureau of Land Management (BLM). NRCS was responsible for survey design and methodology, data collection and analysis, and this report. Fieldwork was completed in July and August of '21, '23, and '24. Soil names and descriptions were approved in 2024. Unless indicated otherwise, maps and supporting documentation in this report refer to conditions in the survey area in 2024.

Maps in this report may be copied without permission. However, enlargement of these maps could cause misunderstanding of the detail of mapping. If enlarged,



Figure 1.1: Fish Creek taken from R44 helicopter. Notice the progression of ice-wedge polygon development from the floodplains to uplands.

maps do not show the small areas of contrasting soils and vegetation that could have been shown at a larger scale.

All programs and services of the Natural Resources Conservation Service and the Bureau of Land Management are offered on a nondiscriminatory basis, without regard to race, color, national origin, gender, religion, age, disability, political beliefs, sexual orientation, marital or family status.

```
##  
## Attaching package: 'dplyr'  
  
## The following objects are masked from 'package:stats':  
##  
##     filter, lag  
  
## The following objects are masked from 'package:base':  
##  
##     intersect, setdiff, setequal, union
```


Chapter 2

Introduction

The Willow area of the National Petroleum Reserve-Alaska (NPR-A) is in the Arctic Coastal Plain of the North Slope of Alaska, approximately 90 miles (145 km) west of Deadhorse. These lands were originally inhabited by various Alaska Native groups, including the Iñupiat people for thousands of years. In 1923, President Warren G. Harding established the Naval Petroleum Reserve No.4 (NPR-4) through an executive order, which brought this area under federal ownership without consultation or compensation to the Alaska Native peoples. In 1976, the Naval Petroleum Reserves Production Act renamed the area to the National Petroleum Reserve-Alaska and transferred the management of the reserve from the United States Navy to the United States Department of Interior, Bureau of Land Management. In recent years, there has been increased interest in developing the oil resources in the NPR-A, exemplified by projects such as the Willow Project. The NPR-A is approximately 24 million acres stretching across the North Slope of Alaska. The soil and vegetation survey area of the Willow area is approximately 500,000 acres on the Eastern most side of the NPR-A.

2.1 Survey Purpose


The primary purpose of the survey was to describe and map the soils and vegetation of the Willow area in the NPR-A. Area soils and vegetation were mapped at a scale of 1:24,000 and detailed description of the map units, soil types, and vegetation cover types were developed.

Chapter 3


General Nature of the Area

3.1 Climate

The Arctic Coastal Plain of Alaska is characterized by a harsh tundra climate, classified as ET in the Köppen system. This region experiences extreme temperature variations, with brutally cold winters where average temperatures often plummet below -22°F, and brief, cool summers barely reaching 37-54°F in July. The average annual temperature ranges from 8 to 14°F. The average freeze-free period is fewer than 5 days to 20 days. Freezing temperatures can occur in any month.



Despite its frigid nature, the coastal plain receives surprisingly little precipitation, typically ranging from 4-10 inches annually. Most of this falls as snow, blanketing the landscape for much of the year. The average annual snowfall is about 50 to 100 centimeters. The low precipitation, coupled with minimal evaporation rates due to the cold, creates a unique hydrologic balance that doesn't quite fit the traditional definition of a desert.



One of the most striking features of this climate is the dramatic swing in daylight hours throughout the year. Summers bring the phenomenon of the midnight sun, with 24 hours of continuous daylight, while winters plunge the region into weeks of polar night. This extreme light regime profoundly impacts biological rhythms and human activities alike. The coastal plain is also known for its windy conditions, with strong easterly winds being common. These winds, combined with the already frigid temperatures, can create dangerously low wind chill factors.

The growing season in the Arctic Coastal Plain is exceptionally brief, usually lasting only 50-60 days. This short window of relatively warmer temperatures and thawed ground supports a unique but fragile tundra ecosystem. Coastal areas are further influenced by sea ice, which is typically present for 8-9 months of the year, affecting local weather patterns and wildlife migrations.

Underlying this harsh surface climate is a layer of continuous permafrost, a defining characteristic of the region. The active layer, which thaws seasonally, is typically shallow, extending only 30-50 cm (12-20 inches) deep. This frozen ground significantly influences the area's ecology and presents unique challenges for construction and resource extraction.

1. Label the heading: # Hello world {#nice-label}.
 - Leave the label off if you like the automated heading generated based on your heading title: for example, # Hello world = # Hello world {#hello-world}.

- To label an un-numbered heading, use: `# Hello world {-#nice-label}` or `{# Hello world .unnumbered}`.
2. Next, reference the labeled heading anywhere in the text using `\@ref(nice-label)`; for example, please see Chapter ??.
- If you prefer text as the link instead of a numbered reference use: any text you want can go here.

3.2 Captioned figures and tables

Figures and tables *with captions* can also be cross-referenced from elsewhere in your book using `\@ref(fig:chunk-label)` and `\@ref(tab:chunk-label)`, respectively.

See Figure 3.1.

```
par(mar = c(4, 4, .1, .1))
plot(pressure, type = 'b', pch = 19)
```




Figure 3.1: Here is a nice figure!

Don't miss Table 3.1.

```
knitr::kable(
  head(pressure, 10), caption = 'Here is a nice table!',
```

Table 3.1: Here is a nice table!

temperature	pressure
0	0.0002
20	0.0012
40	0.0060
60	0.0300
80	0.0900
100	0.2700
120	0.7500
140	1.8500
160	4.2000
180	8.8000

```
booktabs = TRUE  
)
```


Chapter 4

Field Work

For this project getting to the site with enough gear to be self-sustainable was a logistical challenge. Thankfully the BLM has a remote field station, Inigok. Inigok is a

You can add parts to organize one or more book chapters together. Parts can be inserted at the top of an .Rmd file, before the first-level chapter heading in that same file.

Add a numbered part: # (PART) Act one {-} (followed by # A chapter)

Add an unnumbered part: # (PART*) Act one {-} (followed by # A chapter)

Add an appendix as a special kind of un-numbered part: # (APPENDIX) Other stuff {-} (followed by # A chapter). Chapters in an appendix are prepended with letters instead of numbers.

Chapter 5

Footnotes and citations

5.1 Footnotes

Footnotes are put inside the square brackets after a caret ^[] . Like this one ¹.

5.2 Citations

Reference items in your bibliography file(s) using @key.

For example, we are using the **bookdown** package [Xie, 2024] (check out the last code chunk in index.Rmd to see how this citation key was added) in this sample book, which was built on top of R Markdown and **knitr** [Xie, 2015] (this citation was added manually in an external file book.bib). Note that the **.bib** files need to be listed in the index.Rmd with the YAML **bibliography** key.

The RStudio Visual Markdown Editor can also make it easier to insert citations:
<https://rstudio.github.io/visual-markdown-editing/#/citations>

¹This is a footnote.

Chapter 6

Blocks

6.1 Equations

Here is an equation.

$$f(k) = \binom{n}{k} p^k (1-p)^{n-k} \quad (6.1)$$

You may refer to using `\@ref(eq:binom)`, like see Equation (6.1).

6.2 Theorems and proofs

Labeled theorems can be referenced in text using `\@ref(thm:tri)`, for example, check out this smart theorem 6.1.

Theorem 6.1. *For a right triangle, if c denotes the length of the hypotenuse and a and b denote the lengths of the **other** two sides, we have*

$$a^2 + b^2 = c^2$$

Read more here <https://bookdown.org/yihui/bookdown/markdown-extensions-by-bookdown.html>.

6.3 Callout blocks

The R Markdown Cookbook provides more help on how to use custom blocks to design your own callouts: <https://bookdown.org/yihui/rmarkdown-cookbook/custom-blocks.html>

Chapter 7

Sharing your book

7.1 Publishing

HTML books can be published online, see: <https://bookdown.org/yihui/bookdown/publishing.html>

7.2 404 pages

By default, users will be directed to a 404 page if they try to access a webpage that cannot be found. If you'd like to customize your 404 page instead of using the default, you may add either a `_404.Rmd` or `_404.md` file to your project root and use code and/or Markdown syntax.

7.3 Metadata for sharing

Bookdown HTML books will provide HTML metadata for social sharing on platforms like Twitter, Facebook, and LinkedIn, using information you provide in the `index.Rmd` YAML. To setup, set the `url` for your book and the path to your `cover-image` file. Your book's `title` and `description` are also used.

This `gitbook` uses the same social sharing data across all chapters in your book—all links shared will look the same.

Specify your book's source repository on GitHub using the `edit` key under the configuration options in the `_output.yml` file, which allows users to suggest an edit by linking to a chapter's source file.

Read more about the features of this output format here:

<https://pkgs.rstudio.com/bookdown/reference/gitbook.html>

Or use:

```
?bookdown::gitbook
```

Chapter 8

useful code

this is how you commit and push stuff in terminal git add .

git commit -m "commit message"

git push

ctrl + Alt + i = insert r script echo = false means the scrubs dont see code

```
library(tidyr) library(ggplot2) library(readr) library(dplyr) library(tidyverse) library(lubridate)
```

```
df1 <- read_delim("_book/dailyMinTemp.csv", delim = "|")
```

```
data <- df1
```

```
data_long <- data %>% pivot_longer(cols = -c(Water Year, Day), names_to = "Month", values_to = "Temperature")
```

```
month_order <- c("Jan", "Feb", "Mar", "Apr", "May", "Jun", "Jul", "Aug", "Sep", "Oct", "Nov", "Dec") data_long <- data_long %>% mutate(Month = factor(Month, levels = month_order, ordered = TRUE), Date = as.Date(paste("2000", Month, Day, sep = "-"), format = "%Y-%b-%d")) )
```


Chapter 9

**Calculate the average
temperature for each day
across all years**

```
avg_temp <- data_long %>% group_by(Date) %>% summarise(AvgTemperature  
= mean(Temperature, na.rm = TRUE))
```

28CHAPTER 9. CALCULATE THE AVERAGE TEMPERATURE FOR EACH DAY ACROSS ALL Y

Chapter 10

Create the combined line graph

```
ggplot() + # Individual year lines geom_line(data = data_long, aes(x = Date,
y = Temperature), alpha = 0.5) + # Average temperature line geom_line(data
= avg_temp, aes(x = Date, y = AvgTemperature), color = "black", size
= 1.2) + scale_x_date(date_labels = "%b", date_breaks = "1 month") +
scale_color_viridis_d(option = "plasma") + # Color-blind friendly palette
labs(title = "Minimum Daily Temperature 2004-2024", x = "Month", y =
"Temperature (Fahrenheit)") color = "Water Year"

library(tidyr) library(ggplot2) library(readr) library(dplyr) library(tidyverse) li-
brary(lubridate)

df1 <- read_delim("_book/dailyPrecip.csv", delim = "|")

data <- df1 %>% rename(Year = "Water Year")

data_long <- data %>% pivot_longer(cols = -c(Year, Day), names_to =
"Month", values_to = "Precip")

data_long <- data_long %>% mutate( MonthNum = match(Month,
month.abb), Date = make_date(Year, MonthNum, Day), MonthNum = NULL
) %>% mutate(Date = ymd(Date))
```


Chapter 11

Create a function to reorder months from Oct to Sep

```
reorder_month <- function(x) { factor(x, levels = c("Oct", "Nov", "Dec",  
"Jan", "Feb", "Mar", "Apr", "May", "Jun", "Jul", "Aug", "Sep")) }
```

32CHAPTER 11. CREATE A FUNCTION TO REORDER MONTHS FROM OCT TO SEP

Chapter 12

Prepare the data for plotting

```
data_plot <- data_long %>% mutate( WaterYear = ifelse(month(Date) >= 10, Year, Year - 1), Month = reorder_month(Month) ) %>% filter(Day == days_in_month(Date)) %>% arrange(WaterYear, Month)
```


Chapter 13

Calculate average cumulative precipitation

```
avg_precip <- data_plot %>% group_by(Month) %>% summarize(AvgPrecip  
= mean(Precip, na.rm = TRUE))
```

36 CHAPTER 13. CALCULATE AVERAGE CUMULATIVE PRECIPITATION

Chapter 14

Create the plot

```
ggplot() + geom_line(data = data_plot, aes(x = Month, y = Precip,  
group = WaterYear), size = 1, color = "grey80") + geom_line(data =  
avg_precip, aes(x = Month, y = AvgPrecip, group = 1), size = 1.5,  
color = "black") + geom_point(data = avg_precip, aes(x = Month, y =  
AvgPrecip), size = 1, color = "black") + scale_x_discrete(limits = lev-  
els(reorder_month(avg_precip$Month))) + labs( title = "Average Cumulative  
Precipitation by Month", x = "Month", y = "Cumulative Precipitation  
(Inches)" ) + theme_minimal() + theme( legend.position = "none", axis.text.x  
= element_text(angle = 45, hjust = 1), panel.grid.minor = element_blank() )
```


Bibliography

Yihui Xie. *Dynamic Documents with R and knitr*. Chapman and Hall/CRC, Boca Raton, Florida, 2nd edition, 2015. URL <http://yihui.org/knitr/>. ISBN 978-1498716963.

Yihui Xie. *bookdown: Authoring Books and Technical Documents with R Markdown*, 2024. URL <https://github.com/rstudio/bookdown>. R package version 0.40.