

Foundations of Computing

Lecture 22

Arkady Yerukhimovich

April 13, 2023

Outline

1 Lecture 21 Review

2 More \mathcal{NP} -Complete Problems

3 co- \mathcal{NP}

Lecture 21 Review

- \mathcal{P} and \mathcal{NP}
- Polynomial-Time Reductions
- \mathcal{NP} -completeness of SAT

Outline

1 Lecture 21 Review

2 More \mathcal{NP} -Complete Problems

3 co- \mathcal{NP}

What We Already Know

- ① SAT is \mathcal{NP} -complete

What We Already Know

- ① SAT is \mathcal{NP} -complete
- ② 3-SAT is \mathcal{NP} -complete

What We Already Know

- ① SAT is \mathcal{NP} -complete
- ② 3-SAT is \mathcal{NP} -complete
- ③ $3\text{-SAT} \leq_P \text{CLIQUE}$ – So CLIQUE in \mathcal{NP} -complete

What We Already Know

- ① SAT is NP -complete
- ② 3-SAT is NP -complete
- ③ $\text{3-SAT} \leq_P \text{CLIQUE}$ – So CLIQUE in NP -complete
- ④ $\text{3-SAT} \leq_P \text{Vertex Cover}$

What We Already Know

- ① SAT is NP -complete
- ② 3-SAT is NP -complete
- ③ $\text{3-SAT} \leq_P \text{CLIQUE}$ – So CLIQUE in NP -complete
- ④ $\text{3-SAT} \leq_P \text{Vertex Cover}$
- ⑤ $\text{Vertex Cover} \leq_P \text{Independent Set}$

What We Already Know

- ① SAT is NP -complete
- ② 3-SAT is NP -complete
- ③ $\text{3-SAT} \leq_P \text{CLIQUE}$ – So CLIQUE in NP -complete
- ④ $\text{3-SAT} \leq_P \text{Vertex Cover}$
- ⑤ $\text{Vertex Cover} \leq_P \text{Independent Set}$
- ⑥ More on the HW

A Key Tool to Build Reductions



A Key Tool to Build Reductions



Gadgets

A Key Tool to Build Reductions



Gadgets

- Gadgets are structures in the target problem that can simulate structures in the source problem

A Key Tool to Build Reductions



Gadgets

- Gadgets are structures in the target problem that can simulate structures in the source problem
- For example, in proof of $3\text{SAT} \leq_P \text{CLIQUE}$
 - We replaced each clause with set of 3 nodes
 - Edges were used to capture common variables between clauses

A Key Tool to Build Reductions



Gadgets

- Gadgets are structures in the target problem that can simulate structures in the source problem
- For example, in proof of $3\text{SAT} \leq_P \text{CLIQUE}$
 - We replaced each clause with set of 3 nodes
 - Edges were used to capture common variables between clauses
- Similarly in proof of $3\text{SAT} \leq_P \text{Vertex Cover}$
 - We replaced each clause with a triangle and

A Key Tool to Build Reductions



Gadgets

- Gadgets are structures in the target problem that can simulate structures in the source problem
- For example, in proof of $3\text{SAT} \leq_P \text{CLIQUE}$
 - We replaced each clause with set of 3 nodes
 - Edges were used to capture common variables between clauses
- Similarly in proof of $3\text{SAT} \leq_P \text{Vertex Cover}$
 - We replaced each clause with a triangle and
 - each variable with a pair of nodes connected by an edge

NAE-kSAT Problem

NAE-kSAT = { $\langle \phi \rangle \mid \phi$ is in k -CNF and ϕ has a satisfying assignment s.t.
each clause has at least one 0 and at least one 1}

$$(x_1 \vee x_2 \vee x_3) \wedge (x_1 \vee \neg x_2 \vee \neg x_3)$$

NAE-kSAT Problem

$\text{NAE-kSAT} = \{\langle \phi \rangle \mid \phi \text{ is in } k\text{-CNF and } \phi \text{ has a satisfying assignment s.t. each clause has at least one 0 and at least one 1}\}$

Definition:

- x is an NAE-assignment of ϕ if $\phi(x) = 1$ and x does not assign all the same variables to any clause

*Lemma: If x is NAE-assignment
 \bar{x} is also NAE-assignment*

NAE-kSAT Problem

$\text{NAE-kSAT} = \{\langle \phi \rangle \mid \phi \text{ is in } k\text{-CNF and } \phi \text{ has a satisfying assignment s.t. each clause has at least one 0 and at least one 1}\}$

Definition:

- x is an NAE-assignment of ϕ if $\phi(x) = 1$ and x does not assign all the same variables to any clause

Lemma: If x is NAE-assignment of ϕ then \bar{x} is NAE-assignment of ϕ

Proof:

- x must assign at least one 1 and at least one 0 to every clause
- \bar{x} must also have at least one 1 and one 0 in every clause
- This means every clause is satisfied, and ϕ is satisfied since it's CNF

Goal

Prove that NAE-3SAT is \mathcal{NP} -complete: $3\text{SAT} \leq_P \text{NAE-3SAT}$

$3\text{SAT} \leq_P \text{NAE-4SAT}$

$3\text{SAT} \leq_P \text{NAE-4SAT}$

- We need a reduction f that takes 3SAT instance ϕ and converts it into NAE-4SAT instance ϕ'

$3\text{SAT} \leq_P \text{NAE-4SAT}$

- We need a reduction f that takes 3SAT instance ϕ and converts it into NAE-4SAT instance ϕ'
 - If ϕ is satisfiable, ϕ' is NAE-satisfiable

$3SAT \leq_P NAE\text{-}4SAT$

- We need a reduction f that takes 3SAT instance ϕ and converts it into NAE-4SAT instance ϕ'
 - If ϕ is satisfiable, ϕ' is NAE-satisfiable
 - If ϕ' is NAE-satisfiable, ϕ is satisfiable

$$\begin{cases} A \Rightarrow D \\ \neg B \Rightarrow \neg A \end{cases}$$

f is s.t.

$\text{if } x \in 3SAT \Rightarrow f(x) \in NAE\text{-}4SAT$

$\text{if } x \notin 3SAT \Rightarrow f(x) \notin NAE\text{-}4SAT$



$\rightarrow \text{if } f(x) \in NAE\text{-}4SAT \Rightarrow x \in 3SAT$

$3\text{SAT} \leq_P \text{NAE-4SAT}$

- We need a reduction f that takes 3SAT instance ϕ and converts it into NAE-4SAT instance ϕ'
 - If ϕ is satisfiable, ϕ' is NAE-satisfiable
 - If ϕ' is NAE-satisfiable, ϕ is satisfiable
 - Note that this must hold for every clause of ϕ , ϕ'

3SAT \leq_P NAE-4SAT

- We need a reduction f that takes 3SAT instance ϕ and converts it into NAE-4SAT instance ϕ'
 - If ϕ is satisfiable, ϕ' is NAE-satisfiable
 - If ϕ' is NAE-satisfiable, ϕ is satisfiable
 - Note that this must hold for every clause of ϕ , ϕ'
- Idea: Can we build a “gadget” for each clause of ϕ to enforce this condition?

$$(x_1 \vee x_2 \vee x_3) \rightarrow$$

3SAT \leq_P NAE-4SAT

- We need a reduction f that takes 3SAT instance ϕ and converts it into NAE-4SAT instance ϕ'
 - If ϕ is satisfiable, ϕ' is NAE-satisfiable
 - If ϕ' is NAE-satisfiable, ϕ is satisfiable
 - Note that this must hold for every clause of ϕ , ϕ'
- Idea: Can we build a “gadget” for each clause of ϕ to enforce this condition?

$$(x_1 \vee x_2 \vee x_3) \rightarrow (x_1 \vee x_2 \vee x_3 \vee S)$$

3SAT \leq_P NAE-4SAT

- We need a reduction f that takes 3SAT instance ϕ and converts it into NAE-4SAT instance ϕ'
 - If ϕ is satisfiable, ϕ' is NAE-satisfiable
 - If ϕ' is NAE-satisfiable, ϕ is satisfiable
 - Note that this must hold for every clause of ϕ , ϕ'
- Idea: Can we build a “gadget” for each clause of ϕ to enforce this condition?

$$(x_1 \vee x_2 \vee x_3) \rightarrow (x_1 \vee x_2 \vee x_3 \vee S)$$

- Why this works:

3SAT \leq_P NAE-4SAT

- We need a reduction f that takes 3SAT instance ϕ and converts it into NAE-4SAT instance ϕ'
 - If ϕ is satisfiable, ϕ' is NAE-satisfiable
 - If ϕ' is NAE-satisfiable, ϕ is satisfiable
 - Note that this must hold for every clause of ϕ , ϕ'
- Idea: Can we build a “gadget” for each clause of ϕ to enforce this condition?

$$(x_1 \vee x_2 \vee x_3) \rightarrow (x_1 \vee x_2 \vee x_3 \vee S)$$

- Why this works:
 - (\Rightarrow) If $(x_1 \vee x_2 \vee x_3) = 1$

3SAT \leq_P NAE-4SAT

- We need a reduction f that takes 3SAT instance ϕ and converts it into NAE-4SAT instance ϕ'
 - If ϕ is satisfiable, ϕ' is NAE-satisfiable
 - If ϕ' is NAE-satisfiable, ϕ is satisfiable
 - Note that this must hold for every clause of ϕ , ϕ'
- Idea: Can we build a “gadget” for each clause of ϕ to enforce this condition?

$$(x_1 \vee x_2 \vee x_3) \rightarrow (x_1 \vee x_2 \vee x_3 \vee S)$$

- Why this works:
 - (\Rightarrow) If $(x_1 \vee x_2 \vee x_3) = 1$ at least one $x_i = 1$, so $(x_1 \vee x_2 \vee x_3 \vee S) = 1$. Set $S = 0$ to make it NAE-assignment

3SAT \leq_P NAE-4SAT

- We need a reduction f that takes 3SAT instance ϕ and converts it into NAE-4SAT instance ϕ'
 - If ϕ is satisfiable, ϕ' is NAE-satisfiable
 - If ϕ' is NAE-satisfiable, ϕ is satisfiable
 - Note that this must hold for every clause of ϕ , ϕ'
- Idea: Can we build a “gadget” for each clause of ϕ to enforce this condition?

$$(x_1 \vee x_2 \vee x_3) \rightarrow (x_1 \vee x_2 \vee x_3 \vee S)$$

- Why this works:
 - (\Rightarrow) If $(x_1 \vee x_2 \vee x_3) = 1$ at least one $x_i = 1$, so $(x_1 \vee x_2 \vee x_3 \vee S) = 1$.
Set $S = 0$ to make it NAE-assignment
 - (\Leftarrow) If $(x_1 \vee x_2 \vee x_3 \vee S) = 1$

3SAT \leq_P NAE-4SAT

- We need a reduction f that takes 3SAT instance ϕ and converts it into NAE-4SAT instance ϕ'
 - If ϕ is satisfiable, ϕ' is NAE-satisfiable
 - If ϕ' is NAE-satisfiable, ϕ is satisfiable
 - Note that this must hold for every clause of ϕ , ϕ'
- Idea: Can we build a “gadget” for each clause of ϕ to enforce this condition?

$$(x_1 \vee x_2 \vee x_3) \rightarrow (x_1 \vee x_2 \vee x_3 \vee S)$$

- Why this works:
 - (\Rightarrow) If $(x_1 \vee x_2 \vee x_3) = 1$ at least one $x_i = 1$, so $(x_1 \vee x_2 \vee x_3 \vee S) = 1$.
Set $S = 0$ to make it NAE-assignment
 - (\Leftarrow) If $(x_1 \vee x_2 \vee x_3 \vee S) = 1$
 - If $S = 0$, then at least one $x_i = 1$, so $(x_1 \vee x_2 \vee x_3) = 1$

3SAT \leq_P NAE-4SAT

- We need a reduction f that takes 3SAT instance ϕ and converts it into NAE-4SAT instance ϕ'
 - If ϕ is satisfiable, ϕ' is NAE-satisfiable
 - If ϕ' is NAE-satisfiable, ϕ is satisfiable
 - Note that this must hold for every clause of ϕ , ϕ'
- Idea: Can we build a “gadget” for each clause of ϕ to enforce this condition?

$$(x_1 \vee x_2 \vee x_3) \rightarrow (x_1 \vee x_2 \vee x_3 \vee S)$$

- Why this works:
 - (\Rightarrow) If $(x_1 \vee x_2 \vee x_3) = 1$ at least one $x_i = 1$, so $(x_1 \vee x_2 \vee x_3 \vee S) = 1$.
Set $S = 0$ to make it NAE-assignment
 - (\Leftarrow) If $(x_1 \vee x_2 \vee x_3 \vee S) = 1$
 - If $S = 0$, then at least one $x_i = 1$, so $(x_1 \vee x_2 \vee x_3) = 1$
 - If $S = 1$, then $(\overline{x_1} \vee \overline{x_2} \vee \overline{x_3} \vee 0)$ is also NAE-assignment. So, $(\overline{x_1} \vee \overline{x_2} \vee \overline{x_3}) = 1$

NAE-4SAT \leq_P NAE-3SAT

$\text{NAE-4SAT} \leq_P \text{NAE-3SAT}$

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE

NAE-4SAT \leq_P NAE-3SAT

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE
- Observations: Starting with $(x_1 \vee x_2 \vee x_3 \vee x_4)$

NAE-4SAT \leq_P NAE-3SAT

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE
- Observations: Starting with $(x_1 \vee x_2 \vee x_3 \vee x_4)$
 - We know that not all x_i have the same value

NAE-4SAT \leq_P NAE-3SAT

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE
- Observations: Starting with $(x_1 \vee x_2 \vee x_3 \vee x_4)$
 - We know that not all x_i have the same value
 - At least one of x_i is a 1 and one is a 0

NAE-4SAT \leq_P NAE-3SAT

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE
- Observations: Starting with $(x_1 \vee x_2 \vee x_3 \vee x_4)$
 - We know that not all x_i have the same value
 - At least one of x_i is a 1 and one is a 0
 - Idea: Let's split the variables into two clauses:

$$(x_1 \vee x_2 \vee \bar{x}_3) \wedge (x_1 \vee \bar{x}_2 \vee \bar{x}_3)$$

NAE-4SAT \leq_P NAE-3SAT

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE
- Observations: Starting with $(x_1 \vee x_2 \vee x_3 \vee x_4)$
 - We know that not all x_i have the same value
 - At least one of x_i is a 1 and one is a 0
 - Idea: Let's split the variables into two clauses:
$$(x_1 \vee x_2 \vee x_3 \vee x_4) \rightarrow (x_1 \vee x_2 \vee z_i) \wedge (x_3 \vee x_4 \vee \bar{z}_i)$$

NAE-4SAT \leq_P NAE-3SAT

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE
- Observations: Starting with $(x_1 \vee x_2 \vee x_3 \vee x_4)$
 - We know that not all x_i have the same value
 - At least one of x_i is a 1 and one is a 0
 - Idea: Let's split the variables into two clauses:
$$(x_1 \vee x_2 \vee x_3 \vee x_4) \rightarrow (x_1 \vee x_2 \vee z_i) \wedge (x_3 \vee x_4 \vee \bar{z}_i)$$
- Why this works:

NAE-4SAT \leq_P NAE-3SAT

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE
- Observations: Starting with $(x_1 \vee x_2 \vee x_3 \vee x_4)$
 - We know that not all x_i have the same value
 - At least one of x_i is a 1 and one is a 0
 - Idea: Let's split the variables into two clauses:
$$(x_1 \vee x_2 \vee x_3 \vee x_4) \rightarrow (x_1 \vee x_2 \vee z_i) \wedge (x_3 \vee x_4 \vee \bar{z}_i)$$

- Why this works:
 - (\Leftarrow) If $(x_1 \vee x_2 \vee z_i)$ and $(x_3 \vee x_4 \vee \bar{z}_i)$ are both NAE

NAE-4SAT \leq_P NAE-3SAT

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE
- Observations: Starting with $(x_1 \vee x_2 \vee x_3 \vee x_4)$
 - We know that not all x_i have the same value
 - At least one of x_i is a 1 and one is a 0
 - Idea: Let's split the variables into two clauses:
$$(x_1 \vee x_2 \vee x_3 \vee x_4) \rightarrow (x_1 \vee x_2 \vee z_i) \wedge (x_3 \vee x_4 \vee \bar{z}_i)$$
- Why this works:
 - (\Leftarrow) If $(x_1 \vee x_2 \vee z_i)$ and $(x_3 \vee x_4 \vee \bar{z}_i)$ are both NAE, then $(x_1 \vee x_2 \vee x_3 \vee x_4)$ is NAE
 - (\Rightarrow) If $(x_1 \vee x_2 \vee x_3 \vee x_4)$ is NAE

NAE-4SAT \leq_P NAE-3SAT

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE
- Observations: Starting with $(x_1 \vee x_2 \vee x_3 \vee x_4)$
 - We know that not all x_i have the same value
 - At least one of x_i is a 1 and one is a 0
 - Idea: Let's split the variables into two clauses:
$$(x_1 \vee x_2 \vee x_3 \vee x_4) \rightarrow (x_1 \vee x_2 \vee z_i) \wedge (x_3 \vee x_4 \vee \bar{z}_i)$$
- Why this works:
 - (\Leftarrow) If $(x_1 \vee x_2 \vee z_i)$ and $(x_3 \vee x_4 \vee \bar{z}_i)$ are both NAE, then $(x_1 \vee x_2 \vee x_3 \vee x_4)$ is NAE
 - (\Rightarrow) If $(x_1 \vee x_2 \vee x_3 \vee x_4)$ is NAE
 - If $x_1 \neq x_2$:

NAE-4SAT \leq_P NAE-3SAT

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE
- Observations: Starting with $(x_1 \vee x_2 \vee x_3 \vee x_4)$
 - We know that not all x_i have the same value
 - At least one of x_i is a 1 and one is a 0
 - Idea: Let's split the variables into two clauses:
$$(x_1 \vee x_2 \vee x_3 \vee x_4) \rightarrow (x_1 \vee x_2 \vee z_i) \wedge (x_3 \vee x_4 \vee \bar{z}_i)$$
- Why this works:
 - (\Leftarrow) If $(x_1 \vee x_2 \vee z_i)$ and $(x_3 \vee x_4 \vee \bar{z}_i)$ are both NAE, then $(x_1 \vee x_2 \vee x_3 \vee x_4)$ is NAE
 - (\Rightarrow) If $(x_1 \vee x_2 \vee x_3 \vee x_4)$ is NAE
 - If $x_1 \neq x_2$: Set $z_i = x_3$

NAE-4SAT \leq_P NAE-3SAT

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE
- Observations: Starting with $(x_1 \vee x_2 \vee x_3 \vee x_4)$
 - We know that not all x_i have the same value
 - At least one of x_i is a 1 and one is a 0
 - Idea: Let's split the variables into two clauses:
$$(x_1 \vee x_2 \vee x_3 \vee x_4) \rightarrow (x_1 \vee x_2 \vee z_i) \wedge (x_3 \vee x_4 \vee \bar{z}_i)$$
- Why this works:
 - (\Leftarrow) If $(x_1 \vee x_2 \vee z_i)$ and $(x_3 \vee x_4 \vee \bar{z}_i)$ are both NAE, then $(x_1 \vee x_2 \vee x_3 \vee x_4)$ is NAE
 - (\Rightarrow) If $(x_1 \vee x_2 \vee x_3 \vee x_4)$ is NAE
 - If $x_1 \neq x_2$: Set $z_i = x_3$
 - If $x_1 \neq x_3$:

NAE-4SAT \leq_P NAE-3SAT

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE
- Observations: Starting with $(x_1 \vee x_2 \vee x_3 \vee x_4)$
 - We know that not all x_i have the same value
 - At least one of x_i is a 1 and one is a 0
 - Idea: Let's split the variables into two clauses:
$$(x_1 \vee x_2 \vee x_3 \vee x_4) \rightarrow (x_1 \vee x_2 \vee z_i) \wedge (x_3 \vee x_4 \vee \bar{z}_i)$$
- Why this works:
 - (\Leftarrow) If $(x_1 \vee x_2 \vee z_i)$ and $(x_3 \vee x_4 \vee \bar{z}_i)$ are both NAE, then $(x_1 \vee x_2 \vee x_3 \vee x_4)$ is NAE
 - (\Rightarrow) If $(x_1 \vee x_2 \vee x_3 \vee x_4)$ is NAE
 - If $x_1 \neq x_2$: Set $z_i = x_3$
 - If $x_1 \neq x_3$: Set $z_i = x_3$

NAE-4SAT \leq_P NAE-3SAT

- Need a gadget to convert 4-CNF clause to CNF clauses that preserves NAE
- Observations: Starting with $(x_1 \vee x_2 \vee x_3 \vee x_4)$
 - We know that not all x_i have the same value
 - At least one of x_i is a 1 and one is a 0
 - Idea: Let's split the variables into two clauses:
$$(x_1 \vee x_2 \vee x_3 \vee x_4) \rightarrow (x_1 \vee x_2 \vee z_i) \wedge (x_3 \vee x_4 \vee \bar{z}_i)$$
- Why this works:
 - (\Leftarrow) If $(x_1 \vee x_2 \vee z_i)$ and $(x_3 \vee x_4 \vee \bar{z}_i)$ are both NAE, then $(x_1 \vee x_2 \vee x_3 \vee x_4)$ is NAE
 - (\Rightarrow) If $(x_1 \vee x_2 \vee x_3 \vee x_4)$ is NAE
 - If $x_1 \neq x_2$: Set $z_i = x_3$
 - If $x_1 \neq x_3$: Set $z_i = x_3$

Theorem

$$3SAT \leq_P NAE-4SAT \leq_P NAE-3SAT$$



3-Coloring


Definition

An undirected graph G is 3-colorable, if can assign colors $\{0, 1, 2\}$ to all nodes, such that no edges have the same color on both ends.

3-Coloring

Definition


An undirected graph G is 3-colorable, if we can assign colors $\{0, 1, 2\}$ to all nodes, such that no edges have the same color on both ends.




3-Coloring

Definition

An undirected graph G is 3-colorable, if we can assign colors $\{0, 1, 2\}$ to all nodes, such that no edges have the same color on both ends.




NAE-3SAT \leq_P 3-Coloring



NAE-3SAT \leq_P 3-Coloring


Gadgets:



- For each variable x_i :

NAE-3SAT \leq_P 3-Coloring

Gadgets:



Conclusions

- Many useful problems are \mathcal{NP} -complete

Conclusions

- Many useful problems are NP -complete
- But, as long as $\mathcal{P} \neq \text{NP}$, these are hard

Conclusions

- Many useful problems are NP -complete
- But, as long as $\mathcal{P} \neq \text{NP}$, these are hard
- Given a problem L , you should:

Conclusions

- Many useful problems are NP -complete
- But, as long as $\mathcal{P} \neq \text{NP}$, these are hard
- Given a problem L , you should:
 - ➊ Try to solve it ($L \in \mathcal{P}$)

Conclusions

- Many useful problems are NP -complete
- But, as long as $\mathcal{P} \neq \text{NP}$, these are hard
- Given a problem L , you should:
 - ➊ Try to solve it ($L \in \mathcal{P}$)
 - ➋ Try to prove NP -complete

Conclusions

- Many useful problems are NP -complete
- But, as long as $\mathcal{P} \neq \text{NP}$, these are hard
- Given a problem L , you should:
 - ① Try to solve it ($L \in \mathcal{P}$)
 - ② Try to prove NP -complete
- But, you must be careful

Conclusions

- Many useful problems are NP -complete
- But, as long as $\mathcal{P} \neq \text{NP}$, these are hard
- Given a problem L , you should:
 - ① Try to solve it ($L \in \mathcal{P}$)
 - ② Try to prove NP -complete
- But, you must be careful

3-Coloring is NP -complete, but 2-Coloring $\in \mathcal{P}$

Outline

1 Lecture 21 Review

2 More \mathcal{NP} -Complete Problems

3 co- \mathcal{NP}

Are All Problems in \mathcal{NP} ?

Question

Do all languages have poly-size proofs?

Are All Problems in \mathcal{NP} ?

Question

Do all languages have poly-size proofs?

Consider the following language:

UNSAT

$$\text{UNSAT} = \{\langle \phi \rangle \mid \phi \text{ is not satisfiable}\}$$

Are All Problems in \mathcal{NP} ?

Question

Do all languages have poly-size proofs?

Consider the following language:

UNSAT

$$\text{UNSAT} = \{\langle \phi \rangle \mid \phi \text{ is not satisfiable}\}$$

Problems like UNSAT are in co- \mathcal{NP}

\mathcal{P} , \mathcal{NP} and co- \mathcal{NP}

\mathcal{P}

$L \in \mathcal{P}$ if there exists poly-time DTM M s.t $M(x) = [x \in L]$

\mathcal{P} , \mathcal{NP} and co- \mathcal{NP}

\mathcal{P}

$L \in \mathcal{P}$ if there exists poly-time DTM M s.t $M(x) = [x \in L]$

\mathcal{NP}

$L \in \mathcal{NP}$ if there exists poly-time DTM V s.t. for $x \in L$ there exists a witness w s.t. $V(x, w) = 1$

\mathcal{P} , \mathcal{NP} and co- \mathcal{NP}

\mathcal{P}

$L \in \mathcal{P}$ if there exists poly-time DTM M s.t $M(x) = [x \in L]$

\mathcal{NP}

$L \in \mathcal{NP}$ if there exists poly-time DTM V s.t. for $x \in L$ there exists a witness w s.t. $V(x, w) = 1$

co- \mathcal{NP}

$L \in \text{co-}\mathcal{NP}$ if there exists poly-time DTM V s.t. for $x \in L$ for all w , $V(x, w) = 0$

\mathcal{P} , \mathcal{NP} and co- \mathcal{NP}

\mathcal{P}

$L \in \mathcal{P}$ if there exists poly-time DTM M s.t $M(x) = [x \in L]$

\mathcal{NP}

$L \in \mathcal{NP}$ if there exists poly-time DTM V s.t. for $x \in L$ there exists a witness w s.t. $V(x, w) = 1$

co- \mathcal{NP}

$L \in \text{co-}\mathcal{NP}$ if there exists poly-time DTM V s.t. for $x \in L$ for all w , $V(x, w) = 0$

Question:

Is $\mathcal{NP} = \text{co-}\mathcal{NP}$?

The Complexity Zoo

- There are many other complexity classes

The Complexity Zoo

- There are many other complexity classes
- We know some relationships between classes

The Complexity Zoo

- There are many other complexity classes
- We know some relationships between classes
- But, most big questions (e.g., $\mathcal{P} = \mathcal{NP}$, $\mathcal{NP} = \text{co-}\mathcal{NP}$, does PH collapse) are still not known!!!

The Complexity Zoo

- There are many other complexity classes
- We know some relationships between classes
- But, most big questions (e.g., $\mathcal{P} = \mathcal{NP}$, $\mathcal{NP} = \text{co-}\mathcal{NP}$, does PH collapse) are still not known!!!

Complexity Zoo

The complexity zoo (https://complexityzoo.net/Complexity_Zoo) now has 546 complexity classes.