

Foundations of Computing

Lecture 7

Arkady Yerukhimovich

February 7, 2023

- 1 Lecture 6 Review
- 2 The Pumping Lemma for Regular Languages
- 3 Using the Pumping Lemma
- 4 Using Closure Properties

Lecture 6 Review

- Nonregular languages
- Proving The NFA pumping lemma

Lecture 6 Review

- Nonregular languages
- Proving The NFA pumping lemma

Today

Using the pumping lemma to prove languages are not regular.

HW2 Problem 4

Let L be a regular language, prove that the following languages are regular.

- ① $NOPREFIX(L) = \{w \in L \mid \text{no proper prefix of } w \text{ is a member of } L\}$
- ② $NOEXTEND(L) = \{w \in L \mid w \text{ is not a proper prefix of any string in } L\}$

HW2 Problem 4

Let L be a regular language, prove that the following languages are regular.

- ① $NOPREFIX(L) = \{w \in L \mid \text{no proper prefix of } w \text{ is a member of } L\}$
- ② $NOEXTEND(L) = \{w \in L \mid w \text{ is not a proper prefix of any string in } L\}$

Example:

- $L = \{00, 11, 001, 101\}$
- $NOPREFIX(L) = \{00, 11, 101\}$
- $NOEXTEND(L) = \{11, 001, 101\}$

HW2 Problem 4

Let L be a regular language, prove that the following languages are regular.

① $NOPREFIX(L) = \{w \in L \mid \text{no proper prefix of } w \text{ is a member of } L\}$

HW2 Problem 4

Let L be a regular language, prove that the following languages are regular.

- 1 $NOPREFIX(L) = \{w \in L \mid \text{no proper prefix of } w \text{ is a member of } L\}$
- 2 $NOEXTEND(L) = \{w \in L \mid w \text{ is not a proper prefix of any string in } L\}$

Outline

- 1 Lecture 6 Review
- 2 The Pumping Lemma for Regular Languages**
- 3 Using the Pumping Lemma
- 4 Using Closure Properties

The Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three pieces $w = xyz$ satisfying:

The Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three pieces $w = xyz$ satisfying:

- 1 For each $i \geq 0$, $xy^iz \in L$

The Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three pieces $w = xyz$ satisfying:

- 1 For each $i \geq 0$, $xy^iz \in L$
- 2 $|y| > 0$, and

The Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three pieces $w = xyz$ satisfying:

- 1 For each $i \geq 0$, $xy^iz \in L$
- 2 $|y| > 0$, and
- 3 $|xy| \leq p$

The Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three pieces $w = xyz$ satisfying:

- 1 For each $i \geq 0$, $xy^iz \in L$
- 2 $|y| > 0$, and
- 3 $|xy| \leq p$

- We saw how to prove the pumping lemma last week
- Today we will learn how to use it

Outline

- 1 Lecture 6 Review
- 2 The Pumping Lemma for Regular Languages
- 3 Using the Pumping Lemma**
- 4 Using Closure Properties

The Proof Procedure

To use the pumping lemma to prove that L is not regular, we do the following:

The Proof Procedure

To use the pumping lemma to prove that L is not regular, we do the following:

- 1 Assume that L is regular

The Proof Procedure

To use the pumping lemma to prove that L is not regular, we do the following:

- 1 Assume that L is regular
- 2 Use pumping lemma to guarantee pumping length p , s.t. all w with $|w| > p$ can be pumped

The Proof Procedure

To use the pumping lemma to prove that L is not regular, we do the following:

- 1 Assume that L is regular
- 2 Use pumping lemma to guarantee pumping length p , s.t. all w with $|w| > p$ can be pumped
- 3 Choose $w \in L$ with $|w| \geq p$

The Proof Procedure

To use the pumping lemma to prove that L is not regular, we do the following:

- 1 Assume that L is regular
- 2 Use pumping lemma to guarantee pumping length p , s.t. all w with $|w| > p$ can be pumped
- 3 Choose $w \in L$ with $|w| \geq p$
- 4 Demonstrate that w cannot be pumped
 - For each possible division $w = xyz$ (with $|y| > 0$ and $|xy| \leq p$), find an integer i such that $xy^iz \notin L$

The Proof Procedure

To use the pumping lemma to prove that L is not regular, we do the following:

- ① Assume that L is regular
- ② Use pumping lemma to guarantee pumping length p , s.t. all w with $|w| > p$ can be pumped
- ③ Choose $w \in L$ with $|w| \geq p$
- ④ Demonstrate that w cannot be pumped
 - For each possible division $w = xyz$ (with $|y| > 0$ and $|xy| \leq p$), find an integer i such that $xy^iz \notin L$
- ⑤ Contradiction!!!

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p \in L$

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p \in L$
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p \in L$
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- 4 Complete proof by considering all possible values for y

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p \in L$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- ④ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xyyz$ has more 0s than 1s, so $w \notin L$

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p \in L$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- ④ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xyyz$ has more 0s than 1s, so $w \notin L$
 - y consists of only 1s – then $xyyz$ has more 1s than 0s, so $w \notin L$

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p \in L$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- ④ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xyyz$ has more 0s than 1s, so $w \notin L$
 - y consists of only 1s – then $xyyz$ has more 1s than 0s, so $w \notin L$
 - y consists of both 0s and 1s – then $xyyz$ has 0s alternating with 1s more than once, so $w \notin L$

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p \in L$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- ④ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xyyz$ has more 0s than 1s, so $w \notin L$
 - y consists of only 1s – then $xyyz$ has more 1s than 0s, so $w \notin L$
 - y consists of both 0s and 1s – then $xyyz$ has 0s alternating with 1s more than once, so $w \notin L$
- ⑤ Contradiction – hence, L is not regular

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p \in L$

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p \in L$
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p \in L$
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- 4 Problem: We have to consider all possible splits $w = xyz$, but if $y = 0^m 1^m$, then w can be pumped – not leading to contradiction

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p \in L$
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- 4 Problem: We have to consider all possible splits $w = xyz$, but if $y = 0^m 1^m$, then w can be pumped – not leading to contradiction
- 5 Solution: Use condition that $|xy| \leq p$

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p \in L$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- ④ Problem: We have to consider all possible splits $w = xyz$, but if $y = 0^m 1^m$, then w can be pumped – not leading to contradiction
- ⑤ Solution: Use condition that $|xy| \leq p$
 - Since $w = 0^p 1^p$ and $|xy| \leq p$, we know that y must be in first p symbols

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p \in L$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- ④ Problem: We have to consider all possible splits $w = xyz$, but if $y = 0^m 1^m$, then w can be pumped – not leading to contradiction
- ⑤ Solution: Use condition that $|xy| \leq p$
 - Since $w = 0^p 1^p$ and $|xy| \leq p$, we know that y must be in first p symbols
 - But, this means that y must be all 0s

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p \in L$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- ④ Problem: We have to consider all possible splits $w = xyz$, but if $y = 0^m 1^m$, then w can be pumped – not leading to contradiction
- ⑤ Solution: Use condition that $|xy| \leq p$
 - Since $w = 0^p 1^p$ and $|xy| \leq p$, we know that y must be in first p symbols
 - But, this means that y must be all 0s
- ⑥ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xyyz$ has more 0s than 1s, so $w \notin L$

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p \in L$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- ④ Problem: We have to consider all possible splits $w = xyz$, but if $y = 0^m 1^m$, then w can be pumped – not leading to contradiction
- ⑤ Solution: Use condition that $|xy| \leq p$
 - Since $w = 0^p 1^p$ and $|xy| \leq p$, we know that y must be in first p symbols
 - But, this means that y must be all 0s
- ⑥ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xyyz$ has more 0s than 1s, so $w \notin L$
- ⑦ Contradiction – hence, L is not regular

Example 2 – A Much Simpler Proof

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Using closure properties:

Example 2 – A Much Simpler Proof

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Using closure properties:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is not regular

Example 2 – A Much Simpler Proof

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Using closure properties:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is not regular
- 2 Observe that $L_1 = L \cap 0^* 1^*$

Example 2 – A Much Simpler Proof

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Using closure properties:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is not regular
- 2 Observe that $L_1 = L \cap 0^* 1^*$
- 3 Easy to see that $0^* 1^*$ is regular

Example 2 – A Much Simpler Proof

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Using closure properties:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is not regular
- 2 Observe that $L_1 = L \cap 0^* 1^*$
- 3 Easy to see that $0^* 1^*$ is regular
- 4 Since regular languages are closed under \cap , if L is regular then L_1 must be regular

Example 2 – A Much Simpler Proof

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Using closure properties:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is not regular
- 2 Observe that $L_1 = L \cap 0^* 1^*$
- 3 Easy to see that $0^* 1^*$ is regular
- 4 Since regular languages are closed under \cap , if L is regular then L_1 must be regular
- 5 Since we know L_1 is nonregular, this means that L must be nonregular

Example 3

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- 1 Assume L is regular, and let p be the pumping length

Example 3

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- 1 Assume L is regular, and let p be the pumping length
- 2 What w should we choose?

Example 3

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- 1 Assume L is regular, and let p be the pumping length
- 2 What w should we choose?
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i

Example 3

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- 1 Assume L is regular, and let p be the pumping length
- 2 What w should we choose?
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- 4 Recall that our goal is to prove that above is not true. Specifically, we want

Example 3

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- 1 Assume L is regular, and let p be the pumping length
- 2 What w should we choose?
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- 4 Recall that our goal is to prove that above is not true. Specifically, we want $xy^i z = 0^{m'} 1^{n'}$ with $m' = n'$.

Example 3

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- 1 Assume L is regular, and let p be the pumping length
- 2 What w should we choose?
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- 4 Recall that our goal is to prove that above is not true. Specifically, we want $xy^i z = 0^{m'} 1^{n'}$ with $m' = n'$.
- 5 Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{m-(\alpha+\beta)} 1^n$

Example 3

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- 1 Assume L is regular, and let p be the pumping length
- 2 What w should we choose?
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- 4 Recall that our goal is to prove that above is not true. Specifically, we want $xy^i z = 0^{m'} 1^{n'}$ with $m' = n'$.
- 5 Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{m-(\alpha+\beta)} 1^n$
- 6 Consider what happens when we pump k times:

Example 3

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- 1 Assume L is regular, and let p be the pumping length
- 2 What w should we choose?
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- 4 Recall that our goal is to prove that above is not true. Specifically, we want $xy^i z = 0^{m'} 1^{n'}$ with $m' = n'$.
- 5 Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{m-(\alpha+\beta)} 1^n$
- 6 Consider what happens when we pump k times: $xy^k z = 0^{\alpha+k\beta+m-(\alpha+\beta)} 1^n$.

Example 3

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- ① Assume L is regular, and let p be the pumping length
- ② What w should we choose?
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- ④ Recall that our goal is to prove that above is not true. Specifically, we want $xy^i z = 0^{m'} 1^{n'}$ with $m' = n'$.
- ⑤ Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{m-(\alpha+\beta)} 1^n$
- ⑥ Consider what happens when we pump k times: $xy^k z = 0^{\alpha+k\beta+m-(\alpha+\beta)} 1^n$.
We need $\alpha + k\beta + m - (\alpha + \beta) = m + (k - 1)\beta = n$ for a contradiction

Example 3

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- ① Assume L is regular, and let p be the pumping length
- ② What w should we choose?
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- ④ Recall that our goal is to prove that above is not true. Specifically, we want $xy^i z = 0^{m'} 1^{n'}$ with $m' = n'$.
- ⑤ Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{m-(\alpha+\beta)} 1^n$
- ⑥ Consider what happens when we pump k times: $xy^k z = 0^{\alpha+k\beta+m-(\alpha+\beta)} 1^n$.
We need $\alpha + k\beta + m - (\alpha + \beta) = m + (k - 1)\beta = n$ for a contradiction
Equivalently, we need $k = (n - m)/\beta + 1$ to be an integer

Example 3

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- ① Assume L is regular, and let p be the pumping length
- ② What w should we choose?
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- ④ Recall that our goal is to prove that above is not true. Specifically, we want $xy^i z = 0^{m'} 1^{n'}$ with $m' = n'$.
- ⑤ Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{m-(\alpha+\beta)} 1^n$
- ⑥ Consider what happens when we pump k times: $xy^k z = 0^{\alpha+k\beta+m-(\alpha+\beta)} 1^n$.
We need $\alpha + k\beta + m - (\alpha + \beta) = m + (k - 1)\beta = n$ for a contradiction
Equivalently, we need $k = (n - m)/\beta + 1$ to be an integer
- ⑦ We only know $\beta \leq p$, how can we guarantee $(n - m)$ is divisible by β ?

Example 3

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- ① Assume L is regular, and let p be the pumping length
- ② What w should we choose?
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$ for all i
- ④ Recall that our goal is to prove that above is not true. Specifically, we want $xy^i z = 0^{m'} 1^{n'}$ with $m' = n'$.
- ⑤ Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{m-(\alpha+\beta)} 1^n$
- ⑥ Consider what happens when we pump k times: $xy^k z = 0^{\alpha+k\beta+m-(\alpha+\beta)} 1^n$.
We need $\alpha + k\beta + m - (\alpha + \beta) = m + (k - 1)\beta = n$ for a contradiction
Equivalently, we need $k = (n - m)/\beta + 1$ to be an integer
- ⑦ We only know $\beta \leq p$, how can we guarantee $(n - m)$ is divisible by β ?
- ⑧ Set $n = 2p!$, $m = p!$, can guarantee $(n - m) = p!$ is divisible by β , so there is k s.t. $xy^k z \notin L$

Hints for Using the Pumping Lemma

To use the pumping lemma, need to do the following

- Identify what it means for $x \notin L$
- Choose w such that any valid split xyz can lead to a contradiction
- Prove that $w' = xy^kz \notin L$ for some k

Choosing w is often tricky, requires intuition and some trial and error.

Outline

- 1 Lecture 6 Review
- 2 The Pumping Lemma for Regular Languages
- 3 Using the Pumping Lemma
- 4 Using Closure Properties

Using Closure Properties of Regular Languages

We have seen a number of closure properties of REs

- ① Closure under complement: \overline{L} is regular if L is
- ② Closure under union: $L_1 \cup L_2$ is regular if L_1, L_2 are
- ③ Closure under intersection: $L_1 \cap L_2$ is regular if L_1, L_2 are
- ④ Closure under reversal: L^R is regular if L is
- ⑤ NOPREFIX, NOEXTEND
- ⑥ There are many more (e.g., set difference, cross product, ...)

Using Closure Properties of Regular Languages

We have seen a number of closure properties of REs

- 1 Closure under complement: \overline{L} is regular if L is
- 2 Closure under union: $L_1 \cup L_2$ is regular if L_1, L_2 are
- 3 Closure under intersection: $L_1 \cap L_2$ is regular if L_1, L_2 are
- 4 Closure under reversal: L^R is regular if L is
- 5 NOPREFIX, NOEXTEND
- 6 There are many more (e.g., set difference, cross product, ...)

Important

- It is much easier to prove non-regularity using closure properties
- Try this first before you turn to pumping lemma

Exercise

Prove that the following language is nonregular:

$$L = \{0^i 1^j 2^i 3^j \mid i, j > 0\}$$

What's Next?

- We will add (a little) memory to our machines to recognize a richer class of languages