

Foundations of Computing

Lecture 6

Arkady Yerukhimovich

January 30, 2025

Outline

- 1 Lecture 5 Review
- 2 A Non-regular Language
- 3 The Pumping Lemma for Regular Languages
- 4 Proving the Pumping Lemma
- 5 Using the Pumping Lemma

Lecture 5 Review

- Regular expressions
- Equivalence of regular expressions and NFAs/DFAs

Quiz Solutions

For each of the following languages over $\Sigma = \{a, b\}$, give two strings that are in the language and two strings not in the language.

① $a^* \cup b^*$

② $(aa \cup bb)^*$

③ $\Sigma^* a \Sigma^* b \Sigma^* a \Sigma^*$

Outline

- 1 Lecture 5 Review
- 2 A Non-regular Language**
- 3 The Pumping Lemma for Regular Languages
- 4 Proving the Pumping Lemma
- 5 Using the Pumping Lemma

What We Know So Far

The following four things are equivalent:

- 1 Regular languages
- 2 Languages decided by DFAs
- 3 Languages decided by NFAs
- 4 Languages described by regular expressions

What We Know So Far

The following four things are equivalent:

- 1 Regular languages
- 2 Languages decided by DFAs
- 3 Languages decided by NFAs
- 4 Languages described by regular expressions

Are all languages regular?

What We Know So Far

The following four things are equivalent:

- 1 Regular languages
- 2 Languages decided by DFAs
- 3 Languages decided by NFAs
- 4 Languages described by regular expressions

Are all languages regular?

Today we will see that there are languages that are not regular

The F in DFA/NFA

The F in DFA/NFA

Important

In a *Finite* Automaton, the number of states is **finite**

The F in DFA/NFA

Important

In a *Finite* Automaton, the number of states is **finite**

This means that:

- The number of states is fixed independently of the input size

The F in DFA/NFA

Important

In a *Finite* Automaton, the number of states is **finite**

This means that:

- The number of states is fixed independently of the input size
- An automaton must be able to process strings w s.t. $|w| > |Q|$

The F in DFA/NFA

Important

In a *Finite* Automaton, the number of states is **finite**

This means that:

- The number of states is fixed independently of the input size
- An automaton must be able to process strings w s.t. $|w| > |Q|$
- Thus, a finite automaton cannot store its whole input

A Nonregular Language

Consider the following language:

$$L = \{0^n 1^n \mid n \geq 0\}$$

Let's try to build a DFA for L :

A Nonregular Language

Consider the following language:

$$L = \{0^n 1^n \mid n \geq 0\}$$

Let's try to build a DFA for L :

The Problem

We need to count the number of 0s, but this is unbounded so can't have a state for each value

The Need for a Proof

What we just saw

Intuition: An NFA/DFA cannot count unbounded inputs

The Need for a Proof

What we just saw

Intuition: An NFA/DFA cannot count unbounded inputs

Why isn't this a proof?

The Need for a Proof

What we just saw

Intuition: An NFA/DFA cannot count unbounded inputs

Why isn't this a proof?

Consider the following language:

$L' = \{w \mid w \text{ has an equal number of occurrences of } 01 \text{ and } 10 \text{ as substrings}\}$

General Proof Structure

We will prove that a language L is not regular by contradiction

General Proof Structure

We will prove that a language L is not regular by contradiction

- 1 Assume L is regular – there is a NFA/DFA M accepting it

General Proof Structure

We will prove that a language L is not regular by contradiction

- 1 Assume L is regular – there is a NFA/DFA M accepting it
- 2 Pick a particular string $w \in L$

General Proof Structure

We will prove that a language L is not regular by contradiction

- 1 Assume L is regular – there is a NFA/DFA M accepting it
- 2 Pick a particular string $w \in L$
- 3 Show that if $M(w) = 1$ then there exists a string $w' \notin L$ s.t. $M(w') = 1$

General Proof Structure

We will prove that a language L is not regular by contradiction

- 1 Assume L is regular – there is a NFA/DFA M accepting it
- 2 Pick a particular string $w \in L$
- 3 Show that if $M(w) = 1$ then there exists a string $w' \notin L$ s.t. $M(w') = 1$
- 4 Conclude that L is not regular since any M that accepts all strings in L must also accept strings not in L

Outline

- 1 Lecture 5 Review
- 2 A Non-regular Language
- 3 The Pumping Lemma for Regular Languages**
- 4 Proving the Pumping Lemma
- 5 Using the Pumping Lemma

The Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three substrings $w = xyz$ satisfying:

The Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three substrings $w = xyz$ satisfying:

- 1 For each $i \geq 0$, $xy^iz \in L$

The Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three substrings $w = xyz$ satisfying:

- 1 For each $i \geq 0$, $xy^iz \in L$
- 2 $|y| > 0$, and

The Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three substrings $w = xyz$ satisfying:

- 1 For each $i \geq 0$, $xy^iz \in L$
- 2 $|y| > 0$, and
- 3 $|xy| \leq p$

The Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three substrings $w = xyz$ satisfying:

- 1 For each $i \geq 0$, $xy^iz \in L$
- 2 $|y| > 0$, and
- 3 $|xy| \leq p$

Let's write this using math:

The Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three substrings $w = xyz$ satisfying:

- 1 For each $i \geq 0$, $xy^iz \in L$
- 2 $|y| > 0$, and
- 3 $|xy| \leq p$

Let's write this using math:

Next steps:

- 1 Prove the pumping lemma
- 2 Show how to use the pumping lemma to prove languages nonregular

Outline

- 1 Lecture 5 Review
- 2 A Non-regular Language
- 3 The Pumping Lemma for Regular Languages
- 4 Proving the Pumping Lemma**
- 5 Using the Pumping Lemma

The Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three substrings $w = xyz$ satisfying:

- 1 For each $i \geq 0$, $xy^iz \in L$
- 2 $|y| > 0$, and
- 3 $|xy| \leq p$

Proof Intuition

Let $M = (Q, \Sigma, \delta, q_1, F)$ be a DFA that recognizes L and let $p = |Q|$

Proof Intuition

Let $M = (Q, \Sigma, \delta, q_1, F)$ be a DFA that recognizes L and let $p = |Q|$

- If for all $w \in L$, $|w| < p$, then we are done

Proof Intuition

Let $M = (Q, \Sigma, \delta, q_1, F)$ be a DFA that recognizes L and let $p = |Q|$

- If for all $w \in L$, $|w| < p$, then we are done
- Suppose $w \in L$ s.t. $|w| = n \geq p$

Proof Intuition


Let $M = (Q, \Sigma, \delta, q_1, F)$ be a DFA that recognizes L and let $p = |Q|$

- If for all $w \in L$, $|w| < p$, then we are done
- Suppose $w \in L$ s.t. $|w| = n \geq p$
 - If we run $M(w)$, it will go through $n + 1$ states – since it transitions on each symbol of w


Proof Intuition

Let $M = (Q, \Sigma, \delta, q_1, F)$ be a DFA that recognizes L and let $p = |Q|$

- If for all $w \in L$, $|w| < p$, then we are done
- Suppose $w \in L$ s.t. $|w| = n \geq p$
 - If we run $M(w)$, it will go through $n + 1$ states – since it transitions on each symbol of w
 - Since $n + 1 > p$, there must be some state that is visited twice




Proof Intuition



Divide $w = xyz$ as follows:


Proof Intuition



Divide $w = xyz$ as follows:

- x is the part of w until $M(w)$ visits q_5


Proof Intuition



Divide $w = xyz$ as follows:

- x is the part of w until $M(w)$ visits q_5
- y is the part of w between the two visits to q_5

Proof Intuition



Divide $w = xyz$ as follows:

- x is the part of w until $M(w)$ visits q_5
- y is the part of w between the two visits to q_5
- z is the part of w after the second visit to q_5

Proof Intuition

We divided w into xyz as follows:

- x is the part of w until $M(w)$ visits q_5
- y is the part of w between the two visits to q_5
- z is the part of w after the second visit to q_5

Proof Intuition

We divided w into xyz as follows:

- x is the part of w until $M(w)$ visits q_5
- y is the part of w between the two visits to q_5
- z is the part of w after the second visit to q_5

We show that the properties of the pumping lemma hold:

Proof Intuition

We divided w into xyz as follows:

- x is the part of w until $M(w)$ visits q_5
- y is the part of w between the two visits to q_5
- z is the part of w after the second visit to q_5

We show that the properties of the pumping lemma hold:

- 1 For each $i \geq 0$, $xy^iz \in L$

Proof Intuition

We divided w into xyz as follows:

- x is the part of w until $M(w)$ visits q_5
- y is the part of w between the two visits to q_5
- z is the part of w after the second visit to q_5

We show that the properties of the pumping lemma hold:

- 1 For each $i \geq 0$, $xy^iz \in L$

Proof: y takes M from q_5 back to q_5 . So, if you run $M(xyyz)$, it would just run this cycle twice...

Proof Intuition

We divided w into xyz as follows:

- x is the part of w until $M(w)$ visits q_5
- y is the part of w between the two visits to q_5
- z is the part of w after the second visit to q_5

We show that the properties of the pumping lemma hold:

- 1 For each $i \geq 0$, $xy^iz \in L$

Proof: y takes M from q_5 back to q_5 . So, if you run $M(xyyz)$, it would just run this cycle twice...

- 2 $|y| > 0$

Proof Intuition

We divided w into xyz as follows:

- x is the part of w until $M(w)$ visits q_5
- y is the part of w between the two visits to q_5
- z is the part of w after the second visit to q_5

We show that the properties of the pumping lemma hold:

- 1 For each $i \geq 0$, $xy^iz \in L$

Proof: y takes M from q_5 back to q_5 . So, if you run $M(xyyz)$, it would just run this cycle twice...

- 2 $|y| > 0$

Proof: ≥ 1 character of w must be between the two visits of q_5

Proof Intuition

We divided w into xyz as follows:

- x is the part of w until $M(w)$ visits q_5
- y is the part of w between the two visits to q_5
- z is the part of w after the second visit to q_5

We show that the properties of the pumping lemma hold:

- 1 For each $i \geq 0$, $xy^iz \in L$

Proof: y takes M from q_5 back to q_5 . So, if you run $M(xyyz)$, it would just run this cycle twice...

- 2 $|y| > 0$

Proof: ≥ 1 character of w must be between the two visits of q_5

- 3 $|xy| \leq p$

Proof Intuition

We divided w into xyz as follows:

- x is the part of w until $M(w)$ visits q_5
- y is the part of w between the two visits to q_5
- z is the part of w after the second visit to q_5

We show that the properties of the pumping lemma hold:

- 1 For each $i \geq 0$, $xy^iz \in L$

Proof: y takes M from q_5 back to q_5 . So, if you run $M(xyyz)$, it would just run this cycle twice...

- 2 $|y| > 0$

Proof: ≥ 1 character of w must be between the two visits of q_5

- 3 $|xy| \leq p$

Proof: if q_5 is the first repetition in $M(w)$, then this repetition must occur in the first $p + 1$ states, hence $|xy| \leq p$

Making it Formal

Let $M = (Q, \Sigma, \delta, q_1, F)$ be a DFA recognizing L

Making it Formal

Let $M = (Q, \Sigma, \delta, q_1, F)$ be a DFA recognizing L

- Let $w = w_1 w_2 \dots w_n \in L$

Making it Formal

Let $M = (Q, \Sigma, \delta, q_1, F)$ be a DFA recognizing L

- Let $w = w_1 w_2 \dots w_n \in L$
- Let r_1, r_2, \dots, r_{n+1} be the sequence of states visited by $M(w)$
($r_{i+1} = \delta(r_i, w_i)$)

Making it Formal

Let $M = (Q, \Sigma, \delta, q_1, F)$ be a DFA recognizing L

- Let $w = w_1 w_2 \dots w_n \in L$
- Let r_1, r_2, \dots, r_{n+1} be the sequence of states visited by $M(w)$
($r_{i+1} = \delta(r_i, w_i)$)
- In first $p + 1$ elements of this sequence there must be a repeated element $r_j = r_k$, so $k \leq p + 1$

Making it Formal

Let $M = (Q, \Sigma, \delta, q_1, F)$ be a DFA recognizing L

- Let $w = w_1 w_2 \dots w_n \in L$
- Let r_1, r_2, \dots, r_{n+1} be the sequence of states visited by $M(w)$
($r_{i+1} = \delta(r_i, w_i)$)
- In first $p + 1$ elements of this sequence there must be a repeated element $r_j = r_k$, so $k \leq p + 1$
- Let $x = w_1 \dots w_{j-1}$, $y = w_j \dots w_{k-1}$, and $z = w_k \dots w_n$

Making it Formal

Let $M = (Q, \Sigma, \delta, q_1, F)$ be a DFA recognizing L

- Let $w = w_1 w_2 \dots w_n \in L$
- Let r_1, r_2, \dots, r_{n+1} be the sequence of states visited by $M(w)$ ($r_{i+1} = \delta(r_i, w_i)$)
- In first $p + 1$ elements of this sequence there must be a repeated element $r_j = r_k$, so $k \leq p + 1$
- Let $x = w_1 \dots w_{j-1}$, $y = w_j \dots w_{k-1}$, and $z = w_k \dots w_n$

Observe that:

- x takes M from $r_1 = q_1$ to r_j , y takes M from r_j to r_k , and z takes M from r_k to r_{n+1} , which is an accept state. So, M must accept $xy^i z$ for $i \geq 0$

Making it Formal

Let $M = (Q, \Sigma, \delta, q_1, F)$ be a DFA recognizing L

- Let $w = w_1 w_2 \dots w_n \in L$
- Let r_1, r_2, \dots, r_{n+1} be the sequence of states visited by $M(w)$
($r_{i+1} = \delta(r_i, w_i)$)
- In first $p + 1$ elements of this sequence there must be a repeated element $r_j = r_k$, so $k \leq p + 1$
- Let $x = w_1 \dots w_{j-1}$, $y = w_j \dots w_{k-1}$, and $z = w_k \dots w_n$

Observe that:

- x takes M from $r_1 = q_1$ to r_j , y takes M from r_j to r_k , and z takes M from r_k to r_{n+1} , which is an accept state. So, M must accept $xy^i z$ for $i \geq 0$
- $j \neq k$ so, $|y| > 0$

Making it Formal

Let $M = (Q, \Sigma, \delta, q_1, F)$ be a DFA recognizing L

- Let $w = w_1 w_2 \dots w_n \in L$
- Let r_1, r_2, \dots, r_{n+1} be the sequence of states visited by $M(w)$ ($r_{i+1} = \delta(r_i, w_i)$)
- In first $p + 1$ elements of this sequence there must be a repeated element $r_j = r_k$, so $k \leq p + 1$
- Let $x = w_1 \dots w_{j-1}$, $y = w_j \dots w_{k-1}$, and $z = w_k \dots w_n$

Observe that:

- x takes M from $r_1 = q_1$ to r_j , y takes M from r_j to r_k , and z takes M from r_k to r_{n+1} , which is an accept state. So, M must accept $xy^i z$ for $i \geq 0$
- $j \neq k$ so, $|y| > 0$
- $k \leq p + 1$, so $|xy| \leq p$

The Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three substrings $w = xyz$ satisfying:

- 1 For each $i \geq 0$, $xy^iz \in L$
- 2 $|y| > 0$, and
- 3 $|xy| \leq p$

Mathematically:

$$\forall w \in L, |w| \geq p \exists \text{ substrings } w = xyz \text{ s.t. } \forall i \geq 0, xy^iz \in L$$

Let's negate this:

$$\exists w \in L, |w| \geq p \forall \text{ substrings } w = xyz \exists i \geq 0, \text{ s.t. } xy^iz \notin L$$

Outline

- 1 Lecture 5 Review
- 2 A Non-regular Language
- 3 The Pumping Lemma for Regular Languages
- 4 Proving the Pumping Lemma
- 5 Using the Pumping Lemma

Negated Pumping Lemma

$$\exists w \in L, |w| \geq p \forall \text{ substrings } w = xyz \exists i \geq 0, \text{ s.t. } xy^iz \notin L$$

To use the pumping lemma to prove that L is not regular, we do the following:

Negated Pumping Lemma

$$\exists w \in L, |w| \geq p \forall \text{ substrings } w = xyz \exists i \geq 0, \text{ s.t. } xy^iz \notin L$$

To use the pumping lemma to prove that L is not regular, we do the following:

- 1 Assume that L is regular

Negated Pumping Lemma

$$\exists w \in L, |w| \geq p \forall \text{ substrings } w = xyz \exists i \geq 0, \text{ s.t. } xy^iz \notin L$$

To use the pumping lemma to prove that L is not regular, we do the following:

- 1 Assume that L is regular
- 2 By pumping lemma there exists pumping length p , s.t. all w with $|w| > p$ can be pumped

Negated Pumping Lemma

$$\exists w \in L, |w| \geq p \ \forall \text{ substrings } w = xyz \ \exists i \geq 0, \text{ s.t. } xy^iz \notin L$$

To use the pumping lemma to prove that L is not regular, we do the following:

- 1 Assume that L is regular
- 2 By pumping lemma there exists pumping length p , s.t. all w with $|w| \geq p$ can be pumped
- 3 Choose a particular $w \in L$ with $|w| \geq p$

Negated Pumping Lemma

$$\exists w \in L, |w| \geq p \ \forall \text{ substrings } w = xyz \ \exists i \geq 0, \text{ s.t. } xy^iz \notin L$$

To use the pumping lemma to prove that L is not regular, we do the following:

- ① Assume that L is regular
- ② By pumping lemma there exists pumping length p , s.t. all w with $|w| \geq p$ can be pumped
- ③ Choose a particular $w \in L$ with $|w| \geq p$
- ④ Demonstrate that w cannot be pumped:
 - For each possible division $w = xyz$, find an i such that $xy^iz \notin L$

Negated Pumping Lemma

$$\exists w \in L, |w| \geq p \ \forall \text{ substrings } w = xyz \ \exists i \geq 0, \text{ s.t. } xy^iz \notin L$$

To use the pumping lemma to prove that L is not regular, we do the following:

- 1 Assume that L is regular
- 2 By pumping lemma there exists pumping length p , s.t. all w with $|w| \geq p$ can be pumped
- 3 Choose a particular $w \in L$ with $|w| \geq p$
- 4 Demonstrate that w cannot be pumped:
 - For each possible division $w = xyz$, find an i such that $xy^iz \notin L$
- 5 Contradiction – Pumping lemma must hold for any regular L

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p$

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p$
- 3 By pumping lemma, w can be split into xyz s.t. $xy^i z \in L$

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p$
- 3 By pumping lemma, w can be split into xyz s.t. $xy^i z \in L$
- 4 Complete proof by considering all possible values for y

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p$
- ③ By pumping lemma, w can be split into xyz s.t. $xy^i z \in L$
- ④ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xyyz$ has more 0s than 1s, so $w \notin L$

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p$
- ③ By pumping lemma, w can be split into xyz s.t. $xy^i z \in L$
- ④ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xxyz$ has more 0s than 1s, so $w \notin L$
 - y consists of only 1s – then $xxyz$ has more 1s than 0s, so $w \notin L$

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p$
- ③ By pumping lemma, w can be split into xyz s.t. $xy^i z \in L$
- ④ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xxyz$ has more 0s than 1s, so $w \notin L$
 - y consists of only 1s – then $xxyz$ has more 1s than 0s, so $w \notin L$
 - y consists of both 0s and 1s – then $xxyz$ has 0s alternating with 1s more than once, so $w \notin L$

Example 1

Consider $L = \{0^n 1^n \mid n \geq 0\}$, prove L is not regular.

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p$
- ③ By pumping lemma, w can be split into xyz s.t. $xy^i z \in L$
- ④ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xxyz$ has more 0s than 1s, so $w \notin L$
 - y consists of only 1s – then $xxyz$ has more 1s than 0s, so $w \notin L$
 - y consists of both 0s and 1s – then $xxyz$ has 0s alternating with 1s more than once, so $w \notin L$
- ⑤ Contradiction – hence, L is not regular

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p$

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p$
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p$
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$
- 4 Problem: If $y = 0^m 1^m$, then w can be pumped – no contradiction

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p$
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$
- 4 Problem: If $y = 0^m 1^m$, then w can be pumped – no contradiction
- 5 Solution: Use condition that $|xy| \leq p$

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$
- ④ Problem: If $y = 0^m 1^m$, then w can be pumped – no contradiction
- ⑤ Solution: Use condition that $|xy| \leq p$
 - Since $w = 0^p 1^p$ and $|xy| \leq p$, we know that y must be in first p symbols

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$
- ④ Problem: If $y = 0^m 1^m$, then w can be pumped – no contradiction
- ⑤ Solution: Use condition that $|xy| \leq p$
 - Since $w = 0^p 1^p$ and $|xy| \leq p$, we know that y must be in first p symbols
 - But, this means that y must be all 0s

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$
- ④ Problem: If $y = 0^m 1^m$, then w can be pumped – no contradiction
- ⑤ Solution: Use condition that $|xy| \leq p$
 - Since $w = 0^p 1^p$ and $|xy| \leq p$, we know that y must be in first p symbols
 - But, this means that y must be all 0s
- ⑥ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xyyz$ has more 0s than 1s, so $w \notin L$

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$
- ④ Problem: If $y = 0^m 1^m$, then w can be pumped – no contradiction
- ⑤ Solution: Use condition that $|xy| \leq p$
 - Since $w = 0^p 1^p$ and $|xy| \leq p$, we know that y must be in first p symbols
 - But, this means that y must be all 0s
- ⑥ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xyyz$ has more 0s than 1s, so $w \notin L$
- ⑦ Contradiction – hence, L is not regular

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

A simpler proof:

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

A simpler proof:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is nonregular

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

A simpler proof:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is nonregular
- 2 Observe that $L_1 = L \cap 0^* 1^*$

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

A simpler proof:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is nonregular
- 2 Observe that $L_1 = L \cap 0^* 1^*$
- 3 Easy to see that $0^* 1^*$ is regular

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

A simpler proof:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is nonregular
- 2 Observe that $L_1 = L \cap 0^* 1^*$
- 3 Easy to see that $0^* 1^*$ is regular
- 4 Since regular languages are closed under \cap , if L is regular then L_1 must be regular

Example 2

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

A simpler proof:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is nonregular
- 2 Observe that $L_1 = L \cap 0^* 1^*$
- 3 Easy to see that $0^* 1^*$ is regular
- 4 Since regular languages are closed under \cap , if L is regular then L_1 must be regular
- 5 Since we know L_1 is nonregular, this means that L must be nonregular

Exercise

Prove that the following language is nonregular:

$$L = \{0^i 1^j 2^i 3^j \mid i, j > 0\}$$

What's Next?

- We will get plenty of practice with proving languages nonregular
- We will add (a small amount of) memory to our machines to recognize a richer class of languages