

Foundations of Computing

Lecture 7

Arkady Yerukhimovich

February 4, 2025

Outline

- 1 Announcements
- 2 Lecture 6 Review
- 3 Proving Languages Not Regular
- 4 Proving L Not Regular Using the Pumping Lemma
- 5 Proving L Not Regular Using Closure Properties

First Midterm

- First midterm exam will be in class on Thursday, February 20
- Tuesday, February 18 will be a review lecture
- 5 points bonus for anyone who attends ACM Hackathon

Important

If you need to miss lecture on February 20, let me know ASAP.

Outline

- 1 Announcements
- 2 Lecture 6 Review
- 3 Proving Languages Not Regular
- 4 Proving L Not Regular Using the Pumping Lemma
- 5 Proving L Not Regular Using Closure Properties

Lecture 6 Review

- Nonregular languages
- Proving The NFA pumping lemma
- Using the pumping lemma

Lecture 6 Review

- Nonregular languages
- Proving The NFA pumping lemma
- Using the pumping lemma

Today

- Some more examples proving languages are not regular
- Going beyond regular languages

HW2 Problem 4

Let L be a regular language, prove that the following languages are regular.

- ① $NOPREFIX(L) = \{w \in L \mid \text{no proper prefix of } w \text{ is a member of } L\}$
- ② $NOEXTEND(L) = \{w \in L \mid w \text{ is not a proper prefix of any string in } L\}$

HW2 Problem 4

Let L be a regular language, prove that the following languages are regular.

- ① $NOPREFIX(L) = \{w \in L \mid \text{no proper prefix of } w \text{ is a member of } L\}$
- ② $NOEXTEND(L) = \{w \in L \mid w \text{ is not a proper prefix of any string in } L\}$

Example:


- $L = \{00, 11, 001, 101\}$
- $NOPREFIX(L) = \{00, 11, 101\}$
- $NOEXTEND(L) = \{11, 001, 101\}$

$001 \notin NP(L)$
 $001 \in L$ $00 \in L$

HW2 Problem 4

Let L be a regular language, prove that the following languages are regular.

- ① $NOPREFIX(L) = \{w \in L \mid \text{no proper prefix of } w \text{ is a member of } L\}$




$\bar{w} \in L$ and a proper prefix of $\bar{w} \in L$

HW2 Problem 4

Let L be a regular language, prove that the following languages are regular.

- 1 $NOPREFIX(L) = \{w \in L \mid \text{no proper prefix of } w \text{ is a member of } L\}$
- 2 $NOEXTEND(L) = \{w \in L \mid w \text{ is not a proper prefix of any string in } L\}$



Outline

- 1 Announcements
- 2 Lecture 6 Review
- 3 Proving Languages Not Regular**
- 4 Proving L Not Regular Using the Pumping Lemma
- 5 Proving L Not Regular Using Closure Properties

Outline

- 1 Announcements
- 2 Lecture 6 Review
- 3 Proving Languages Not Regular
- 4 Proving L Not Regular Using the Pumping Lemma**
- 5 Proving L Not Regular Using Closure Properties

The Regular Language Pumping Lemma

Pumping Lemma

If L is a regular language, then there exists an integer p (the pumping length) where any string $w \in L$ such that $|w| \geq p$ can be divided into three pieces $w = xyz$ satisfying:

- 1 For each $i \geq 0$, $xy^iz \in L$
- 2 $|y| > 0$, and
- 3 $|xy| \leq p$

The Proof Procedure

To use the pumping lemma to prove that L is not regular, we do the following:

The Proof Procedure

To use the pumping lemma to prove that L is not regular, we do the following:

- 1 Assume that L is regular

The Proof Procedure

To use the pumping lemma to prove that L is not regular, we do the following:

- 1 Assume that L is regular
- 2 Use pumping lemma to guarantee pumping length p , s.t. all w with $|w| > p$ can be pumped – Note: proof must work for all p

The Proof Procedure

To use the pumping lemma to prove that L is not regular, we do the following:

- 1 Assume that L is regular
- 2 Use pumping lemma to guarantee pumping length p , s.t. all w with $|w| > p$ can be pumped – Note: proof must work for all p
- 3 Choose $w \in L$ with $|w| \geq p$

The Proof Procedure

To use the pumping lemma to prove that L is not regular, we do the following:

- 1 Assume that L is regular
- 2 Use pumping lemma to guarantee pumping length p , s.t. all w with $|w| > p$ can be pumped – Note: proof must work for all p
- 3 Choose $w \in L$ with $|w| \geq p$
- 4 Demonstrate that w cannot be pumped
 - For each possible division $w = xyz$ (with $|y| > 0$ and $|xy| \leq p$), find an integer i such that $xy^iz \notin L$

The Proof Procedure

To use the pumping lemma to prove that L is not regular, we do the following:

- ① Assume that L is regular
- ② Use pumping lemma to guarantee pumping length p , s.t. all w with $|w| > p$ can be pumped – Note: proof must work for all p
- ③ Choose $w \in L$ with $|w| \geq p$
- ④ Demonstrate that w cannot be pumped
 - For each possible division $w = xyz$ (with $|y| > 0$ and $|xy| \leq p$), find an integer i such that $xy^iz \notin L$
- ⑤ Contradiction!!!

Prior Examples

We've already seen how to prove:

- $L = \{0^n 1^n \mid n \geq 0\}$ is not regular

Let's try something a little harder

Another Example

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Another Example

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies

Another Example

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p$

Another Example

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p$
- 3 By pumping lemma, $w = xyz$ s.t. $xy^iz \in L$

$$y = 0^x$$

$$z = 1^p$$

$$y = 01$$

$$00010111$$

Another Example

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p$
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$
- 4 Problem: If $y = 0^m 1^m$, then w can be pumped – no contradiction

Another Example

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- 1 Assume L is regular, and let p be the pumping length this implies
- 2 Choose $w = 0^p 1^p$
- 3 By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$
- 4 Problem: If $y = 0^m 1^m$, then w can be pumped – no contradiction
- 5 Solution: Use condition that $|xy| \leq p$

Another Example

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$
- ④ Problem: If $y = 0^m 1^m$, then w can be pumped – no contradiction
- ⑤ Solution: Use condition that $|xy| \leq p$
 - Since $w = 0^p 1^p$ and $|xy| \leq p$, we know that y must be in first p symbols

Another Example

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$
- ④ Problem: If $y = 0^m 1^m$, then w can be pumped – no contradiction
- ⑤ Solution: Use condition that $|xy| \leq p$
 - Since $w = 0^p 1^p$ and $|xy| \leq p$, we know that y must be in first p symbols
 - But, this means that y must be all 0s

Another Example

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$
- ④ Problem: If $y = 0^m 1^m$, then w can be pumped – no contradiction
- ⑤ Solution: Use condition that $|xy| \leq p$
 - Since $w = 0^p 1^p$ and $|xy| \leq p$, we know that y must be in first p symbols
 - But, this means that y must be all 0s
- ⑥ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xyyz$ has more 0s than 1s, so $w \notin L$

Another Example

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

Proof:

- ① Assume L is regular, and let p be the pumping length this implies
- ② Choose $w = 0^p 1^p$
- ③ By pumping lemma, $w = xyz$ s.t. $xy^i z \in L$
- ④ Problem: If $y = 0^m 1^m$, then w can be pumped – no contradiction
- ⑤ Solution: Use condition that $|xy| \leq p$
 - Since $w = 0^p 1^p$ and $|xy| \leq p$, we know that y must be in first p symbols
 - But, this means that y must be all 0s
- ⑥ Complete proof by considering all possible values for y
 - y consists of only 0s – then $xyyz$ has more 0s than 1s, so $w \notin L$
- ⑦ Contradiction – hence, L is not regular

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- 1 Assume L is regular, and let p be the pumping length

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- 1 Assume L is regular, and let p be the pumping length
- 2 For any sufficiently long w , by pumping lemma, there must be a partition $w = xyz$ s.t. $xy^i z \in L$ for all i

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- 1 Assume L is regular, and let p be the pumping length
- 2 For any sufficiently long w , by pumping lemma, there must be a partition $w = xyz$ s.t. $xy^i z \in L$ for all i
- 3 Goal: Show that for all partitions, $xy^i z \notin L$ for some $i \geq 0$

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- 1 Assume L is regular, and let p be the pumping length
- 2 For any sufficiently long w , by pumping lemma, there must be a partition $w = xyz$ s.t. $xy^i z \in L$ for all i
- 3 Goal: Show that for all partitions, $xy^i z \notin L$ for some $i \geq 0$
That is, $xy^i z = 0^{m'} 1^{n'}$ with $m' = n'$.

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

- 1 Assume L is regular, and let p be the pumping length
- 2 For any sufficiently long w , by pumping lemma, there must be a partition $w = xyz$ s.t. $xy^i z \in L$ for all i
- 3 Goal: Show that for all partitions, $xy^i z \notin L$ for some $i \geq 0$
That is, $xy^i z = 0^{m'} 1^{n'}$ with $m' = n'$.

Question

What w should we choose?

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{m'}$.

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{m'}$.

- 1 Suppose we choose $w = 0^p 1^{p+1}$, then since $|xy| \leq p$,
 $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{p-(\alpha+\beta)} 1^{p+1}$

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{m'}$.

- 1 Suppose we choose $w = 0^p 1^{p+1}$, then since $|xy| \leq p$,
 $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{p-(\alpha+\beta)} 1^{p+1}$
- 2 Consider what happens when we pump k times:

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{m'}$.

- 1 Suppose we choose $w = 0^p 1^{p+1}$, then since $|xy| \leq p$,
 $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{p-(\alpha+\beta)} 1^{p+1}$

$\beta \geq 1$

- 2 Consider what happens when we pump k times:

$$xy^k z = 0^{\alpha+k\beta+p-(\alpha+\beta)} 1^{p+1}.$$

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{n'}$.

- 1 Suppose we choose $w = 0^p 1^{p+1}$, then since $|xy| \leq p$,
 $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{p-(\alpha+\beta)} 1^{p+1}$
- 2 Consider what happens when we pump k times:

$$xy^k z = 0^{\alpha+k\beta+p-(\alpha+\beta)} 1^{p+1}.$$

For this to give a contradiction we need

$$m' = n', \text{ i.e. } \cancel{\alpha} + k\beta + p - \cancel{(\alpha + \beta)} = \underline{p + (k-1)\beta} = p + 1$$

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{n'}$.

- 1 Suppose we choose $w = 0^p 1^{p+1}$, then since $|xy| \leq p$,
 $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{p-(\alpha+\beta)} 1^{p+1}$
- 2 Consider what happens when we pump k times:

$$xy^k z = 0^{\alpha+k\beta+p-(\alpha+\beta)} 1^{p+1}.$$

For this to give a contradiction we need

$$m' = n', \text{ i.e. } \alpha + k\beta + p - (\alpha + \beta) = p + (k - 1)\beta = p + 1$$

Equivalently, we need

$$(k - 1)\beta = 1$$

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{m'}$.

- 1 Suppose we choose $w = 0^p 1^{p+1}$, then since $|xy| \leq p$,
 $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{p-(\alpha+\beta)} 1^{p+1}$
- 2 Consider what happens when we pump k times:

$$xy^k z = 0^{\alpha+k\beta+p-(\alpha+\beta)} 1^{p+1}.$$

For this to give a contradiction we need

$$m' = n', \text{ i.e. } \alpha + k\beta + p - (\alpha + \beta) = p + (k - 1)\beta = p + 1$$

Equivalently, we need

$$(k - 1)\beta = 1$$

- 3 But, we can't control β , so this w does not work

Let's try again!!!

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{m'}$.

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{m'}$.

- 1 Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$,
 $z = 0^{m-(\alpha+\beta)} 1^n$

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{m'}$.

- 1 Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{m-(\alpha+\beta)} 1^n$
- 2 Consider what happens when we pump k times:

$$xy^k z = 0^{\overbrace{\alpha + k\beta + m - (\alpha + \beta)}^{\text{from } x \text{ and } z}} 1^n.$$

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{n'}$.

- 1 Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$,
 $z = 0^{m-(\alpha+\beta)} 1^n$
- 2 Consider what happens when we pump k times:

$$xy^k z = 0^{\cancel{\alpha} + k\beta + m - (\cancel{\alpha} + \beta)} 1^n.$$

We need a k s.t. $m + (k - 1)\beta = n$ for a contradiction

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{m'}$.

- 1 Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{m-(\alpha+\beta)} 1^n$
- 2 Consider what happens when we pump k times:

$$xy^k z = 0^{\alpha+k\beta+m-(\alpha+\beta)} 1^n.$$

We need a k s.t. $m + (k-1)\beta = n$ for a contradiction
Equivalently, we need $k = 1 + (n-m)/\beta$ to be an integer

$1 + \frac{n-m}{\beta}$ $1 \leq \beta \leq p$

$p!$

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{m'}$.

- 1 Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$,
 $z = 0^{m-(\alpha+\beta)} 1^n$
- 2 Consider what happens when we pump k times:

$$xy^k z = 0^{\alpha+k\beta+m-(\alpha+\beta)} 1^n.$$

We need a k s.t. $m + (k - 1)\beta = n$ for a contradiction

Equivalently, we need $k = 1 + (n - m)/\beta$ to be an integer

- 3 We only know $\beta \leq p$, how can we guarantee $(n - m)$ is divisible by β ?

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{m'}$.

- 1 Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{m-(\alpha+\beta)} 1^n$
- 2 Consider what happens when we pump k times:

$$xy^k z = 0^{\alpha+k\beta+m-(\alpha+\beta)} 1^n.$$

We need a k s.t. $m + (k - 1)\beta = n$ for a contradiction

Equivalently, we need $k = 1 + (n - m)/\beta$ to be an integer

- 3 We only know $\beta \leq p$, how can we guarantee $(n - m)$ is divisible by β ?

Hint: What number is divisible by all integers $\leq p$?

A More Challenging Example

Consider $L = \{0^m 1^n \mid m \neq n\}$, prove L is not regular

Goal

Pick a w s.t. for all partitions $w = xyz$, for some $i \geq 0$, $xy^i z = 0^{m'} 1^{m'}$.

- 1 Suppose we choose $w = 0^m 1^n$ with $m \geq p$, then $x = 0^\alpha$, $y = 0^\beta$, $z = 0^{m-(\alpha+\beta)} 1^n$
- 2 Consider what happens when we pump k times:

$$xy^k z = 0^{\alpha+k\beta+m-(\alpha+\beta)} 1^n.$$

We need a k s.t. $m + (k - 1)\beta = n$ for a contradiction

Equivalently, we need $k = 1 + (n - m)/\beta$ to be an integer

- 3 We only know $\beta \leq p$, how can we guarantee $(n - m)$ is divisible by β ?

Hint: What number is divisible by all integers $\leq p$?

- 4 Set $n = 2(p!)$, $m = p!$, then $(n - m) = p!$ is divisible by β , so there is k s.t. $xy^k z \notin L$

Hints for Using the Pumping Lemma

To use the pumping lemma, need to do the following

- Identify what it means for $x \notin L$
- Choose w such that any valid split xyz can lead to a contradiction
- Prove that $w' = xy^kz \notin L$ for some k

Choosing w is often tricky, requires intuition and some trial and error.

Outline

- 1 Announcements
- 2 Lecture 6 Review
- 3 Proving Languages Not Regular
- 4 Proving L Not Regular Using the Pumping Lemma
- 5 Proving L Not Regular Using Closure Properties

Proving Non-Regularity Using Closure Properties

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

A simpler proof:

Proving Non-Regularity Using Closure Properties

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

A simpler proof:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is nonregular

Proving Non-Regularity Using Closure Properties

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

A simpler proof:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is nonregular
- 2 Observe that $L_1 = L \cap 0^* 1^*$

Proving Non-Regularity Using Closure Properties

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

A simpler proof:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is nonregular
- 2 Observe that $L_1 = L \cap 0^* 1^*$
- 3 Easy to see that $0^* 1^*$ is regular
- 4 Since regular languages are closed under \cap , if L is regular then L_1 must be regular



Proving Non-Regularity Using Closure Properties

Consider $L = \{w \mid w \text{ has an equal number of 0s and 1s}\}$, prove L is not regular

A simpler proof:

- 1 We already proved that $L_1 = \{0^n 1^n \mid n \geq 0\}$ is nonregular
- 2 Observe that $L_1 = L \cap 0^* 1^*$
- 3 Easy to see that $0^* 1^*$ is regular
- 4 Since regular languages are closed under \cap , if L is regular then L_1 must be regular
- 5 Since we know L_1 is nonregular, this means that L must be nonregular

1. Assume L is regular
2. $\Rightarrow L_1$ is regular
contradiction

Using Closure Properties of Regular Languages

We have seen a number of closure properties of REs

- ① Closure under complement: \overline{L} is regular if L is
- ② Closure under union: $L_1 \cup L_2$ is regular if L_1, L_2 are
- ③ Closure under intersection: $L_1 \cap L_2$ is regular if L_1, L_2 are
- ④ Closure under reversal: L^R is regular if L is
- ⑤ NOPREFIX, NOEXTEND
- ⑥ There are many more (e.g., set difference, cross product, ...)

Using Closure Properties of Regular Languages

We have seen a number of closure properties of REs

- 1 Closure under complement: \overline{L} is regular if L is
- 2 Closure under union: $L_1 \cup L_2$ is regular if L_1, L_2 are
- 3 Closure under intersection: $L_1 \cap L_2$ is regular if L_1, L_2 are
- 4 Closure under reversal: L^R is regular if L is
- 5 NOPREFIX, NOEXTEND
- 6 There are many more (e.g., set difference, cross product, ...)

Important

- It is often much easier to prove non-regularity using closure properties
- Try this first before you turn to pumping lemma

Exercise

Prove that the following language is nonregular:

$$L = \{0^i 1^j 2^i 3^j \mid i, j > 0\}$$