

# Foundations of Computing

## Lecture 16

Arkady Yerukhimovich

March 18, 2025

# Outline

- 1 Lecture 15 Review
- 2 Proofs by Reduction
- 3 Example Proofs by Reduction

# Lecture 15 Review

- Countable and Uncountable Sets
  - Diagonalization
- Proving  $A_{TM}$  is Undecidable

# Announcements

## Homework 6 is out

- Due at 5PM on Monday, March 24
- Early deadline by midnight on Friday, March 21.

# Announcements

## Homework 6 is out

- Due at 5PM on Monday, March 24
- Early deadline by midnight on Friday, March 21.

## Exam 2

- Exam 2 will be in class next Thursday, March 27
- Next Tuesday lecture and Wednesday lab will be for review
- You will again be allowed 2 pieces of paper with notes

# Announcements

## Homework 6 is out

- Due at 5PM on Monday, March 24
- Early deadline by midnight on Friday, March 21.

## Exam 2

- Exam 2 will be in class next Thursday, March 27
- Next Tuesday lecture and Wednesday lab will be for review
- You will again be allowed 2 pieces of paper with notes
- Exam will cover the following topics:
  - Turing Machines
  - Countable and uncountable sets
  - Decidable, Turing-recognizable Languages
  - Proofs by reduction
  - Everything we cover this week
- CFL Pumping Lemma will not be on the exam

# Outline

- 1 Lecture 15 Review
- 2 Proofs by Reduction
- 3 Example Proofs by Reduction

# Another Way to Prove Undecidability

## Reductions Between Problems

There is a reduction from a problem  $A$  to a problem  $B$  if we can use a solution to problem  $B$  to solve problem  $A$

$$A \leq B$$



# Another Way to Prove Undecidability

## Reductions Between Problems

There is a reduction from a problem  $A$  to a problem  $B$  if we can use a solution to problem  $B$  to solve problem  $A$

$$A \leq B$$

## Intuition

$A \leq B$  means that:

- problem  $A$  is no harder than problem  $B$ .

# Another Way to Prove Undecidability

## Reductions Between Problems

There is a reduction from a problem  $A$  to a problem  $B$  if we can use a solution to problem  $B$  to solve problem  $A$

$$A \leq B$$

## Intuition

$A \leq B$  means that:

- problem  $A$  is no harder than problem  $B$ .
- Equivalently, problem  $B$  is no easier than problem  $A$

## Main Observation

Suppose that  $A \leq B$ , then:

- If  $A$  is undecidable
- $B$  must also be undecidable

## Main Observation

Suppose that  $A \leq B$ , then:

- If  $A$  is undecidable
- $B$  must also be undecidable

Proof: (by contradiction)

# Reductions and Undecidability

## Main Observation

Suppose that  $A \leq B$ , then:

- If  $A$  is undecidable
- $B$  must also be undecidable

Proof: (by contradiction)

- Suppose that  $B$  is decidable

-  $\exists$   $TM_B$  that decides  $B$   
 $TM_B$  Halts on all inputs  
and outputs if  
 $x \in B$  or not

# Reductions and Undecidability

## Main Observation

Suppose that  $A \leq B$ , then:

- If  $A$  is undecidable
- $B$  must also be undecidable

Proof: (by contradiction)

- Suppose that  $B$  is decidable
- Since  $A \leq B$ , there exists an algorithm (i.e., a reduction) that uses a solution to  $B$  to solve  $A$

*TM<sub>B</sub> decides B*

# Reductions and Undecidability

I know  $A_{TM}$  is undecidable  $A_{TM} \leq L$

## Main Observation

Suppose that  $A \leq B$ , then:

- If  $A$  is undecidable
- $B$  must also be undecidable

Proof: (by contradiction)

- Suppose that  $B$  is decidable
- Since  $A \leq B$ , there exists an algorithm (i.e., a reduction) that uses a solution to  $B$  to solve  $A$
- But, this means that  $A$  is decidable by running the reduction using the decider machine for  $B$ .

$TMA(x)$

1. Run  $R(x)$
2. whenever

$TM_B$ ,  $R$

$R$  needs to answer if  $x' \in B$ , run  $TM_B(x')$

# Outline

- 1 Lecture 15 Review
- 2 Proofs by Reduction
- 3 Example Proofs by Reduction



# Undecidability of $HALT_{TM}$

$$HALT_{TM} = \{ \langle M, w \rangle \mid M \text{ is a TM and } M \text{ halts on input } w \}$$

# Undecidability of $HALT_{TM}$

$$HALT_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ halts on input } w\}$$

Theorem:  $HALT$  is undecidable

# Undecidability of $HALT_{TM}$

$$HALT_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ halts on input } w\}$$

Theorem:  $HALT$  is undecidable

Proof Sketch:

- Recall that  $A_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M(w) = 1\}$  is undecidable.

# Undecidability of $HALT_{TM}$

$$HALT_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ halts on input } w\}$$

Theorem:  $HALT$  is undecidable

Proof Sketch:

- Recall that  $A_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M(w) = 1\}$  is undecidable.
- We show that  $A_{TM} \leq HALT_{TM}$

# Undecidability of $HALT_{TM}$

$$HALT_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ halts on input } w\}$$

Theorem:  $HALT$  is undecidable

Proof Sketch:

- Recall that  $A_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M(w) = 1\}$  is undecidable.
- We show that  $A_{TM} \leq HALT_{TM}$
- This shows that  $HALT_{TM}$  is also undecidable

$$HALT_{TM} \leq A_{TM}$$

---

# Undecidability of $HALT_{TM}$

$$HALT_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ halts on input } w\}$$

Theorem:  $HALT$  is undecidable

Proof Sketch:

- Recall that  $A_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M(w) = 1\}$  is undecidable.
- We show that  $A_{TM} \leq HALT_{TM}$
- This shows that  $HALT_{TM}$  is also undecidable

Proof:

Construct reduction  $R$  that decides  $A_{TM}$  given a TM  $D$  that decides  $HALT$

# Undecidability of $HALT_{TM}$

$$HALT_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ halts on input } w\}$$

Theorem:  $HALT$  is undecidable

Proof Sketch:

- Recall that  $A_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M(w) = 1\}$  is undecidable.
- We show that  $A_{TM} \leq HALT_{TM}$
- This shows that  $HALT_{TM}$  is also undecidable

Proof:

Construct reduction  $R$  that decides  $A_{TM}$  given a TM  $D$  that decides  $HALT$   
On input  $\langle M, w \rangle$ ,  $R$  does the following:

# Undecidability of $HALT_{TM}$

$$HALT_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ halts on input } w\}$$

Theorem:  $HALT$  is undecidable

Proof Sketch:

- Recall that  $A_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M(w) = 1\}$  is undecidable.
- We show that  $A_{TM} \leq HALT_{TM}$
- This shows that  $HALT_{TM}$  is also undecidable

Proof:

Construct reduction  $R$  that decides  $A_{TM}$  given a TM  $D$  that decides  $HALT$   
On input  $\langle M, w \rangle$ ,  $R$  does the following:

- Run  $D(\langle M, w \rangle)$



# Undecidability of $HALT_{TM}$

$$HALT_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ halts on input } w\}$$

Theorem:  $HALT$  is undecidable

Proof Sketch:

- Recall that  $A_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M(w) = 1\}$  is undecidable.
- We show that  $A_{TM} \leq HALT_{TM}$
- This shows that  $HALT_{TM}$  is also undecidable

Proof:

Construct reduction  $R$  that decides  $A_{TM}$  given a TM  $D$  that decides  $HALT$   
On input  $\langle M, w \rangle$ ,  $R$  does the following:


- Run  $D(\langle M, w \rangle)$
- If  $D$  rejects –  $M(w)$  doesn't halt – halt and reject

# Undecidability of $HALT_{TM}$

$$HALT_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M \text{ halts on input } w\}$$

Theorem:  $HALT$  is undecidable

Proof Sketch:

- Recall that  $A_{TM} = \{\langle M, w \rangle \mid M \text{ is a TM and } M(w) = 1\}$  is undecidable.
- We show that  $A_{TM} \leq HALT_{TM}$  
- This shows that  $HALT_{TM}$  is also undecidable

Proof:

Construct reduction  $R$  that decides  $A_{TM}$  given a TM  $D$  that decides  $HALT$   
On input  $\langle M, w \rangle$ ,  $R$  does the following:

- Run  $D(\langle M, w \rangle)$
- If  $D$  rejects –  $M(w)$  doesn't halt – halt and reject
- if  $D$  accepts –  $M(w)$  halts – Simulate  $M(w)$  until it halts, and output whatever  $M$  outputs

# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

Proof Sketch:

- We show that  $A_{TM} \leq REGULAR_{TM}$

# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

Proof Sketch:

- We show that  $A_{TM} \leq REGULAR_{TM}$  ( $M, w$ )
- Specifically, reduction builds another TM  $M'$  s.t.

# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

Proof Sketch:

$(M, w)$

- We show that  $A_{TM} \leq REGULAR_{TM}$
- Specifically, reduction builds another TM  $M'$  s.t.
  - If  $M$  accepts  $w$ ,  $M'$  recognizes  $\Sigma^*$  – regular language

# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

Proof Sketch:

- We show that  $A_{TM} \leq REGULAR_{TM}$
- Specifically, reduction builds another TM  $M'$  s.t.
  - If  $M$  accepts  $w$ ,  $M'$  recognizes  $\Sigma^*$  – regular language
  - If  $M$  does not accept  $w$ ,  $M'$  recognizes  $\{0^n 1^n\}$  – not regular



# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

Proof Sketch:

- We show that  $A_{TM} \leq REGULAR_{TM}$
- Specifically, reduction builds another TM  $M'$  s.t.
  - If  $M$  accepts  $w$ ,  $M'$  recognizes  $\Sigma^*$  – regular language
  - If  $M$  does not accept  $w$ ,  $M'$  recognizes  $\{0^n 1^n\}$  – not regular
- If we can decide whether  $M'$  recognizes a regular language or not, can use that to decide whether  $M$  accepts  $w$  or not

# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

Proof:

# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

$$A_{TM} \leq REGULAR_{TM}$$

Proof:

Reduction  $R$  that decides  $A_{TM}$  given a TM  $D$  that decides  $REGULAR_{TM}$

# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

Proof:

Reduction  $R$  that decides  $A_{TM}$  given a TM  $D$  that decides  $REGULAR_{TM}$

On input  $\langle M, w \rangle$ :

# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

Proof:

Reduction  $R$  that decides  $A_{TM}$  given a TM  $D$  that decides  $REGULAR_{TM}$

On input  $\langle M, w \rangle$ :

- 1 Construct TM  $M'_{\langle M, w \rangle}$  s.t.  $M'_{\langle M, w \rangle}(x)$  is as follows:

# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

Proof:

Reduction  $R$  that decides  $A_{TM}$  given a TM  $D$  that decides  $REGULAR_{TM}$

On input  $\langle M, w \rangle$ :

- 1 Construct TM  $M'_{\langle M, w \rangle}$  s.t.  $M'_{\langle M, w \rangle}(x)$  is as follows:
  - If  $x = 0^n 1^n$ , accept

# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

Proof:

Reduction  $R$  that decides  $A_{TM}$  given a TM  $D$  that decides  $REGULAR_{TM}$

On input  $\langle M, w \rangle$ :

- 1 Construct TM  $M'_{\langle M, w \rangle}$  s.t.  $M'_{\langle M, w \rangle}(x)$  is as follows:
  - If  $x = 0^n 1^n$ , accept
  - If  $x$  does not have this form, run  $M(w)$  and accept if it accepts

# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

Proof:

Reduction  $R$  that decides  $A_{TM}$  given a TM  $D$  that decides  $REGULAR_{TM}$

On input  $\langle M, w \rangle$ :

- ① Construct TM  $M'_{\langle M, w \rangle}$  s.t.  $M'_{\langle M, w \rangle}(x)$  is as follows:
  - If  $x = 0^n 1^n$ , accept
  - If  $x$  does not have this form, run  $M(w)$  and accept if it accepts
- ② Run  $D$  on input  $\langle M' \rangle$

$M(w) = 1$  , what  $x$  does  $M'$  accept?  $\hookrightarrow$   
 $M(w) \neq 1$  ,  $0^n 1^n$



# Other Undecidable Languages

$$REGULAR_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } L(M) \text{ is a regular language}\}$$

Theorem:  $REGULAR_{TM}$  is undecidable

Proof:

Reduction  $R$  that decides  $A_{TM}$  given a TM  $D$  that decides  $REGULAR_{TM}$

On input  $\langle M, w \rangle$ :

- 1 Construct TM  $M'_{\langle M, w \rangle}$  s.t.  $M'_{\langle M, w \rangle}(x)$  is as follows:
  - If  $x = 0^n 1^n$ , accept
  - If  $x$  does not have this form, run  $M(w)$  and accept if it accepts
- 2 Run  $D$  on input  $\langle M' \rangle$ 
  - If  $M(w) = 1$ , then  $M'_{\langle M, w \rangle}$  accepts all  $x \in \Sigma^*$  – regular
  - If  $M(w) \neq 1$ ,  $M'_{\langle M, w \rangle}$  accepts the language  $0^n 1^n$  – not regular
- 3 Output what  $D$  outputs

# Other Undecidable Languages – Exercise

$$EMPTY - STRING_{TM} = \{\langle M \rangle \mid M \text{ is a TM and } M(\epsilon) = 1\}$$

Think about:

- What direction should the reduction go?
- What language should the reduction use?