

Cryptography

Lecture 19

Arkady Yerukhimovich

November 4, 2024

- 1 Lecture 18 Review
- 2 Crypto Hardness Assumptions (Chapters 8.2, 8.3)
- 3 Assumptions in Cyclic Groups (Chapters 8.2, 8.3)

Lecture 18 Review

- The Group \mathbb{Z}_N^*
- Chinese Remainder Theorem
- Modular Arithmetic by Hand

Modular Arithmetic Without a Calculator

To evaluate exponentiation $\text{mod } N$ use the following steps:

- If N is not prime, apply the Chinese Remainder Theorem
- Reduce $\text{mod } \phi(N)$ in the exponent
- Reduce $\text{mod } N$ in the base

Useful Hints:

- Sometimes useful to use negative numbers
- look for things that are easy to compute (e.g., 1^{53})

- 1 Lecture 18 Review
- 2 **Crypto Hardness Assumptions (Chapters 8.2, 8.3)**
- 3 Assumptions in Cyclic Groups (Chapters 8.2, 8.3)

What Are Hardness Assumptions?

- As we've discussed before, all crypto primitives rely on computational hardness
- Thus, we need to assume that some problem is hard to compute
- We have seen such assumptions before: E.g., Existence of PRG, PRF, CRHF

What Are Hardness Assumptions?

- As we've discussed before, all crypto primitives rely on computational hardness
- Thus, we need to assume that some problem is hard to compute
- We have seen such assumptions before: E.g., Existence of PRG, PRF, CRHF
- Going forward, we will instead use hard problems from number theory and mathematics
 - Some of these problems have been studied for 1000s of years
 - Easy to state and widely understood
 - Still believed to be hard for all PPT machines

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

GenMod(1^n):

- Find n -bit primes p, q , compute $N = pq$
- Output (N, p, q)

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

GenMod(1^n):

- Find n -bit primes p, q , compute $N = pq$
- Output (N, p, q)

Consider the following game between an adversary \mathcal{A} and a challenger:

Factor $_{\mathcal{A}, \text{GenMod}}(n)$

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

GenMod(1^n):

- Find n -bit primes p, q , compute $N = pq$
- Output (N, p, q)

Consider the following game between an adversary \mathcal{A} and a challenger:

Factor $_{\mathcal{A}, \text{GenMod}}(n)$

- The challenger runs $(N, p, q) \leftarrow \text{GenMod}(1^n)$ and sends N to \mathcal{A}

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

GenMod(1^n):

- Find n -bit primes p, q , compute $N = pq$
- Output (N, p, q)

Consider the following game between an adversary \mathcal{A} and a challenger:

Factor $_{\mathcal{A}, \text{GenMod}}(n)$

- The challenger runs $(N, p, q) \leftarrow \text{GenMod}(1^n)$ and sends N to \mathcal{A}
- \mathcal{A} outputs two primes p', q'

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

GenMod(1^n):

- Find n -bit primes p, q , compute $N = pq$
- Output (N, p, q)

Consider the following game between an adversary \mathcal{A} and a challenger:

Factor $_{\mathcal{A}, \text{GenMod}}(n)$

- The challenger runs $(N, p, q) \leftarrow \text{GenMod}(1^n)$ and sends N to \mathcal{A}
- \mathcal{A} outputs two primes p', q'
- We say that $\text{Factor}_{\mathcal{A}, \text{GenMod}}(n) = 1$ (i.e., \mathcal{A} wins) if $p' \cdot q' = N$.

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

GenMod(1^n):

- Find n -bit primes p, q , compute $N = pq$
- Output (N, p, q)

Consider the following game between an adversary \mathcal{A} and a challenger:

Factor $_{\mathcal{A}, \text{GenMod}}(n)$

- The challenger runs $(N, p, q) \leftarrow \text{GenMod}(1^n)$ and sends N to \mathcal{A}
- \mathcal{A} outputs two primes p', q'
- We say that $\text{Factor}_{\mathcal{A}, \text{GenMod}}(n) = 1$ (i.e., \mathcal{A} wins) if $p' \cdot q' = N$.

Definition: Factoring is hard relative to GenMod if for all PPT \mathcal{A} it holds that

$$\Pr[\text{Factor}_{\mathcal{A}, \text{GenMod}}(n) = 1] \leq \text{negl}(n)$$

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

Observations:

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

Observations:

- Factoring can easily be done in $O(\sqrt{N})$ divisions – just try all numbers less than \sqrt{N}

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

Observations:

- Factoring can easily be done in $O(\sqrt{N})$ divisions – just try all numbers less than \sqrt{N}
- No known way to factor in time polynomial in $\|N\| = \log_2 N$

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

Observations:

- Factoring can easily be done in $O(\sqrt{N})$ divisions – just try all numbers less than \sqrt{N}
- No known way to factor in time polynomial in $||N||$
- Requires ability to efficiently sample n -bit primes

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

Observations:

- Factoring can easily be done in $O(\sqrt{N})$ divisions – just try all numbers less than \sqrt{N}
- No known way to factor in time polynomial in $||N||$
- Requires ability to efficiently sample n -bit primes
 - Prime number theorem: The fraction of n -bit integers that are prime is at least $1/3n$

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

Observations:

- Factoring can easily be done in $O(\sqrt{N})$ divisions – just try all numbers less than \sqrt{N}
- No known way to factor in time polynomial in $||N||$
- Requires ability to efficiently sample n -bit primes
 - Prime number theorem: The fraction of n -bit integers that are prime is at least $1/3n$
 - So, can sample integers at random, and test if they are prime

Factoring Assumption

Factoring Problem

Given $N = pq$ when p and q are n -bit primes, find p and q

Observations:

- Factoring can easily be done in $O(\sqrt{N})$ divisions – just try all numbers less than \sqrt{N}
- No known way to factor in time polynomial in $||N||$
- Requires ability to efficiently sample n -bit primes
 - Prime number theorem: The fraction of n -bit integers that are prime is at least $1/3n$
 - So, can sample integers at random, and test if they are prime
 - Miller-Rabin primality test – efficiently test if a number is prime

RSA Assumption (Rivest, Shamir, Adleman '77)

Given $N = pq$, Integer $e > 1$ s.t. $\gcd(e, \phi(N)) = 1$, we know that $f_e(x) = x^e$ is a permutation over \mathbb{Z}_N^*

RSA Problem

Given (N, e) and $y \in \mathbb{Z}_N^*$, compute $[y^{1/e} \bmod N]$

$$f_e(x) = y = x^e \bmod N$$

RSA Assumption (Rivest, Shamir, Adleman '77)

Given $N = pq$, Integer $e > 1$ s.t. $\gcd(e, \phi(N)) = 1$, we know that $f_e(x) = x^e$ is a permutation over \mathbb{Z}_N^*

RSA Problem

Given (N, e) and $y \in \mathbb{Z}_N^*$, compute $[y^{1/e} \bmod N]$

Observations:

RSA Assumption (Rivest, Shamir, Adleman '77)

Given $N = pq$, Integer $e > 1$ s.t. $\gcd(e, \phi(N)) = 1$, we know that $f_e(x) = x^e$ is a permutation over \mathbb{Z}_N^*

RSA Problem

Given (N, e) and $y \in \mathbb{Z}_N^*$, compute $[y^{1/e} \bmod N]$

Observations:

- Since $\gcd(e, \phi(N)) = 1$, there is an integer $d = e^{-1} \bmod \phi(N)$

RSA Assumption (Rivest, Shamir, Adleman '77)

Given $N = pq$, Integer $e > 1$ s.t. $\gcd(e, \phi(N)) = 1$, we know that $f_e(x) = x^e$ is a permutation over \mathbb{Z}_N^*

RSA Problem

Given (N, e) and $y \in \mathbb{Z}_N^*$, compute $[y^{1/e} \bmod N]$

Observations:

- Since $\gcd(e, \phi(N)) = 1$, there is an integer $d = e^{-1} \bmod \phi(N)$
- The function $f_d(x) = x^d$ is also a permutation over \mathbb{Z}_N^*

RSA Assumption (Rivest, Shamir, Adleman '77)

Given $N = pq$, Integer $e > 1$ s.t. $\gcd(e, \phi(N)) = 1$, we know that $f_e(x) = x^e$ is a permutation over \mathbb{Z}_N^*

RSA Problem

Given (N, e) and $y \in \mathbb{Z}_N^*$, compute $[y^{1/e} \bmod N]$

Observations:

- Since $\gcd(e, \phi(N)) = 1$, there is an integer $d = e^{-1} \bmod \phi(N)$
- The function $f_d(x) = x^d$ is also a permutation over \mathbb{Z}_N^*
- Moreover, f_d is the inverse permutation of f_e

$$(x^e)^d = x^{[ed \bmod \phi(N)]} = x \bmod N$$

RSA Assumption (Rivest, Shamir, Adleman '77)

Given $N = pq$, Integer $e > 1$ s.t. $\gcd(e, \phi(N)) = 1$, we know that $f_e(x) = x^e$ is a permutation over \mathbb{Z}_N^*

RSA Problem

Given (N, e) and $y \in \mathbb{Z}_N^*$, compute $[y^{1/e} \bmod N]$

Observations:

- Since $\gcd(e, \phi(N)) = 1$, there is an integer $d = e^{-1} \bmod \phi(N)$
- The function $f_d(x) = x^d$ is also a permutation over \mathbb{Z}_N^*
- Moreover, f_d is the inverse permutation of f_e

$$(x^e)^d = x^{[ed \bmod \phi(N)]} = x \bmod N$$

- RSA problem is easy if know any of $d, \phi(N), p, q$

RSA Assumption (Rivest, Shamir, Adleman '77)

Given $N = pq$, Integer $e > 1$ s.t. $\gcd(e, \phi(N)) = 1$, we know that $f_e(x) = x^e$ is a permutation over \mathbb{Z}_N^*

RSA Problem

Given (N, e) and $y \in \mathbb{Z}_N^*$, compute $[y^{1/e} \bmod N]$

Observations:

- Since $\gcd(e, \phi(N)) = 1$, there is an integer $d = e^{-1} \bmod \phi(N)$
- The function $f_d(x) = x^d$ is also a permutation over \mathbb{Z}_N^*
- Moreover, f_d is the inverse permutation of f_e

$$(x^e)^d = x^{[ed \bmod \phi(N)]} = x \bmod N$$

- RSA problem is easy if know any of $d, \phi(N), p, q$
- RSA is potentially easier than factoring

RSA Assumption

GenRSA(1^n):

- $(N, p, q) \leftarrow \text{GenMod}(1^n)$, let $\phi(N) = (p - 1)(q - 1)$
- Choose $e > 1$ s.t. $\gcd(e, \phi(N)) = 1$
- Compute $d = [e^{-1} \bmod \phi(N)]$
- Output (N, e, d)

RSA Assumption

GenRSA(1^n):

- $(N, p, q) \leftarrow \text{GenMod}(1^n)$, let $\phi(N) = (p - 1)(q - 1)$
- Choose $e > 1$ s.t. $\gcd(e, \phi(N)) = 1$
- Compute $d = [e^{-1} \bmod \phi(N)]$
- Output (N, e, d)

Consider the following game between an adversary \mathcal{A} and a challenger:

RSALnv $_{\mathcal{A}, \text{GenRSA}}(n)$

RSA Assumption

GenRSA(1^n):

- $(N, p, q) \leftarrow \text{GenMod}(1^n)$, let $\phi(N) = (p - 1)(q - 1)$
- Choose $e > 1$ s.t. $\gcd(e, \phi(N)) = 1$
- Compute $d = [e^{-1} \bmod \phi(N)]$
- Output (N, e, d)

Consider the following game between an adversary \mathcal{A} and a challenger:

RSAINV $_{\mathcal{A}, \text{GenRSA}}(n)$

- The challenger runs $(N, e, d) \leftarrow \text{GenRSA}(1^n)$, chooses $y \leftarrow \mathbb{Z}_N^*$, and sends (N, e, y) to \mathcal{A}

RSA Assumption

GenRSA(1^n):

- $(N, p, q) \leftarrow \text{GenMod}(1^n)$, let $\phi(N) = (p - 1)(q - 1)$
- Choose $e > 1$ s.t. $\gcd(e, \phi(N)) = 1$
- Compute $d = [e^{-1} \bmod \phi(N)]$
- Output (N, e, d)

Consider the following game between an adversary \mathcal{A} and a challenger:

RSInv $_{\mathcal{A}, \text{GenRSA}}(n)$

- The challenger runs $(N, e, d) \leftarrow \text{GenRSA}(1^n)$, chooses $y \leftarrow \mathbb{Z}_N^*$, and sends (N, e, y) to \mathcal{A}
- \mathcal{A} outputs $x \in \mathbb{Z}_N^*$

RSA Assumption

GenRSA(1^n):

- $(N, p, q) \leftarrow \text{GenMod}(1^n)$, let $\phi(N) = (p - 1)(q - 1)$
- Choose $e > 1$ s.t. $\gcd(e, \phi(N)) = 1$
- Compute $d = [e^{-1} \bmod \phi(N)]$
- Output (N, e, d)

Consider the following game between an adversary \mathcal{A} and a challenger:

RSAINV $_{\mathcal{A}, \text{GenRSA}}(n)$

- The challenger runs $(N, e, d) \leftarrow \text{GenRSA}(1^n)$, chooses $y \leftarrow \mathbb{Z}_N^*$, and sends (N, e, y) to \mathcal{A}
- \mathcal{A} outputs $x \in \mathbb{Z}_N^*$
- We say that $\text{RSAINV}_{\mathcal{A}, \text{GenRSA}}(n) = 1$ (i.e., \mathcal{A} wins) if $x^e = y \bmod N$.

RSA Assumption

GenRSA(1^n):

- $(N, p, q) \leftarrow \text{GenMod}(1^n)$, let $\phi(N) = (p - 1)(q - 1)$
- Choose $e > 1$ s.t. $\gcd(e, \phi(N)) = 1$
- Compute $d = [e^{-1} \bmod \phi(N)]$
- Output (N, e, d)

Consider the following game between an adversary \mathcal{A} and a challenger:

RSAINV $_{\mathcal{A}, \text{GenRSA}}(n)$

- The challenger runs $(N, e, d) \leftarrow \text{GenRSA}(1^n)$, chooses $y \leftarrow \mathbb{Z}_N^*$, and sends (N, e, y) to \mathcal{A}
- \mathcal{A} outputs $x \in \mathbb{Z}_N^*$
- We say that $\text{RSAINV}_{\mathcal{A}, \text{GenRSA}}(n) = 1$ (i.e., \mathcal{A} wins) if $x^e = y \bmod N$.

Definition: RSA is hard relative to GenRSA if for all PPT \mathcal{A} it holds that

$$\Pr[\text{RSAINV}_{\mathcal{A}, \text{GenRSA}}(n) = 1] \leq \text{negl}(n)$$

- 1 Lecture 18 Review
- 2 Crypto Hardness Assumptions (Chapters 8.2, 8.3)
- 3 Assumptions in Cyclic Groups (Chapters 8.2, 8.3)

Discrete Log Assumption

Let G be a cyclic group of order q with generator g

Discrete Log Problem

Given $h \in G$, find $0 \leq x \leq q - 1$ s.t. $g^x = h$. We say $x = \log_g h$

$$G = \{g^0, g^1, g^2, g^3, \dots, g^{|G|-1}\} \quad |G| = q$$

Discrete Log Assumption

Let G be a cyclic group of order q with generator g

Discrete Log Problem

Given $h \in G$, find $0 \leq x \leq q - 1$ s.t. $g^x = h$. We say $x = \log_g h$

Gen(1^n):

- Pick n -bit prime q , group G of order q , generator g . Output (G, q, g)

Discrete Log Assumption

Let G be a cyclic group of order q with generator g

Discrete Log Problem

Given $h \in G$, find $0 \leq x \leq q - 1$ s.t. $g^x = h$. We say $x = \log_g h$

$\text{Gen}(1^n)$:

- Pick n -bit prime q , group G of order q , generator g . Output (G, q, g)

Consider the following game between an adversary \mathcal{A} and a challenger:

$\text{DLog}_{\mathcal{A}, \text{Gen}}(n)$

Discrete Log Assumption

Let G be a cyclic group of order q with generator g

Discrete Log Problem

Given $h \in G$, find $0 \leq x \leq q - 1$ s.t. $g^x = h$. We say $x = \log_g h$

$\text{Gen}(1^n)$:

- Pick n -bit prime q , group G of order q , generator g . Output (G, q, g)

Consider the following game between an adversary \mathcal{A} and a challenger:

$\text{DLog}_{\mathcal{A}, \text{Gen}}(n)$

- Challenger runs $(G, q, g) \leftarrow \text{Gen}(1^n)$, $h \leftarrow G$, sends (G, q, g, h) to \mathcal{A}

Discrete Log Assumption

Let G be a cyclic group of order q with generator g

Discrete Log Problem

Given $h \in G$, find $0 \leq x \leq q - 1$ s.t. $g^x = h$. We say $x = \log_g h$

$\text{Gen}(1^n)$:

- Pick n -bit prime q , group G of order q , generator g . Output (G, q, g)

Consider the following game between an adversary \mathcal{A} and a challenger:

$\text{DLog}_{\mathcal{A}, \text{Gen}}(n)$

- Challenger runs $(G, q, g) \leftarrow \text{Gen}(1^n)$, $h \leftarrow G$, sends (G, q, g, h) to \mathcal{A}
- \mathcal{A} outputs $x \in \mathbb{Z}_q$

Discrete Log Assumption

Let G be a cyclic group of order q with generator g

Discrete Log Problem

Given $h \in G$, find $0 \leq x \leq q - 1$ s.t. $g^x = h$. We say $x = \log_g h$

$\text{Gen}(1^n)$:

- Pick n -bit prime q , group G of order q , generator g . Output (G, q, g)
- Consider the following game between an adversary \mathcal{A} and a challenger:

$\text{DLog}_{\mathcal{A}, \text{Gen}}(n)$

- Challenger runs $(G, q, g) \leftarrow \text{Gen}(1^n)$, $h \leftarrow G$, sends (G, q, g, h) to \mathcal{A}
- \mathcal{A} outputs $x \in \mathbb{Z}_q$
- We say that $\text{DLog}_{\mathcal{A}, \text{Gen}}(n) = 1$ (i.e., \mathcal{A} wins) if $g^x = h$.

Discrete Log Assumption

Let G be a cyclic group of order q with generator g

Discrete Log Problem

Given $h \in G$, find $0 \leq x \leq q - 1$ s.t. $g^x = h$. We say $x = \log_g h$

$\text{Gen}(1^n)$:

- Pick n -bit prime q , group G of order q , generator g . Output (G, q, g)

Consider the following game between an adversary \mathcal{A} and a challenger:

$\text{DLog}_{\mathcal{A}, \text{Gen}}(n)$

- Challenger runs $(G, q, g) \leftarrow \text{Gen}(1^n)$, $h \leftarrow G$, sends (G, q, g, h) to \mathcal{A}
- \mathcal{A} outputs $x \in \mathbb{Z}_q$
- We say that $\text{DLog}_{\mathcal{A}, \text{Gen}}(n) = 1$ (i.e., \mathcal{A} wins) if $g^x = h$.

Definition: DLog is hard relative to Gen if for all PPT \mathcal{A} it holds that

$$\Pr[\text{DLog}_{\mathcal{A}, \text{Gen}}(n) = 1] \leq \text{negl}(n)$$

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

$$h_1 = g^x$$

$$h_2 = g^y$$

$$g^x \cdot g^y = g^{x+y}$$

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

Variant 1: Computational Diffie-Hellman

- Hard to find g^{xy}

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

Variant 1: Computational Diffie-Hellman

- Hard to find g^{xy}

Consider the following game between an adversary \mathcal{A} and a challenger:

Computation Diffie-Hellman: $\text{CDH}_{\mathcal{A}, \text{Gen}}(n)$

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

Variant 1: Computational Diffie-Hellman

- Hard to find g^{xy}

Consider the following game between an adversary \mathcal{A} and a challenger:

Computation Diffie-Hellman: $\text{CDH}_{\mathcal{A}, \text{Gen}}(n)$

- Challenger runs $(G, q, g) \leftarrow \text{Gen}(1^n)$, $x, y \leftarrow \mathbb{Z}_q$, sends $(G, q, g, h_1 = g^x, h_2 = g^y)$ to \mathcal{A}

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

Variant 1: Computational Diffie-Hellman

- Hard to find g^{xy}

Consider the following game between an adversary \mathcal{A} and a challenger:

Computation Diffie-Hellman: $\text{CDH}_{\mathcal{A}, \text{Gen}}(n)$

- Challenger runs $(G, q, g) \leftarrow \text{Gen}(1^n)$, $x, y \leftarrow \mathbb{Z}_q$, sends $(G, q, g, h_1 = g^x, h_2 = g^y)$ to \mathcal{A}
- \mathcal{A} outputs $h_3 \in G$

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

Variant 1: Computational Diffie-Hellman

- Hard to find g^{xy}

Consider the following game between an adversary \mathcal{A} and a challenger:

Computation Diffie-Hellman: $\text{CDH}_{\mathcal{A}, \text{Gen}}(n)$

- Challenger runs $(G, q, g) \leftarrow \text{Gen}(1^n)$, $x, y \leftarrow \mathbb{Z}_q$, sends $(G, q, g, h_1 = g^x, h_2 = g^y)$ to \mathcal{A}
- \mathcal{A} outputs $h_3 \in G$
- We say that $\text{CDH}_{\mathcal{A}, \text{Gen}}(n) = 1$ (i.e., \mathcal{A} wins) if $h_3 = g^{xy}$.

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

Variant 1: Computational Diffie-Hellman

- Hard to find g^{xy}

Consider the following game between an adversary \mathcal{A} and a challenger:

Computational Diffie-Hellman: $\text{CDH}_{\mathcal{A}, \text{Gen}}(n)$

- Challenger runs $(G, q, g) \leftarrow \text{Gen}(1^n)$, $x, y \leftarrow \mathbb{Z}_q$, sends $(G, q, g, h_1 = g^x, h_2 = g^y)$ to \mathcal{A}
- \mathcal{A} outputs $h_3 \in G$
- We say that $\text{CDH}_{\mathcal{A}, \text{Gen}}(n) = 1$ (i.e., \mathcal{A} wins) if $h_3 = g^{xy}$.

Definition: CDH is hard if for all PPT \mathcal{A} : $\Pr[\mathcal{A} \text{ wins}] \leq \text{negl}(n)$

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

Variant 2: Decisional Diffie-Hellman

- Hard to distinguish g^{xy} from a random element in G

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

Variant 2: Decisional Diffie-Hellman

- Hard to distinguish g^{xy} from a random element in G

Consider the following game between an adversary \mathcal{A} and a challenger:

Decisional Diffie-Hellman: $\text{DDH}_{\mathcal{A}, \text{Gen}}(n)$

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

Variant 2: Decisional Diffie-Hellman

- Hard to distinguish g^{xy} from a random element in G

Consider the following game between an adversary \mathcal{A} and a challenger:

Decisional Diffie-Hellman: $\text{DDH}_{\mathcal{A}, \text{Gen}}(n)$

- Challenger runs $(G, q, g) \leftarrow \text{Gen}(1^n)$, $x, y \leftarrow \mathbb{Z}_q$, $b \leftarrow \{0, 1\}$

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

Variant 2: Decisional Diffie-Hellman

- Hard to distinguish g^{xy} from a random element in G

Consider the following game between an adversary \mathcal{A} and a challenger:

Decisional Diffie-Hellman: $\text{DDH}_{\mathcal{A}, \text{Gen}}(n)$

- Challenger runs $(G, q, g) \leftarrow \text{Gen}(1^n)$, $x, y \leftarrow \mathbb{Z}_q$, $b \leftarrow \{0, 1\}$
- If $b = 0$, send $(G, q, g, g^x, g^y, g^{xy})$ to \mathcal{A} .
If $b = 1$, choose $z \leftarrow \mathbb{Z}_q$, and send (G, q, g, g^x, g^y, g^z) to \mathcal{A}

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

Variant 2: Decisional Diffie-Hellman

- Hard to distinguish g^{xy} from a random element in G

Consider the following game between an adversary \mathcal{A} and a challenger:

Decisional Diffie-Hellman: $\text{DDH}_{\mathcal{A}, \text{Gen}}(n)$

- Challenger runs $(G, q, g) \leftarrow \text{Gen}(1^n)$, $x, y \leftarrow \mathbb{Z}_q$, $b \leftarrow \{0, 1\}$
- If $b = 0$, send $(G, q, g, g^x, g^y, g^{xy})$ to \mathcal{A} .
If $b = 1$, choose $z \leftarrow \mathbb{Z}_q$, and send (G, q, g, g^x, g^y, g^z) to \mathcal{A}
- \mathcal{A} outputs bit b'

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

Variant 2: Decisional Diffie-Hellman

- Hard to distinguish g^{xy} from a random element in G

Consider the following game between an adversary \mathcal{A} and a challenger:

Decisional Diffie-Hellman: $\text{DDH}_{\mathcal{A}, \text{Gen}}(n)$

- Challenger runs $(G, q, g) \leftarrow \text{Gen}(1^n)$, $x, y \leftarrow \mathbb{Z}_q$, $b \leftarrow \{0, 1\}$
- If $b = 0$, send $(G, q, g, g^x, g^y, g^{xy})$ to \mathcal{A} .
If $b = 1$, choose $z \leftarrow \mathbb{Z}_q$, and send (G, q, g, g^x, g^y, g^z) to \mathcal{A}
- \mathcal{A} outputs bit b'
- We say that $\text{DDH}_{\mathcal{A}, \text{Gen}}(n) = 1$ (i.e., \mathcal{A} wins) if $b = b'$.

Diffie-Hellman Assumptions

Let G be a cyclic group of order q with generator g

Diffie-Hellman Problem

Given $h_1 = g^x$, $h_2 = g^y$, find g^{xy}

Variant 2: Decisional Diffie-Hellman

- Hard to distinguish g^{xy} from a random element in G

Consider the following game between an adversary \mathcal{A} and a challenger:

Decisional Diffie-Hellman: $\text{DDH}_{\mathcal{A}, \text{Gen}}(n)$

- Challenger runs $(G, q, g) \leftarrow \text{Gen}(1^n)$, $x, y \leftarrow \mathbb{Z}_q$, $b \leftarrow \{0, 1\}$
- If $b = 0$, send $(G, q, g, g^x, g^y, g^{xy})$ to \mathcal{A} .
If $b = 1$, choose $z \leftarrow \mathbb{Z}_q$, and send (G, q, g, g^x, g^y, g^z) to \mathcal{A}
- \mathcal{A} outputs bit b'
- We say that $\text{DDH}_{\mathcal{A}, \text{Gen}}(n) = 1$ (i.e., \mathcal{A} wins) if $b = b'$.

Definition: DDH is hard if for all PPT \mathcal{A} : $\Pr[\mathcal{A} \text{ wins}] \leq 1/2 + \text{negl}(n)$

Relationship Between Cyclic Group Assumptions

Relative Hardness of The Problems

$DLog > CDH > DDH$

Relationship Between Cyclic Group Assumptions

Relative Hardness of The Problems

$DLog > CDH > DDH$

1 DLog vs. CDH

DLog: Given g^x find x

CDH: Given g^x, g^y find g^{xy}

$$\begin{array}{c} \downarrow \\ x \end{array} \quad (g^y)^x = g^{xy}$$

Relationship Between Cyclic Group Assumptions

Relative Hardness of The Problems

$DLog > CDH > DDH$

① DLog vs. CDH

- If \mathcal{A} can solve DLog, then given g^x, g^y , he can find x, y and compute g^{xy} (solve CDH)
- The reverse direction doesn't seem true

Relationship Between Cyclic Group Assumptions

Relative Hardness of The Problems

$DLog > CDH > DDH$

① DLog vs. CDH

- If \mathcal{A} can solve DLog, then given g^x, g^y , he can find x, y and compute g^{xy} (solve CDH)
- The reverse direction doesn't seem true

② CDH vs. DDH

CDH: $g^x, g^y \rightarrow g^{xy}$

DDH: g^x, g^y, g^{xz} look random

Relationship Between Cyclic Group Assumptions

Relative Hardness of The Problems

$DLog > CDH > DDH$

① DLog vs. CDH

- If \mathcal{A} can solve DLog, then given g^x, g^y , he can find x, y and compute g^{xy} (solve CDH)
- The reverse direction doesn't seem true

② CDH vs. DDH

- If \mathcal{A} given g^x, g^y can find g^{xy} , then he can distinguish this from g^z for a random $z \leftarrow \mathbb{Z}_q$ (solve DDH)
- The reverse direction doesn't seem true

Relationship Between Cyclic Group Assumptions

Relative Hardness of The Problems

$$DLog > CDH > DDH$$

1 DLog vs. CDH

- If \mathcal{A} can solve DLog, then given g^x, g^y , he can find x, y and compute g^{xy} (solve CDH)
- The reverse direction doesn't seem true

2 CDH vs. DDH

- If \mathcal{A} given g^x, g^y can find g^{xy} , then he can distinguish this from g^z for a random $z \leftarrow \mathbb{Z}_q$ (solve DDH)
- The reverse direction doesn't seem true

Strength of Assumption

Since DDH is the easiest problem, assuming it is secure is the strongest assumption

Preference for Prime-Order Groups

Note that \mathbb{Z}_p^* for p prime, $p > 2$, has order $p - 1$

- $p - 1$ is not prime
- So $G = \mathbb{Z}_p^*$ is not a prime-order group

Preference for Prime-Order Groups

Note that \mathbb{Z}_p^* for p prime, $p > 2$, has order $p - 1$

- $p - 1$ is not prime
- So $G = \mathbb{Z}_p^*$ is not a prime-order group

Reasons to prefer prime-order groups:

- Easy to find a generator, all $g \in G$ are generators.

Preference for Prime-Order Groups

Note that \mathbb{Z}_p^* for p prime, $p > 2$, has order $p - 1$

- $p - 1$ is not prime
- So $G = \mathbb{Z}_p^*$ is not a prime-order group

Reasons to prefer prime-order groups:

- Easy to find a generator, all $g \in G$ are generators.
- DDH is easy in composite-order groups!!!

Preference for Prime-Order Groups

Note that \mathbb{Z}_p^* for p prime, $p > 2$, has order $p - 1$

- $p - 1$ is not prime
- So $G = \mathbb{Z}_p^*$ is not a prime-order group

Reasons to prefer prime-order groups:

- Easy to find a generator, all $g \in G$ are generators.
- DDH is easy in composite-order groups!!!

Group of Quadratic Residues mod p

Preference for Prime-Order Groups

Note that \mathbb{Z}_p^* for p prime, $p > 2$, has order $p - 1$

- $p - 1$ is not prime
- So $G = \mathbb{Z}_p^*$ is not a prime-order group

Reasons to prefer prime-order groups:

- Easy to find a generator, all $g \in G$ are generators.
- DDH is easy in composite-order groups!!!

Group of Quadratic Residues mod p

Let $p = 2q + 1$ with both p and q prime.

Preference for Prime-Order Groups

Note that \mathbb{Z}_p^* for p prime, $p > 2$, has order $p - 1$

- $p - 1$ is not prime
- So $G = \mathbb{Z}_p^*$ is not a prime-order group

$$\begin{aligned} x^2, g^2 \\ x^2 \cdot g^2 &= (x \cdot g)^2 \\ |\mathbb{Z}_p^*| = p - 1 = 2q \end{aligned}$$

Reasons to prefer prime-order groups:

- Easy to find a generator, all $g \in G$ are generators.
- DDH is easy in composite-order groups!!!

Group of Quadratic Residues mod p

Let $p = 2q + 1$ with both p and q prime.

$$G = \{[h^2 \bmod p] \mid h \in \mathbb{Z}_p^*\}$$

is a group of prime order q

Preference for Prime-Order Groups

Note that \mathbb{Z}_p^* for p prime, $p > 2$, has order $p - 1$

- $p - 1$ is not prime
- So $G = \mathbb{Z}_p^*$ is not a prime-order group

Reasons to prefer prime-order groups:

- Easy to find a generator, all $g \in G$ are generators.
- DDH is easy in composite-order groups!!!

Given $g, h = g^x$ find x

$$g^x = \underbrace{g + g + g + \dots + g}_x =$$
$$= gx$$

Group of Quadratic Residues mod p

Let $p = 2q + 1$ with both p and q prime.

$$G = \{[h^2 \bmod p] \mid h \in \mathbb{Z}_p^*\}$$

is a group of prime order q