

**WARNING: MISBEHAVIOR AT EXAM TIME WILL LEAD TO SERIOUS CONSEQUENCE.**

## Probability and Statistics Exam paper B (2021–2022-2)

- Notice:**
1. Make sure that you have filled the form on the left side of seal line.
  2. Write your answers on the exam answer sheet .
  3. This is a close-book exam.
  4. The exam with full score of 100 points lasts 120 minutes.

Question No.	I	II	III	IV	V	VI	VII	VIII	IX	Sum
Score										

I. (10 points) We roll a die  $n$  times. Let  $A_{ij}$  be the event that the  $i$  th and  $j$  th rolls produce the same number. Show that the events  $\{A_{ij} : 1 \leq i < j \leq n\}$  are pairwise independent but not independent.

Score

**Solution.**

Suppose  $i < j$  and  $m < n$ . If  $j < m$ , then  $A_{ij}$  and  $A_{mn}$  are determined by distinct independent rolls, and are therefore independent. For the case  $j = m$  we have that

$$P(A_{ij} \cap A_{jn}) = P(i \text{ th, } j \text{ th, and } n \text{ th rolls show same number}).$$

$$= \sum_{r=1}^6 \frac{1}{6} P(j \text{ th and } n \text{ th rolls both show } r \mid i \text{ th shows } r) = \frac{1}{36} = P(A_{ij}) P(A_{jn})$$

as required. However, if  $i \neq j \neq k$ ,

$$P(A_{ij} \cap A_{jk} \cap A_{ik}) = \frac{1}{36} \neq \frac{1}{216} = P(A_{ij}) P(A_{jk}) P(A_{ik}).$$

II. (10 points) Two fair dice are rolled.

- (a) Compute the probability of the two dice have different scores.
- (b) Show that the event that their sum is 7 is independent of the score shown by the first die.

Score

**Solution.**

- (a) The probability of the two dice have different scores is  $\frac{6 \cdot 5}{6^2} = \frac{5}{6}$ .
- (b) It is implied by

$$P(1 \text{ st shows } r \text{ and sum is } 7) = \frac{1}{36} = \frac{1}{6} \cdot \frac{1}{6} = P(1 \text{ st shows } r) P(\text{sum is } 7).$$

III. (10 points) Individuals **A** and **B** begin to play a sequence of chess games. Let  $S = \{A \text{ wins a game}\}$ , and suppose that outcomes of successive games are independent with  $P(S) = p$  and  $P(F) = 1 - p$  (they never draw). They will play until one of them wins ten games. Let  $X$  = the number of games played (with possible values 10, 11,  $\dots$ , 19).

Score
-------

(a) For  $x = 10, 11, \dots, 19$ , obtain an expression for  $p(x) = P(X = x)$

(b) If a draw is possible, with  $p = P(S), q = P(F), 1 - p - q = P(\text{draw})$ , what is  $P(20 \leq X)$ ?

**Solution.**

(a)  $P(X = x) = P(\text{A wins in } x \text{ games}) + P(\text{B wins in } x \text{ games})$

$$\begin{aligned} &= P(9S' \text{ s in } 1^{\text{st}} x - 1 \cap S \text{ on the } x^{\text{th}}) + P(9F' \text{ s in } 1^{\text{st}} x - 1 \cap F \text{ on the } x^{\text{th}}) \\ &= \binom{x-1}{9} p^9 (1-p)^{(x-1)-9} \cdot p + \binom{x-1}{9} (1-p)^9 p^{(x-1)-9} \cdot (1-p) \\ &= \binom{x-1}{9} [p^{10} (1-p)^{x-10} + (1-p)^{10} p^{x-10}] \end{aligned}$$

(b) Possible values of  $X$  are now all positive integers  $\geq 10 : 10, 11, 12, \dots$

Similar to case (a), we have

$P(X = x) = P(\text{A wins in } x \text{ games}) + P(\text{B wins in } x \text{ games})$

$= P(9S' \text{ s in } 1^{\text{st}} x - 1 \cap S \text{ on the } x^{\text{th}}) + P(9F' \text{ s in } 1^{\text{st}} x - 1 \cap F \text{ on the } x^{\text{th}})$

$$= \binom{x-1}{9} p^9 (1-p)^{(x-1)-9} \cdot p + \binom{x-1}{9} q^9 (1-q)^{(x-1)-9} \cdot q$$

$$= \binom{x-1}{9} [p^{10} (1-p)^{x-10} + q^{10} (1-q)^{x-10}] \text{ . Finally,}$$

$$P(X \geq 20) = 1 - P(X < 20) = 1 - \sum_{x=10}^{19} \binom{x-1}{9} [p^{10} (1-p)^{x-10} + q^{10} (1-q)^{x-10}] \text{ .}$$

IV. (10 points) A 12-in. bar that is clamped at both ends is to be subjected to an increasing amount of stress until it snaps. Let  $Y$  = the distance from the left end at which the break occurs. Suppose  $Y$  has pdf

Score
-------

$$f(y) = \begin{cases} \left(\frac{1}{24}\right) y \left(1 - \frac{y}{12}\right) & 0 \leq y \leq 12 \\ 0 & \text{otherwise} \end{cases}$$

Compute the following:

(a) The cdf of  $Y$ .

(b) The expected length of the shorter segment  $\min(Y, 12 - Y)$  when the break occurs.

**Solution.**

(a) For  $0 \leq y \leq 12, F(y) = \frac{1}{24} \int_0^y \left(u - \frac{u^2}{12}\right) du = \frac{1}{24} \left(\frac{u^2}{2} - \frac{u^3}{36}\right) \Big|_0^y = \frac{y^2}{48} - \frac{y^3}{864}$ .

(b) The shorter segment has length equal to  $\min(Y, 12 - Y)$ , and

$$\begin{aligned} E[\min(Y, 12 - Y)] &= \int_0^{12} \min(y, 12 - y) \cdot f(y) dy = \int_0^6 \min(y, 12 - y) \cdot f(y) dy \\ &+ \int_6^{12} \min(y, 12 - y) \cdot f(y) dy = \int_0^6 y \cdot f(y) dy + \int_6^{12} (12 - y) \cdot f(y) dy = \frac{90}{24} = 3.75 \text{ inches.} \end{aligned}$$

V. (20 points) Suppose that  $X$  and  $Y$  are two independent rv's, both of which has uniform distribution in  $(0, 2)$ .

Score

- Determine the joint pdf of  $X$  and  $Y$ .
- Compute the probability  $P(X + Y \leq 1)$ .
- Compute the probability  $P(X \leq Y)$ .
- Compute  $V(X - Y)$ .

**Solution.**

(a) Since  $X$  and  $Y$  are independent, their joint pdf is

$$f(x, y) = \begin{cases} f_X(x)f_Y(y) = \frac{1}{4}, & 0 \leq x \leq 2, 0 \leq y \leq 2; \\ 0, & \text{otherwise.} \end{cases}$$

(b)  $P(X + Y \leq 1) = \int_0^1 \left[ \int_0^{1-x} \frac{1}{4} dy \right] dx = \frac{1}{4} \times \frac{1}{2} = \frac{1}{8}.$

(c)  $P(X \leq Y) = \int_0^2 \left[ \int_x^2 \frac{1}{4} dy \right] dx = \frac{1}{4} \times 2 = \frac{1}{2}.$

(d)  $V(X - Y) = V(X) + V(Y) = 2V(X) = 2 \times \frac{2^2}{12} = \frac{2}{3}.$

VI. (10 points) Suppose the expected tensile strength of type-A steel is 100ksi and the standard deviation of tensile strength is 8ksi. For type-B steel, suppose the expected tensile strength is 95ksi and the standard deviation of tensile strength is 7ksi, respectively. Let  $\bar{X}$  = the sample average tensile strength of a random sample of 40 type-A specimens, and let  $\bar{Y}$  = the sample average tensile strength of a random sample of 35 type-B specimen. Use the Central Limit Theorem to answer the following questions.

Score

- What are the approximate distributions of  $\bar{X}$  and  $\bar{Y}$  respectively?
- Calculate  $P(\bar{X} - \bar{Y} \geq 10)$ . ( $\Phi(1.65) = 0.95$ ,  $\Phi(2.89) = 0.998$ ,  $\Phi(1.96) = 0.975$ .)

**Solution.**

(a) According to the CLT,  $\bar{X}$  has approximately a normal distribution  $N\left(100, \frac{8^2}{40}\right)$ , i.e.  $N(100, 1.6)$ ,  $\bar{Y}$  has approximately a normal distribution  $N\left(95, \frac{7^2}{35}\right)$ , i.e.  $N(95, 1.4)$ .

(b) According to the CLT,  $\bar{X} - \bar{Y}$  has approximately  $N(5, 3)$ .  $P(\bar{X} - \bar{Y} \geq 10) = 1 - P(\bar{X} - \bar{Y} < 10) \approx 1 - \Phi\left(\frac{10-5}{\sqrt{3}}\right) = 1 - \Phi\left(\frac{5}{\sqrt{3}}\right) = 1 - \Phi(2.89) = 1 - 0.998 = 0.002.$

VII. (10 points) Let  $X_1, X_2, \dots, X_n$  be a random sample of size  $n$  from the pdf

Score

$$f(x; \theta) = \begin{cases} \frac{x}{\theta} e^{-\frac{x^2}{2\theta}}, & x > 0 \\ 0, & \text{otherwise} \end{cases}$$

- Use the method of moments to find an estimator for  $\theta$ .
- Find the maximum likelihood estimator for  $\theta$ .

**Solution.**

(a)

$$\begin{aligned} E(X) &= \int_{-\infty}^{+\infty} x f(x, \theta) dx = \int_0^{+\infty} \frac{x^2}{\theta} e^{-\frac{x^2}{2\theta}} dx \\ &= \int_0^{+\infty} x e^{-\frac{x^2}{2\theta}} d\frac{x^2}{2\theta} = x \left( -e^{-\frac{x^2}{2\theta}} \right) \Big|_0^{+\infty} + \int_0^{+\infty} e^{-\frac{x^2}{2\theta}} dx \\ &= 0 + \frac{1}{2} \int_{-\infty}^{+\infty} e^{-\frac{x^2}{2\theta}} dx = \sqrt{\frac{\theta \cdot \pi}{2}} \\ \therefore \theta &= \frac{2E^2(x)}{\pi}. \end{aligned}$$

(or)

$$\begin{aligned} E(X) &= \int_{-\infty}^{+\infty} x f(x, \theta) dx = \int_0^{+\infty} \frac{x^2}{\theta} e^{-\frac{x^2}{2\theta}} dx = \frac{1}{2} \int_{-\infty}^{+\infty} \frac{x^2}{\theta} e^{-\frac{x^2}{2\theta}} dx \\ &= \frac{\sqrt{2\pi}}{2\sqrt{\theta}} \int_{-\infty}^{+\infty} \frac{x^2}{\sqrt{2\pi}\sqrt{\theta}} e^{-\frac{x^2}{2\theta}} dx = \sqrt{\frac{\pi}{2\theta}} \cdot V(Y), \quad (Y \sim N(0, \theta)) \\ &= \sqrt{\frac{\pi}{2\theta}} \cdot (\sqrt{\theta})^2 = \sqrt{\frac{\theta \cdot \pi}{2}} \\ \therefore \theta &= \frac{2E^2(x)}{\pi}. \end{aligned}$$

Thus, the moment estimator is  $\hat{\theta}_M = \frac{2}{\pi} \bar{X}^2$ .

(or using 2nd moment)

$$\begin{aligned} E(X^2) &= \int_{-\infty}^{+\infty} x^2 f(x, \theta) dx = \int_0^{+\infty} \frac{x^3}{\theta} e^{-\frac{x^2}{2\theta}} dx \\ &= \int_0^{+\infty} x^2 e^{-\frac{x^2}{2\theta}} d\frac{x^2}{2\theta} = 2\theta \cdot \int_0^{+\infty} \frac{x^2}{2\theta} e^{-\frac{x^2}{2\theta}} d\left(\frac{x^2}{2\theta}\right) \\ &= 2\theta \cdot \int_0^{+\infty} u e^{-u} du = 2\theta \\ \therefore \theta &= \frac{E(X^2)}{2} \end{aligned}$$

The moment estimator is  $\hat{\theta}_M = \frac{1}{2n} \sum_{i=1}^n X_i^2$ .

(b) The likelihood function is

$$L(\theta) = L(x_1, \dots, x_n, \theta) = \prod_{i=1}^n \frac{x_i}{\theta} \cdot e^{-\sum_{i=1}^n \frac{x_i^2}{2\theta}} \quad (x_i > 0, i = 1, \dots, n).$$

The ln(likelihood) is

$$\ln L(\theta) = \sum_{i=1}^n \ln\left(\frac{x_i}{\theta}\right) - \sum_{i=1}^n \frac{x_i^2}{2\theta} = \sum_{i=1}^n \ln x_i - n \ln \theta - \sum_{i=1}^n \frac{x_i^2}{2\theta} \quad (x_i > 0, i = 1, \dots, n).$$

Let

$$\frac{d \ln L(x_1, \dots, x_n, \theta)}{d\theta} = \frac{-n}{\theta} + \frac{1}{\theta^2} \sum_{i=1}^n \frac{x_i^2}{2} = 0.$$

We get  $\theta = \frac{1}{2n} \sum_{i=1}^n x_i^2$ . So the maximum likelihood estimator of  $\theta$  is

$$\hat{\theta}_L = \frac{1}{2n} \sum_{i=1}^n X_i^2.$$

VIII. (10 points) It is reported that for a sample of 49 kitchens with gas cooking appliances monitored during a one-week period, the sample mean CO<sub>2</sub> level (ppm) was 654. Suppose that the population of CO<sub>2</sub> level of all homes is normal.

Score

(a) Calculate a 95% confidence interval for true average CO<sub>2</sub> level with the sample standard deviation  $s = 168$ . ( $t_{0.05,48} = 1.68$ ,  $t_{0.025,48} = 2.0$ .)

(b) Suppose that  $\sigma = 175$ . What sample size would be necessary to obtain an interval width of at most 50ppm for a confidence level of 95%? ( $z_{0.05} = 1.65$ ,  $z_{0.025} = 1.96$ .)

**Solution.**

(a) With  $n = 49$ ,  $\bar{x} = 654$  and  $t_{\alpha/2, n-1} = t_{0.025, 48} = 2.0$ , the 95% confidence interval for  $\mu$  is

$$\bar{x} \pm t_{\alpha/2, n-1} \frac{s}{\sqrt{n}} = 654 \pm 2.0 \frac{168}{7} = 654 \pm 48 = (606, 702).$$

(b) With  $\sigma = 175$  and  $\mu \in \bar{x} \pm z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$ , the width of CI is  $w = 2z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$ . So

$$n = \left( \frac{2z_{\alpha/2}\sigma}{w} \right)^2 = \left( \frac{2(1.96)(175)}{50} \right)^2 = (13.72)^2 = 188.24,$$

which rounds up to 189.

IX. (10 points) The desired percentage of SiO<sub>2</sub> in a certain type of aluminous cement is 5.5. In a test 16 independently obtained samples are analyzed. Suppose that the percentage of SiO<sub>2</sub> is normally distributed with  $\sigma = 0.3$  and that  $\bar{x} = 5.25$ .

Score

(a) Does this indicate conclusively that the true average percentage less than 5.5? Consider a significance level of  $\alpha = 0.01$ . ( $z_{0.01} = 2.33$ ,  $z_{0.005} = 2.58$ .)

(b) If the true average percentage is  $\mu = 5.3$  and a level  $\alpha = 0.01$  test based on  $n = 16$  is used, what is the probability of rejecting  $H_0$ ? ( $\Phi(0.34) = 0.63$ ,  $\Phi(0.64) = 0.74$ .)

**Solution.**

The hypotheses are  $H_0 : \mu = 5.5$  vs  $H_a : \mu < 5.5$ . The sample mean is  $\bar{x} = 5.25$ .

(a)  $z = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}} = \frac{5.25 - 5.5}{0.3/\sqrt{16}} = -3.33 \leq -z_{\alpha} = -z_{0.01} = -2.33$ . Reject  $H_0$ .

(b) The probability of making a type II error when  $\mu = 5.3$  is

$$\beta(5.3) = 1 - \Phi \left( -z_{\alpha} + \frac{\mu_0 - \mu'}{\sigma/\sqrt{n}} \right) = 1 - \Phi \left( -2.33 + \frac{5.5 - 5.3}{0.3/\sqrt{16}} \right) = 1 - \Phi(0.34).$$

So the probability of rejecting  $H_0$  is  $1 - \beta(5.3) = \Phi(0.34) = 0.63$ .