

Puffin: New System of Equations

November 17, 2015

Compared to the original POP Puffin paper, now we use the peak magnetic fields are used to scale the system of eqns (as opposed to the *rms* values used previously). Instead of p_2 , the energy exchange is modelled directly through the scaled variable $\Gamma_j = \gamma_j/\gamma_r$, so that:

$$\left[\frac{1}{2} \left(\frac{\partial^2}{\partial \bar{x}^2} + \frac{\partial^2}{\partial \bar{y}^2} \right) - \frac{\partial^2}{\partial \bar{z} \partial \bar{z}_2} \right] A_\perp = -\frac{1}{\bar{n}_p} \frac{\partial}{\partial \bar{z}_2} \sum_{j=1}^N \frac{\bar{p}_{\perp j}}{\Gamma_j} (1 + \eta p_{2j}) \delta^3(\bar{x}_j, \bar{y}_j, \bar{z}_{2j}) \quad (1)$$

$$\frac{d\bar{p}_{\perp j}}{d\bar{z}} = \frac{1}{2\rho} \left[i b_\perp - \frac{\eta p_{2j}}{\kappa^2} A_\perp \right] - i \kappa \frac{\bar{p}_{\perp j}}{\Gamma_j} (1 + \eta p_{2j}) b_z \quad (2)$$

$$\frac{d\Gamma_j}{d\bar{z}} = -\rho \frac{(1 + \eta p_{2j})}{\Gamma_j} (\bar{p}_{\perp j} A_{\perp j}^* + c.c.) \quad (3)$$

$$\frac{d\bar{z}_{2j}}{d\bar{z}} = p_{2j} \quad (4)$$

$$\frac{d\bar{x}_j}{d\bar{z}} = \frac{2\rho\kappa}{\sqrt{\eta}\Gamma_j} (1 + \eta p_{2j}) \Re(\bar{p}_{\perp j}) \quad (5)$$

$$\frac{d\bar{y}_j}{d\bar{z}} = -\frac{2\rho\kappa}{\sqrt{\eta}\Gamma_j} (1 + \eta p_{2j}) \Im(\bar{p}_{\perp j}). \quad (6)$$

where $\kappa = \frac{a_w}{2\rho\gamma_r}$, which is actually the scaled natural focussing wavenumber $\kappa = \sqrt{2k_\beta}$.

p_{2j} may be worked out analytically from

$$p_{2j} = \frac{1}{\eta} \left[\left(1 - \frac{(1 + a_w^2 |\bar{p}_{\perp j}|^2)}{\gamma_r^2 \Gamma_j^2} \right)^{-1/2} - 1 \right] \quad (7)$$

at each step. It is NOT output from Puffin.

Outputs from Puffin are the scaled radiation field A_\perp , and electron phase space coords $\bar{x}, \bar{y}, \bar{z}_2, \bar{p}_\perp, \Gamma$, and scaled distance through the undulator \bar{z} . The normalized electron weights are in the file NormChiDataFile.dat (normalised to the peak *spatial* (3D) density).

Scaled parameters are:-

$$\begin{aligned}
\bar{z}_{2j} &= \frac{ct_j - z}{l_c}, & \bar{z} &= \frac{z}{l_g}, \\
\bar{p}_\perp &= \frac{p_\perp}{mca_u}, & A_\perp &= \frac{ea_ul_g}{2mc^2\gamma_0^2\rho}E_\perp, \\
(\bar{x}, \bar{y}) &= \frac{(x, y)}{\sqrt{l_g l_c}}, & l_g &= \frac{\lambda_w}{4\pi\rho}, \\
l_c &= \frac{\lambda_r}{4\pi\rho}, & \Gamma_j &= \frac{\gamma_j}{\gamma_0}, \\
\rho &= \frac{1}{\gamma_0} \left(\frac{a_u \omega_p}{4ck_u} \right)^{2/3}, & a_u &= \frac{eB_0}{mck_u}, \\
\kappa &= \frac{a_u}{2\rho\gamma_0}, & b_\perp &= b_x - ib_y,
\end{aligned}$$

B_0 is the peak magnetic field in the wiggler. $\omega_p = \sqrt{e^2 n_p / \epsilon_0 m}$ is the (non-relativistic) plasma frequency, and n_p is the peak spatial number density of the electron beam ($N_e / \delta_x \delta_y \delta_z$). $E_\perp = E_x - iE_y$ are the x and y radiation electric field vectors. γ_0 is the reference energy (Lorentz factor), which is usually taken as the mean beam energy.

The scaled reference velocity,

$$\eta = \frac{1 - \beta_{zr}}{\beta_{zr}} = \frac{\lambda_r}{\lambda_u} = \frac{l_c}{l_g}, \quad (8)$$

where β_{zr} is some reference velocity scaled to c , which is sensible (but not, strictly speaking, necessary) to take as the mean longitudinal electron velocity in the wiggler, so that

$$\beta_{zr} = \sqrt{\left(1 - \frac{1}{\gamma_r^2} \left(1 + \bar{a}_u^2\right)\right)^{-1/2} - 1}, \quad (9)$$

where \bar{a}_u is the *rms* undulator parameter. This defines the velocity at which the electrons travel in the scaled \bar{z}_2 frame. More generally, η describes an *ideal* resonance condition - electrons resonant with wavelength λ_r will travel with velocity $p_2 = 1$ through the \bar{z}_2 frame.