

1 Supplement — Construction of the *Methanococcus maripaludis*
2 etc-GEM, the Sharpe–Schoolfield E_a audit, and robustness

3 Gabriel Yvon-Durocher
4 *Living Systems Institute, University of Exeter, Stocker Road, Exeter EX4 4QD, UK*
5 2026-07-12

6 **Table of contents**

7	1 Building the methanogen etc-GEM (M1–M6)	1
8	1.1 M1 — base GEM + medium (foundation)	1
9	1.2 M1b — carbon-honest curation	2
10	1.3 M2 / M2b — the sMOMENT ecModel + kcat refinement	2
11	1.4 M3 — thermal envelope + measured maintenance	2
12	1.5 M4 — Bayesian calibration to Jones 1983	3
13	1.6 M6 — the methanogen’s own allocation layer	3
14	2 Why Sharpe–Schoolfield E: the window-sensitivity audit	4
15	3 Mcr-k_{cat} sensitivity	5

16 This Supplement documents the construction of the *M. maripaludis* methanogenesis etc-GEM
17 (the *E. coli* respiration model is described in the companion report `reports/ecoli_tpc/`),
18 the window-independence audit that motivates the Sharpe–Schoolfield E , and the robustness
19 analyses. Numbers are model outputs; honest caveats are stated throughout.

20 **1 Building the methanogen etc-GEM (M1–M6)**

21 The model was built in six phases, each committed and validated (build records in
22 `strains/mmaripaludis/outputs/M*.md`).

23 **1.1 M1 — base GEM + medium (foundation)**

24 The iMR539 reconstruction¹ (688 reactions / 710 metabolites / 539 genes; BioModels
25 BIOMD0000001099) was acquired and validated on a defined H₂/CO₂ medium. Plain

FBA reproduces hydrogenotrophic methanogenesis with the correct **H2:CO2:CH4 = 4:1:1** stoichiometry and a gene-annotated Wolfe cycle. A preliminary audit found **no thermodynamic energy loops**; 85% metabolic GPR coverage; MMP locus tags map to UniProt proteome UP000000590 (sequences fetchable) — ecModel-ready.

1.2 M1b — carbon-honest curation

The published reconstruction could not grow without free uptake of four biomass precursors. Tracing the network, three (Membrane_lipid, NAC, Flagellin) feed *only* the archaellin (flagellum) glycoprotein — a dispensable motility structure — so archaellin was folded out of biomass; the fourth (tRNA(SeCys)) was a **recycling gap** (the incorporation step failed to release the free tRNA), fixed by regenerating it. Organic-carbon leaks were closed and the pinned CH4 exchange freed. The curated model grows autotrophically on H2/CO2 with **99.98% of biomass carbon from CO2**. Two vitamin cofactors (thiamin, NMN) lack biosynthesis and are supplied as documented medium traces (0.02% of carbon).

1.3 M2 / M2b — the sMOMENT ecModel + kcat refinement

An sMOMENT total-protein pool was attached (reusing the E. coli **enzyme_cost** engine): each reaction costs MW/k_{cat} from a shared budget. Per-reaction k_{cat} by source priority: **measured core** (electron-bifurcation complexes from Milton et al. 2018, 25–165 U/mg²; classic biochemistry for Mtr/Fwd/Mcr/etc.), **DLTKcat** sequence predictions for the bulk, and a mean fallback. Mcr — the famously slow terminal enzyme and the E_a target — was sourced carefully (mature-Mcr average specific activity ~20 U/mg, *Methanothermobacter*; k_{cat} 59 s⁻¹, uncertainty range 3–294 s⁻¹). **M2b** corrected a systematic DLTKcat archaeal-underprediction (28% of predictions <1 s⁻¹) with cited literature/BRENDA overrides for the flagged high-pool offenders (pyruvate- and 2-oxoglutarate:ferredoxin oxidoreductases) plus a documented 1 s⁻¹ floor. The grounded pool ($P_{total} \times \sigma \times f_{metab}$, not tuned to μ_{max}) gives an emergent μ that undershoots the observed rate ~4×, expected for a-priori kinetics.

1.4 M3 — thermal envelope + measured maintenance

$kcat(T)$ (MMRT/Eyring) $\times$ native fraction $f_N(T)$ (two-state unfolding keyed on T_m), with per-enzyme optima from a documented **mesophile Topt prior** and T_m from a mesophile prior (the sequence/meltome predictors are the future upgrade). Maintenance is a temperature-dependent NGAM(T) **anchored on the measured** M. maripaludis maintenance energy (7.836 mmol ATP gDW⁻¹ h⁻¹, Goyal et al. 2015³). The emergent TPC matches Jones' **CTmax** (~47 °C) a-priori (Figure 1); the rising limb was too warm/steep and the magnitude maintenance-crushed — the residuals the calibration then addresses.




Fig. 1 | Emergent (nothing-fit) methanogen TPC vs Jones 1983: the falling limb (CTmax) matches a-priori; the rising limb and magnitude are calibrated in M4.

1.5 M4 — Bayesian calibration to Jones 1983

The thermal ecModel was calibrated to the digitised Jones 1983 growth TPC⁴ by ensemble MCMC (emcee; the same machinery as the *E. coli* fit), magnitude-first, with provenance priors. The posterior **reproduces the whole curve** (Figure 2; RMSE 0.005 h⁻¹), reaching the peak (0.184 vs 0.181 h⁻¹) with Topt 37 °C and CTmax 51 °C. The demanded magnitude correction (kcat_scale 7×) is a plausible in-vitro→in-vivo + archaeal-underprediction gap (the single sMOMENT pool lumps the degenerate magnitude levers). The maintenance multiplier stayed near 1, confirming the Goyal-measured NGAM.

1.6 M6 — the methanogen’s own allocation layer

A proteome-sector/growth-law layer was added, grounded in the organism’s **own** allocation physiology: Müller et al. 2021⁵ show *M. maripaludis* holds a **constant ribosomal proteome fraction** and invariant catabolic/anabolic allocation across growth rate (the opposite of the *E. coli*/yeast scaling law), so the growth-law coupling slope is **~0 (flat)**. Sector fractions (f_{metab} 0.55 / f_{maint} 0.30 / f_{bio} 0.15) are grounded in the Xia/Müller proteome. Forward-check: with the sector layer on and the M4 posterior, the TPC is **identical** to the single-pool model ($\max|\Delta\mu| = 0$), so no re-calibration was needed. The directly-measured allocation buffer is **+0.000 eV** (vs *E. coli* -0.130), grounding the allocation asymmetry in each organism’s measured strategy.

Honest caveats carried into the analysis. (i) The Mcr k_{cat} (3–294 s⁻¹) is the dominant single-parameter uncertainty. (ii) **DLTKcat is bacteria-trained**, so archaeal predictions are corrected but remain a prior for the biosynthetic bulk. (iii) The T_m and **Topt** are mesophile priors (no archaeal sequence predictor). (iv) The **sector fractions** are grounded




Fig. 2 | Prior (emergent) vs posterior methanogen TPC against Jones 1983.

but approximate; the flat slope is the robust, cited quantity. The main-text conclusions are shown robust to the parameter that matters most (Mcr; Section 3).

2 Why Sharpe–Schoolfield E : the window-sensitivity audit

The naive Boltzmann–Arrhenius slope of the rising limb depends on the fitting window because the limb is curved. Table 1 recomputes it under five window conventions for every model and observed curve: the spread is **0.42–1.41 eV per curve**, larger than the effects we attribute, and enough to reverse the cross-organism ordering. The Sharpe–Schoolfield E (the high-temperature-inactivation model, main text) is read from the whole shape, is window-independent, and — the decisive test — makes model and observed agree for both organisms (Figure 3).

```
import pandas as pd
w = pd.read_csv("assets/tables/window_sensitivity.csv")
ss = pd.read_csv("assets/tables/sharpe_schoolfield_fits.csv")[["curve", "SS_E_eV", "SS_r2" if
m = w.merge(ss, on="curve", how="left")
print(m.round(3).to_markdown(index=False))
```

Table 1: Full window-sensitivity table: Boltzmann-Arrhenius slope E_a (eV) under five window conventions, and the window-independent Sharpe-Schoolfield E , for every model + observed curve.

curve	full_be-					spread_eV	SS_E_eV	r2
	rel_10_95	low	rel_10_80	rel_30_90	rel_10_50			
ecoli_emer- gent	0.609	0.64	0.802	0.489	1.054	0.565	0.458	0.896
ecoli_tuned	0.956	1.468	1.203	0.756	1.56	0.804	0.677	0.93
ecoli_ob- served_VdL	0.643	0.963	0.683	0.545	0.839	0.418	0.558	0.973
methan_cal- ibrated	1.174	1.462	1.372	0.892	1.785	0.893	1.024	0.957
methan_ob- served_Jones	0.66	2.067	nan	0.781	nan	1.407	1.042	0.95




Fig. 3 | Sharpe–Schoolfield fits overlaid on the model and observed curves for both organisms; the model and observed E agree.

91 3 $\text{Mcr-}k_{\text{cat}}$ sensitivity

92 Table 2 sweeps Mcr across its $3\text{--}294\text{ s}^{-1}$ range. The organism E_a stays **1.01–1.27 eV**, always
 93 **well above E. coli's 0.68**, and the methanogenesis backbone carries **83–103%** of the control
 94 term throughout — so the ordering and the backbone-control finding are robust. Only the
 95 identity of the single leading enzyme shifts (Mcr when slow $\rightarrow$ Fwd when fast).

```
import pandas as pd
m = pd.read_csv("assets/tables/mcr_sweep.csv")
print(m.round(3).to_markdown(index=False))
```

Table 2: Mcr-kcat sensitivity sweep: organism SS-E, backbone fraction of the control term, and the leading enzymes, across 3-294/s.

Mcr_kcat	SS_E	slope	back- bone_frac_of_con- trol	top3	Mcr_C_i	Mcr_con- trib
3	1.274	1.597	1.026	Mcr(1.605);Fwd(0.092);CODH(0.065)	1.391	1.605
10	1.01	1.105	0.986	Mcr(0.876);Fwd(0.167);CODH(0.039)	1.907	0.876
30	1.065	1.132	0.925	Mcr(0.488);Fwd(0.277);Mtr(0.059)	0.599	0.488
58.9	1.056	1.081	0.887	Fwd(0.337);Mcr(0.304);Mtr(0.077)	0.727	0.304
100	1.066	1.153	0.857	Fwd(0.395);Mcr(0.211);Mtr(0.078)	0.778	0.211
294	1.017	1.074	0.833	Fwd(0.427);Mtr(0.086);Mer(0.076)	0.766	0.078

1. Richards, M. A. *et al.* Exploring hydrogenotrophic methanogenesis: A genome scale metabolic reconstruction of *Methanococcus maripaludis*. *Journal of Bacteriology* **198**, 3379–3390 (2016).
2. Milton, R. D. *et al.* Measured specific activities and redox thermodynamics of the flavin-based electron-bifurcation hydrogenase and heterodisulfide-reductase complexes of *Methanococcus maripaludis*. *Biochemistry* (2018).
3. Goyal, N., Widiastuti, H., Karimi, I. A. & Zhou, Z. Flux measurements and maintenance energy for carbon dioxide utilization by *Methanococcus maripaludis*. *Microbial Cell Factories* **14**, 146 (2015).
4. Jones, W. J., Paynter, M. J. B. & Gupta, R. Characterization of *Methanococcus maripaludis* sp. Nov., a new methanogen isolated from salt marsh sediment. *Archives of Microbiology* **135**, 91–97 (1983).
5. Müller, A. L. *et al.* An alternative resource allocation strategy in the chemolithoautotrophic archaeon *Methanococcus maripaludis*. *Proceedings of the National Academy of Sciences* **118**, e2025854118 (2021).