```
iSAT = 1e-9;
iS = 0.1;
    = 20;
xa = 0.4;
xb = 0.5;
xx = xa : (xb-xa)/10000 : xb;
A = 2/3 * R * iSAT;
B = A + R * iS/3;
g1 = 0(x) x;
g2 = 0(x) B - A * exp(x./Vth);
g = 0(x) g1(x) - g2(x);
figure(1);
plot(xx, g1(xx), '-b', ...
    xx, g2(xx), '-r');
x0 = 0;
format long e;
xEX = fzero(g, x0)
Tg = @(x) Vth * log((B-x)./A);
dTg = @(x) -Vth / (B-x);
W = abs(dTg(xEX))
x0 = 0;
itMAX = 1000;
toll = 1e-12;
[x ,nIT, err] = fixed_point(x0,Tg,toll,itMAX);
% stimap(x);
xFIX
     = x(end)
estERR = err(end)
trueERR = xEX - xFIX
C = 1e-6;
t0 = 0;
tf = 0.1;
iSf = 0.1;
f = 100;
tD = 1e-2;
x0 = 0;
iS = 0(t) iSf * sin(2*pi*f * t) .* exp(-t ./ tD);
f = @(t,x) (-3*x) ./ (2*R*C) + (iSAT/C) * (1 - exp(x ./ Vth)) + iS(t) ./ (2*C);
[tm, xm] = ode15s(f, [t0, tf], x0);
deltaXa = max(xm) - min(xm)
dfdx = @(t,x) -3 ./ (2*R*C) - iSAT/(C*Vth) * exp(x ./ Vth);
NT
     = 1000:
tol
      = 1e-12;
maxtit = 100;
[tth,xth,iter_nwt] = crank_nicolson(f,dfdx,t0,tf,x0,(tf-t0)/NT);
deltaXb = max(xth) - min(xth)
```

constants();

```
figure(2);
plot(tm, xm, '-r', ...
    tth, xth, '-b');
legend('ode15s', 'crank\_nicolson');
```