



Coreceptor usage prediction from genotype (version 2.5)

I. General information

Patient:	Study Id:
Birth date:	Viral load:
Sample received:	Sample collected:
Sample IDs: 000001465805	Predicted V3-subtype: B
Sample type:	Report date: July 20, 2024
Physician:	Reported by:

II. Additional clinical parameters used for the predictions:

No clinical parameters given

III. Aligned V3 region

Consensus B: TGT ACA AGA CCC AAC AAC AAT ACA AGA AAA AGT ATA CAT ATA GGA CCA GGG AGA GCA TTT TAT ACA ACA GGA GAA ATA ATA GGA GAT ATA AGA CAA GCA CAT TGT
 Query: C T R P N N N T R K S I H I G P G R A F Y T T G E I I G D I R Q A H C
 C T R P N N N T I R S I T I G P G R K F H A T R R I I G D I R Q A H C
 TGT ACA AGA CCC AAC AAC AAT ACA ATA AGA AGC ATA ACT ATA GGA CCA GGG AGA AAA TTT CAT GCA ACA AGA CGC ATA ATA GGA GAT ATA AGA CAA GCA CAT TGT

IV. Coreceptor prediction

Model	Prediction	FPR	Remarks
Clonal	The CXCR4-coreceptor can be used. CCR5-antagonists like Maraviroc (Celsentri/Selzentry) should not be administered!	1.7%	The 11/25 rule would predict this sequence as an X4-virus.

The significance level was set to: 20% false positive rate

Comments:

(signature)

Mutações: R304I,K305R,R306S,R308T,del310,del311,A316K,317insH,V318A,I320R,G321R,K322I,N325D,M326I