



Coreceptor usage prediction from genotype (version 2.5)

I. General information

Patient:	Study Id:
Birth date:	Viral load:
Sample received:	Sample collected:
Sample IDs: 000001476170	Predicted V3-subtype: C
Sample type:	Report date: July 19, 2024
Physician:	Reported by:

II. Additional clinical parameters used for the predictions:

No clinical parameters given

III. Aligned V3 region

Consensus B: TGT ACA AGA CCC AAC AAC AAT ACA AGA AAA AGT ATA CAT ATA GGA CCA GGG AGA GCA TTT TAT ACA ACA GGA GAA ATA ATA GGA GAT ATA AGA CAA GCA CAT TGT
 Query: C T R P N N N T R K S I H I G P G R A F Y T T G E I I G D I R Q A H C
 TGT ACA CGA CCC AAC AAT AAT ACA AGA AAR AGT ATA AGG ATA GGA CCA GGA CAA GCA TTC TAT GCA ACA GGA GAC ATA ATA GGA GAT ATA AGA CAA GCC TAT TGY

IV. Coreceptor prediction

Model	Prediction	FPR	Remarks
Clonal	Only the CCR5-coreceptor can be used. CCR5-antagonists like Maraviroc (Celsentri/Selzentry) could work.	43%	

The significance level was set to: 20% false positive rate

Comments:

(signature)

Mutações: R306S,del307,del308,Q310R,R311I,R315Q,V318Y,T319A,I320T,K322D,322insI,N325D,M326I,H330Y