

RWorksheet_Benedicto#4a.R

Gabriel R. Benedicto

2024-10-14

1. The table below shows the data about shoe size and height. Create a data frame.

```
shoe_size <- c(6.5, 9.0, 8.5, 8.5, 7.0, 9.0, 9.5, 13.0, 7.5, 10.5, 10.5, 12.0, 10.5,
              13.0, 11.5, 8.5, 5.0, 10.0, 6.5, 7.5, 8.5, 10.5, 8.5, 10.5, 11.0, 9.0, 13.0)
height <- c(66.0, 68.0, 65.0, 65.0, 64.0, 71.0, 72.0, 72.0, 74.5, 67.0, 74.5, 71.0, 71.0,
           77.0, 72.0, 59.0, 62.0, 72.0, 66.0, 64.0, 67.0, 73.0, 69.0, 72.0, 70.0, 69.0, 70.0)
gender <- c("F", "F", "F", "F", "F", "F", "F", "M", "F", "M", "M", "M", "M",
           "M", "M", "F", "F", "M", "F", "F", "F", "M", "F", "M", "M", "M", "M")

data <- data.frame(Shoe_size = shoe_size, Height = height, Gender = gender)
```

data

##	Shoe_size	Height	Gender
## 1	6.5	66.0	F
## 2	9.0	68.0	F
## 3	8.5	65.0	F
## 4	8.5	65.0	F
## 5	7.0	64.0	F
## 6	9.0	71.0	F
## 7	9.5	72.0	F
## 8	13.0	72.0	M
## 9	7.5	74.5	F
## 10	10.5	67.0	M
## 11	10.5	74.5	M
## 12	12.0	71.0	M
## 13	10.5	71.0	M
## 14	13.0	77.0	M
## 15	11.5	72.0	M
## 16	8.5	59.0	F
## 17	5.0	62.0	F
## 18	10.0	72.0	M
## 19	6.5	66.0	F
## 20	7.5	64.0	F
## 21	8.5	67.0	F
## 22	10.5	73.0	M
## 23	8.5	69.0	F
## 24	10.5	72.0	M
## 25	11.0	70.0	M
## 26	9.0	69.0	M
## 27	13.0	70.0	M

- a. Describe the data. The table presents data about individuals' shoe size, height, and gender. The variables are:

-Shoe size: Numeric, represents the size of the shoe. -Height: Numeric, represents the height of the individual (presumably in inches). -Gender: Categorical, either “M” for male or “F” for female.

- b. Create a subset by males and females with their corresponding shoe size and height. What its result? Show the R scripts.

- The result is that it shows the Shoe-size and height of every Male and Female in the data.

```
male_data <- subset(data, Gender == "M", select = c(Shoe_size, Height))
```

```
male_data
```

```
##      Shoe_size Height
## 8          13.0    72.0
## 10         10.5    67.0
## 11         10.5    74.5
## 12         12.0    71.0
## 13         10.5    71.0
## 14         13.0    77.0
## 15         11.5    72.0
## 18         10.0    72.0
## 22         10.5    73.0
## 24         10.5    72.0
## 25         11.0    70.0
## 26          9.0    69.0
## 27         13.0    70.0
```

```
female_data <- subset(data, Gender == "F", select = c(Shoe_size, Height))
```

```
female_data
```

```
##      Shoe_size Height
## 1           6.5    66.0
## 2           9.0    68.0
## 3           8.5    65.0
## 4           8.5    65.0
## 5           7.0    64.0
## 6           9.0    71.0
## 7           9.5    72.0
## 9           7.5    74.5
## 16          8.5    59.0
## 17          5.0    62.0
## 19          6.5    66.0
## 20          7.5    64.0
## 21          8.5    67.0
## 23          8.5    69.0
```

- c. Find the mean of shoe size and height of the respondents. Write the R scripts and its result.

```
mean_shoe_size <- mean(data$Shoe_size)
```

```
mean_shoe_size
```

```
## [1] 9.444444
```

```
mean_height <- mean(data$Height)
```

```
mean_height
```

```
## [1] 69
```

d. Is there a relationship between shoe size and height? Why?

- Yes. for me from personal experience, as the shoesize increase, the height also increases.

2. Construct character vector `months` to a factor with `factor()` and assign the result to `factor_months_vector`. Print out `factor_months_vector` and assert that R prints out the factor levels below the actual values.

```
months_vector <- c("March", "April", "January", "November", "January",  
"September", "October", "September", "November", "August",  
"January", "November", "November", "February", "May", "August",  
"July", "December", "August", "August", "September", "November", "February",  
"April")
```

```
factor_months_vector <- factor(months_vector)
```

```
print(factor_months_vector)
```

```
## [1] March      April      January   November  January   September October  
## [8] September November  August    January   November  November  February  
## [15] May        August     July      December  August    August    September  
## [22] November  February  April  
## 11 Levels: April August December February January July March May ... September
```

```
levels(factor_months_vector)
```

```
## [1] "April"      "August"      "December"    "February"    "January"     "July"  
## [7] "March"      "May"         "November"    "October"     "September"
```

3. Then check the `summary()` of the `months_vector` and `factor_months_vector`. Interpret the results of both vectors. Are they both equally useful in this case? -The result of `months_vector` is that it states the Length, Class and Mode. While the `factor_months_vector` states how many months in the data for example April, April has 2. -I think they are both useful because it is easy for me to understand and determine how many types of data from the raw data itself.

```
summary(months_vector)
```

```
##      Length      Class      Mode  
##           24 character character
```

```
summary(factor_months_vector)
```

```
##      April      August  December  February   January      July      March      May  
##           2          4           1           2           3           1           1           1  
## November  October  September  
##           5          1           3
```

4. Create a vector and factor for the table below.

```
direction_vector <- c("East", "West", "North", "West", "North", "West", "North", "West")
```

```
factor_data <- factor(direction_vector)
```

```
new_order_data <- factor(factor_data, levels = c("East", "West", "North"))  
print(new_order_data)
```

```
## [1] East West North West North West North West  
## Levels: East West North
```

5. Enter the data below in Excel with file name = import_march.csv

a. Import the excel file into the Environment Pane using read.table() function. Write the code.

```
data_excel <- read.table("import_march.csv")
```

b. View the dataset. Write the R scripts and its result.

```
data_excel
```

```
##                               V1
## 1 Students,Strategy1,Strategy2,Strategy3
## 2                               Male,8,10,8
## 3                               ,4,8,6
## 4                               ,0,6,4
## 5                               Female,14,4,15
## 6                               ,10,2,12
## 7                               ,6,0,9
```

6. Full Search

```
exhaustive_search <- function(selected_number) {

  if(selected_number < 1 || selected_number > 50) {
    print("The selected number is beyond the range of 1 to 50")
  } else if(selected_number == 20) {
    print("TRUE")
  } else {
    print(selected_number)
  }
}

selected_number <- readline(prompt = "Select a number from 1 to 50: ")
```

```
## Select a number from 1 to 50:
```

```
exhaustive_search(selected_number)
```

```
## [1] "The selected number is beyond the range of 1 to 50"
```

7. Change

```
min_bills <- function(price) {
  bills <- c(1000, 500, 200, 100, 50)

  num_bills <- 0
  for(bill in bills) {
    count <- price %/% bill
    num_bills <- num_bills + count

    price <- price %% bill
  }

  print(paste("Minimum number of bills needed to purchase a snack: ", num_bills))
}

min_bills(1640)
```

```
## [1] "Minimum number of bills needed to purchase a snack: 3"
```

8. The following is each student's math score for one semester. Based on this, answer the following questions.

a. Create a dataframe from the above table. Write the R codes and its output.

```
students <- data.frame(
  Name = c("Annie", "Thea", "Steve", "Hanna"),
  Grade1 = c(85, 65, 75, 95),
  Grade2 = c(65, 75, 55, 75),
  Grade3 = c(85, 90, 80, 100),
  Grade4 = c(100, 90, 85, 90)
)
```

students

```
##      Name Grade1 Grade2 Grade3 Grade4
## 1 Annie      85      65      85      100
## 2 Thea       65      75      90      90
## 3 Steve      75      55      80      85
## 4 Hanna      95      75     100      90
```

b. Without using the rowMean function, output the average score of students whose average math score over 90 points during the semester. write R code and its output. Example Output: Annie's average grade this semester is 88.75.

```
for (i in 1:nrow(students)) {
  total_score <- students$Grade1[i] + students$Grade2[i] + students$Grade3[i] + students$Grade4[i]
  avg_score <- total_score / 4

  if (avg_score > 90) {
    formatted_output <- sprintf("%s's average grade this semester is %.2f.", students$Name[i], avg_score)
    print(formatted_output)
  }
}
```

C. Without using the mean function, output as follows for the tests in which the average score was less than 80 out of 4 tests. Example output: The nth test was difficult.

```
for (j in 2:5) {
  total_test_score <- sum(students[, j])
  avg_test_score <- total_test_score / nrow(students)

  if (avg_test_score < 80) {
    print(paste("The", j-1, "th test was difficult."))
  }
}
```

```
## [1] "The 2 th test was difficult."
```

D.. Without using the max function, output as follows for students whose highest score for a semester exceeds 90 points. Example Output: Annie's highest grade this semester is 95.

```
for (i in 1:nrow(students)) {

  grades <- c(students$Grade1[i], students$Grade2[i], students$Grade3[i], students$Grade4[i])

  highest_grade <- grades[1]
  for (grade in grades) {
```

```
    if (grade > highest_grade) {  
      highest_grade <- grade  
    }  
  }  
  
  if (highest_grade > 90) {  
    print(paste(students$Name[i], "'s highest grade this semester is", highest_grade))  
  }  
}  
  
## [1] "Annie 's highest grade this semester is 100"  
## [1] "Hanna 's highest grade this semester is 100"
```