Table 7.45D Mean Age at First Substance Use: Among Past Year Initiates Aged 12 to 49; Standard Errors of Averages, 2002-2020

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
ILLICIT DRUGS	nc	nr	nr	nr	nr	nr	nr												
Marijuana	0.19	0.14	0.21	0.22	0.21	0.34	0.35	0.13	0.48	0.18	0.32	0.26	0.28	0.27	0.34	0.28	0.27	0.38	0.51
Cocaine	0.33	0.36	0.45	0.38	0.51	0.52	0.44	0.42	0.96	0.47	0.32	0.36	0.48	0.38	0.38	0.38	0.37	0.61	0.91
Crack	1.35	1.01	1.35	1.28	1.47	2.54	1.68	0.66	2.45	1.05	1.03	2.70	2.78	0.57	1.54	1.51	1.84	2.00	3.73
Heroin	2.51	1.36	2.53	1.55	1.51	1.23	3.18	1.40	1.05	1.22	1.18	1.26	1.28	0.97	1.11	1.34	1.85	2.53	*
Hallucinogens	nc	0.32	0.25	0.37	0.46	0.50	0.67												
LSD	0.49	0.32	0.71	0.78	0.72	0.29	0.31	0.21	0.40	0.27	0.44	0.37	0.33	0.44	0.38	0.40	0.48	0.47	0.88
PCP	0.27	0.49	1.17	0.40	0.61	0.51	0.63	0.80	0.57	0.94	0.50	0.61	0.55	0.68	1.93	1.11	*	0.70	*
Ecstasy	nc	0.37	0.39	0.48	0.51	0.50	1.37												
Inhalants	nc	0.58	0.57	0.69	0.50	0.57	0.95												
Methamphetamine	nc	1.36	1.00	1.10	1.32	1.39	3.92												
Misuse of Prescription																			
Psychotherapeutics ¹	nc	nr	nr	nr	nr	nr	nr												
Pain Relievers ¹	nc	0.58	0.52	0.66	0.64	0.75	1.55												
Stimulants	nc	0.52	0.67	0.63	0.59	0.64	1.30												
Tranquilizers or Sedatives	nc	nr	nr	nr	nr	nr	nr												
Tranquilizers	nc	0.63	0.55	0.63	0.62	0.77	1.13												
Sedatives	nc	1.50	1.14	1.79	1.37	2.47	2.63												
Benzodiazepines																			
Opioids	nc	nr	nr	nr	nr	nr	nr												
Illicit Drugs Other Than																			
Marijuana ²	nc	nr	nr	nr	nr	nr	nr												
CIGARETTES	0.30	0.26	0.19	0.33	0.21	0.12	0.26	0.25	0.21	0.13	0.23	0.22	0.23	0.17	0.16	0.25	0.18	0.26	0.42
Daily Cigarette Use	0.72	0.62	0.40	0.71	0.53	0.45	0.78	0.69	0.34	0.44	0.51	0.65	0.49	0.50	0.74	0.55	0.48	1.07	0.84
SMOKELESS TOBACCO	nc	0.50	0.61	0.48	0.60	0.54	0.79												
CIGARS	0.50	0.45	0.37	0.45	0.42	0.42	0.38	0.44	0.37	0.33	0.38	0.62	0.33	0.33	0.36	0.42	0.33	0.39	0.89
ALCOHOL	0.19	0.10	0.12	0.11	0.11	0.13	0.16	0.13	0.16	0.13	0.19	0.14	0.08	0.13	0.12	0.12	0.09	0.11	0.15

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

^{*=} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions).

NOTE: Methodological limitations preclude the estimation of past year initiates for aggregate substance use categories having at least one prescription psychotherapeutic, including categories for overall illicit drugs, misuse of prescription psychotherapeutics, tranquilizers or sedatives, opioids, and illicit drugs other than marijuana (see Chapter 3 of the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions).

¹ Prescription psychotherapeutic subtypes were revised in 2016; one effect was the comparability of codeine products between 2015 and later years.

² Illicit Drugs Other Than Marijuana excludes respondents who used only marijuana but includes those who used marijuana in addition to other illicit drugs.