Table 3.13C Selected Risk and Protective Factors for Substance Use Disorders and Mental Illness: Among People Aged 12 to 17; by Demographic and Geographic Characteristics, Standard Errors of Numbers in Thousands, 2019 and 2020

	RISK FACTORS				PROTECTIVE FACTORS				
	Youth Argued with at Least One Parent 10 or More Times in the Past Year		Youth Was Approached by Someone Selling Drugs in the Past Month		Youth Had Someone to Talk to about Problems <sup>1</sup>		Youth Attended 25 or More Religious Services in the Past Year		
Demographic/Geographic Characteristic	2019	2020	2019	2020	2019	2020	2019	2020	
TOTAL	98	194	72	105	56	133	122	257	
AGE GROUP									
12-13	55	120	29	48	150	300	76	151	
14-15	62	126	48	78	152	297	85	176	
16-17	65	127	55	64	164	299	79	167	
GENDER									
Male	60	122	54	78	39	93	84	180	
Female	71	144	50	76	40	92	80	171	
HISPANIC ORIGIN AND RACE									
Not Hispanic or Latino	83	162	62	72	44	95	106	217	
White	83	167	55	68	237	389	128	235	
Black or African American	28	53	32	33	111	219	44	109	
AIAN	6	*	4	*	26	*	*	*	
NHOPI	*	*	*	*	*	*	*	*	
Asian	26	54	20	16	82	182	42	88	
Two or More Races	19	34	13	9	43	61	19	18	
Hispanic or Latino	52	105	42	79	34	87	57	129	
COUNTY TYPE									
Large Metro	94	186	63	91	262	489	121	254	
Small Metro	60	117	43	64	195	323	93	172	
Nonmetro	31	49	24	32	132	186	57	113	
Urbanized	20	37	17	22	95	117	38	74	
Less Urbanized	21	33	15	24	86	143	39	78	
Completely Rural	14	*	5	*	45	61	23	*	

<sup>\* =</sup> low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Respondents with unknown Risk or Protective Factors data were excluded from the respective analyses.

<sup>&</sup>lt;sup>1</sup> Respondents were able to select multiple categories of people that they would talk to about a serious problem, including the choice of "nobody." Respondents who reported both a person that they would talk to and that they had nobody to talk to were excluded from the analysis.

Table 3.13D Selected Risk and Protective Factors for Substance Use Disorders and Mental Illness: Among People Aged 12 to 17; by Demographic and Geographic Characteristics, Standard Errors of Percentages, 2019 and 2020

	RISK FACTORS				PROTECTIVE FACTORS				
	Youth Argued with at Least One Parent 10 or More Times in the Past Year		Youth Was Approached by Someone Selling Drugs in the Past Month		Youth Had Someone to Talk to about Problems <sup>1</sup>		Youth Attended 25 or More Religious Services in the Past Year		
Demographic/Geographic Characteristic	2019	2020	2019	2020	2019	2020	2019	2020	
TOTAL	0.40	0.79	0.29	0.42	0.23	0.55	0.51	1.06	
AGE GROUP									
12-13	0.65	1.32	0.35	0.56	0.35	0.96	0.84	1.63	
14-15	0.68	1.33	0.53	0.85	0.42	0.73	0.84	1.68	
16-17	0.68	1.45	0.63	0.80	0.38	1.06	0.79	1.77	
GENDER									
Male	0.49	0.98	0.43	0.62	0.32	0.75	0.69	1.46	
Female	0.59	1.19	0.41	0.62	0.34	0.78	0.68	1.43	
HISPANIC ORIGIN AND RACE									
Not Hispanic or Latino	0.46	0.88	0.33	0.39	0.24	0.52	0.59	1.19	
White	0.56	1.11	0.39	0.50	0.28	0.66	0.74	1.42	
Black or African American	0.78	1.59	0.86	1.06	0.71	1.20	1.12	2.97	
AIAN	3.00	*	2.07	*	3.10	*	*	*	
NHOPI	*	*	*	*	*	*	*	*	
Asian	1.77	2.66	1.37	0.84	0.94	1.21	2.44	4.21	
Two or More Races	1.85	4.66	1.37	1.54	1.16	2.65	1.87	2.89	
Hispanic or Latino	0.87	1.70	0.68	1.25	0.57	1.44	0.97	2.09	
COUNTY TYPE									
Large Metro	0.60	1.10	0.42	0.61	0.32	0.75	0.70	1.51	
Small Metro	0.69	1.36	0.50	0.77	0.41	0.99	0.92	1.87	
Nonmetro	0.79	1.50	0.58	0.97	0.47	0.93	1.21	2.78	
Urbanized	1.12	2.58	0.93	1.58	0.61	1.69	1.86	4.29	
Less Urbanized	1.14	1.94	0.85	1.45	0.78	1.12	1.69	3.91	
Completely Rural	2.47	*	1.20	*	1.22	1.52	3.76	*	

<sup>\* =</sup> low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Respondents with unknown Risk or Protective Factors data were excluded from the respective analyses.

<sup>&</sup>lt;sup>1</sup> Respondents were able to select multiple categories of people that they would talk to about a serious problem, including the choice of "nobody." Respondents who reported both a person that they would talk to and that they had nobody to talk to were excluded from the analysis.