Table 8.38A Major Depressive Episode (MDE) and MDE with Severe Impairment in Past Year: Among People Aged 18 or Older; Receipt of Treatment for Depression in Past Year: Among People Aged 18 or Older with MDE and among People Aged 18 or Older with MDE with Severe Impairment in Past Year; by Gender and Age Group, Numbers in Thousands, 2019 and 2020

Gender/Age Group	Had MDE (2019)	Had MDE (2020)	Had MDE with Severe Impairment ¹ (2019)	Had MDE with Severe Impairment ¹ (2020)	Received Treatment for Depression in Past Year among People with MDE ² (2019)	Received Treatment for Depression in Past Year among People with MDE ² (2020)	Received Treatment for Depression in Past Year among People with MDE with Severe Impairment ^{1,2} (2019)	Received Treatment for Depression in Past Year among People with MDE with Severe Impairment ^{1,2} (2020)
TOTAL	19,357	20,976	13,112	14,845	12,824	13,832	9,467	10,537
18-25	5,045	5,610	3,407	3,978	2,565	3,230	1,921	2,536
26 or Older	14,312	15,366	9,705	10,867	10,258	10,602	7,546	8,001
26-49	8,914	9,168	6,101	6,510	6,135	5,937	4,532	4,475
50 or Older	5,399	6,199	3,604	4,357	4,124	4,665	3,014	3,526
MALE	7,125	7,440	4,646	5,387	4,041	4,193	2,868	3,363
18-25	1,841	1,829	1,236	1,300	793	854	599	682
26 or Older	5,284	5,612	3,409	4,087	3,248	3,339	2,268	2,681
26-49	3,271	3,649	2,169	2,584	1,927	1,978	1,369	1,541
50 or Older	2,013	1,962	1,240	1,503	1,321	*	899	*
FEMALE	12,233	13,536	8,466	9,458	8,783	9,639	6,599	7,174
18-25	3,204	3,781	2,170	2,678	1,772	2,377	1,321	1,854
26 or Older	9,028	9,755	6,296	6,780	7,011	7,262	5,278	5,320
26-49	5,643	5,518	3,932	3,926	4,208	3,958	3,163	2,934
50 or Older	3,386	4,236	2,364	2,854	2,802	3,304	2,115	*

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health:

Methodological Summary and Definitions for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

¹ Respondents with unknown impairment data were excluded.

² Respondents with unknown past year treatment for depression data were excluded.

Table 8.38B Major Depressive Episode (MDE) and MDE with Severe Impairment in Past Year: Among People Aged 18 or Older; Receipt of Treatment for Depression in Past Year: Among People Aged 18 or Older with MDE and among People Aged 18 or Older with MDE with Severe Impairment in Past Year; by Gender and Age Group, Percentages, 2019 and 2020

Gender/Age Group	Had MDE (2019)	Had MDE (2020)	Had MDE with Severe Impairment ¹ (2019)	Had MDE with Severe Impairment ¹ (2020)	Received Treatment for Depression in Past Year among People with MDE ² (2019)	Received Treatment for Depression in Past Year among People with MDE ² (2020)	Received Treatment for Depression in Past Year among People with MDE with Severe Impairment ^{1,2} (2019)	Received Treatment for Depression in Past Year among People with MDE with Severe Impairment ^{1,2} (2020)
TOTAL	7.8	8.4	5.3	6.0	66.3	66.0	72.2	71.0
18-25	15.2	17.0	10.3	12.1	50.9	57.6	56.4	63.8
26 or Older	6.7	7.1	4.5	5.0	71.8	69.0	77.8	73.6
26-49	8.9	9.1	6.1	6.5	68.9	64.8	74.4	68.7
50 or Older	4.7	5.4	3.2	3.8	76.5	75.3	83.6	80.9
MALE	6.0	6.2	3.9	4.5	56.8	56.4	61.8	62.5
18-25	11.1	11.1	7.4	7.9	43.1	46.8	48.5	52.7
26 or Older	5.1	5.4	3.3	3.9	61.6	59.5	66.7	65.6
26-49	6.6	7.3	4.4	5.2	59.2	54.2	63.3	59.6
50 or Older	3.8	3.6	2.3	2.8	65.6	*	72.5	*
FEMALE	9.6	10.5	6.6	7.3	71.9	71.2	78.0	75.8
18-25	19.4	22.9	13.1	16.3	55.3	62.9	60.9	69.2
26 or Older	8.1	8.7	5.7	6.0	77.7	74.5	83.8	78.5
26-49	11.1	10.8	7.8	7.7	74.6	71.7	80.4	74.7
50 or Older	5.6	6.9	3.9	4.7	83.0	78.0	89.5	*

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health:

Methodological Summary and Definitions for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

¹ Respondents with unknown impairment data were excluded.

² Respondents with unknown past year treatment for depression data were excluded.