Table 8.57A Had Serious Thoughts of Suicide in Past Year: Among People Aged 18 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Numbers in Thousands, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)	Aged 26-49 (2019)	Aged 26-49 (2020)	Aged 50+ (2019)	Aged 50+ (2020)
TOTAL	11,951	12,197	3,926	3,771	8,025	8,427	5,331	5,317	2,693	3,110
GEOGRAPHIC REGION	<b>y</b>	,	- ,-	- ,	- 7	.,	-	- ,	,	-, -
Northeast	2,073	1,747	654	557	1,419	1,190	745	817	674	373
Midwest	2,742	2,942	967	882	1,776	2,060	1,128	1,352	648	708
South	4,059	4,083	1,339	1,278	2,720	2,805	1,942	1,670	778	1,135
West	3,076	3,425	966	1,053	2,110	2,372	1,516	1,479	594	893
COUNTY TYPE										
Large Metro	6,412	6,079	2,175	1,954	4,237	4,125	2,908	2,707	1,329	1,418
Small Metro	3,899	4,322	1,255	1,248	2,644	3,074	1,712	1,951	932	1,123
Nonmetro	1,640	1,797	496	569	1,143	1,228	712	660	432	568
Urbanized	719	911	246	268	473	643	294	336	180	307
Less Urbanized	785	788	194	269	590	519	358	310	232	209
Completely Rural	136	97	56	*	80	66	60	14	20	52
POVERTY LEVEL <sup>1</sup>										
Less Than 100%	2,290	2,311	925	857	1,365	1,454	969	957	396	497
100-199%	2,735	3,110	890	995	1,845	2,115	1,004	1,144	841	971
200% or More	6,809	6,720	1,995	1,862	4,814	4,858	3,358	3,216	1,456	1,641
EDUCATION LEVEL										
< High School	1,286	1,303	509	384	777	919	518	494	259	425
High School Graduate	2,952	3,245	1,239	1,199	1,712	2,046	1,093	1,056	619	990
Some College/Associate's										
Degree	4,793	4,713	1,787	1,763	3,006	2,950	2,080	1,905	926	1,045
College Graduate	2,920	2,936	391	424	2,529	2,512	1,639	1,863	890	649
HEALTH INSURANCE <sup>2</sup>										
Private	6,834	6,603	2,349	2,301	4,486	4,301	2,882	2,764	1,604	1,537
Medicaid/CHIP	2,865	2,799	894	801	1,972	1,998	1,344	1,464	627	534
Other <sup>3</sup>	2,234	2,583	294	325	1,940	2,257	636	733	1,305	1,524
No Coverage	1,608	1,575	582	522	1,026	1,053	902	812	124	241

<sup>\* =</sup> low precision: -- = not available: da = does not apply: nc = not comparable due to methodological changes: nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020.

Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Respondents with unknown suicide information were excluded.

<sup>&</sup>lt;sup>1</sup> Respondents aged 18 to 22 who were living in a college dormitory were excluded.

<sup>&</sup>lt;sup>2</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>&</sup>lt;sup>3</sup> Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Table 8.57B Had Serious Thoughts of Suicide in Past Year: Among People Aged 18 or Older; by Age Group and Geographic and Socioeconomic Characteristics, Percentages, 2019 and 2020

Geographic/Socioeconomic Characteristic	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)	Aged 26-49 (2019)	Aged 26-49 (2020)	Aged 50+ (2019)	Aged 50+ (2020)
TOTAL	4.8	4.9	11.8	11.3	3.7	3.9	5.3	5.3	2.4	2.7
GEOGRAPHIC REGION										,
Northeast	4.8	4.0	11.5	9.9	3.8	3.1	4.4	4.8	3.3	1.8
Midwest	5.3	5.7	13.6	12.5	4.0	4.6	5.6	6.7	2.7	2.9
South	4.3	4.3	10.6	10.1	3.3	3.4	5.1	4.3	1.8	2.6
West	5.2	5.7	12.1	13.3	4.1	4.6	6.0	5.8	2.3	3.4
COUNTY TYPE										
Large Metro	4.6	4.5	11.7	11.1	3.5	3.5	4.9	4.7	2.2	2.3
Small Metro	5.2	5.6	12.0	11.4	4.1	4.7	5.9	6.3	2.6	3.2
Nonmetro	4.8	4.8	11.5	11.9	3.8	3.8	5.8	5.1	2.4	2.9
Urbanized	5.0	6.1	12.2	12.0	3.8	5.0	5.5	6.4	2.6	4.1
Less Urbanized	4.9	4.4	10.1	12.3	4.2	3.3	6.4	4.7	2.7	2.3
Completely Rural	3.5	2.1	15.1	*	2.2	1.5	4.5	1.3	0.9	1.6
POVERTY LEVEL <sup>1</sup>										
Less Than 100%	7.2	6.2	11.9	10.8	5.7	5.0	7.3	6.4	3.7	3.5
100-199%	5.8	6.8	11.6	12.9	4.7	5.6	5.3	6.3	4.0	4.9
200% or More	4.0	4.0	11.9	10.8	3.2	3.2	4.9	4.7	1.8	2.0
EDUCATION LEVEL										
< High School	4.3	4.6	11.1	10.3	3.1	3.7	4.9	4.9	1.8	2.9
High School Graduate	4.9	4.7	11.9	11.3	3.4	3.5	5.3	4.4	2.1	2.9
Some College/Associate's										
Degree	6.3	6.2	13.3	12.7	4.8	4.7	6.8	6.5	2.9	3.2
College Graduate	3.6	3.8	8.0	8.6	3.3	3.5	4.3	4.9	2.3	1.9
HEALTH INSURANCE <sup>2</sup>										
Private	4.1	4.2	11.6	11.8	3.1	3.1	4.3	4.3	2.1	2.1
Medicaid/CHIP	8.1	7.1	12.6	10.7	7.0	6.3	8.4	8.4	5.1	3.7
Other <sup>3</sup>	3.3	3.5	11.7	13.6	2.9	3.1	8.8	9.2	2.2	2.4
No Coverage	6.3	5.8	12.2	9.7	4.9	4.8	6.1	5.0	2.1	4.3

<sup>\* =</sup> low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020.

Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Respondents with unknown suicide information were excluded.

<sup>&</sup>lt;sup>1</sup> Respondents aged 18 to 22 who were living in a college dormitory were excluded.

<sup>&</sup>lt;sup>2</sup> Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

<sup>&</sup>lt;sup>3</sup> Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.