Table 10.32C Had Serious Thoughts of Suicide in Past Year: Among People Aged 18 or Older; by Geographic and Socioeconomic Characteristics, Standard Errors of Numbers in Thousands, 2008-2020

| Geographic/Socioeconomic Characteristic | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|------|------|------|------|------|------|------|------|------|------|------|------|------|
| TOTAL | 299 | 299 | 311 | 299 | 305 | 340 | 281 | 280 | 267 | 289 | 282 | 300 | 499 |
| GEOGRAPHIC REGION | | | | | | | | | | | | | |
| Northeast | 123 | 115 | 121 | 125 | 127 | 128 | 108 | 128 | 114 | 112 | 103 | 135 | 180 |
| Midwest | 127 | 129 | 130 | 127 | 115 | 130 | 129 | 118 | 115 | 119 | 120 | 131 | 235 |
| South | 178 | 178 | 193 | 178 | 196 | 225 | 161 | 172 | 154 | 172 | 171 | 171 | 311 |
| West | 156 | 167 | 156 | 165 | 147 | 185 | 148 | 142 | 155 | 164 | 155 | 166 | 247 |
| COUNTY TYPE ¹ | | | | | | | | | | | | | |
| Large Metro | 237 | 239 | 242 | 234 | 242 | 277 | 220 | 231 | 215 | 234 | 229 | 237 | 363 |
| Small Metro | 174 | 159 | 161 | 178 | 172 | 178 | 150 | 154 | 159 | 166 | 157 | 181 | 286 |
| Nonmetro | 87 | 100 | 128 | 105 | 119 | 126 | 112 | 103 | 90 | 121 | 107 | 108 | 200 |
| Urbanized | 42 | 67 | 76 | 59 | 73 | 99 | 72 | 72 | 64 | 80 | 85 | 76 | 162 |
| Less Urbanized | 72 | 69 | 97 | 80 | 74 | 83 | 85 | 69 | 59 | 91 | 68 | 81 | 111 |
| Completely Rural | 26 | 30 | 42 | 39 | 52 | 26 | 30 | 31 | 29 | 38 | 31 | 25 | 34 |
| POVERTY LEVEL ² | | | | | | | | | | | | | |
| Less Than 100% | 108 | 134 | 123 | 159 | 133 | 149 | 122 | 125 | 135 | 125 | 130 | 121 | 208 |
| 100-199% | 137 | 132 | 141 | 135 | 175 | 144 | 144 | 140 | 142 | 132 | 125 | 155 | 271 |
| 200% or More | 232 | 238 | 242 | 226 | 223 | 288 | 203 | 212 | 201 | 235 | 227 | 219 | 362 |
| EDUCATION LEVEL | | | | | | | | | | | | | |
| < High School | nc | 122 | 114 | 95 | 96 | 101 | 187 |
| High School Graduate | nc | 138 | 125 | 142 | 145 | 148 | 274 |
| Some College/Associate's Degree | nc | 164 | 163 | 181 | 165 | 189 | 283 |
| College Graduate | nc | 141 | 139 | 160 | 150 | 164 | 218 |
| HEALTH INSURANCE ³ | | | | | | | | | | | | | |
| Private | 230 | 231 | 233 | 221 | 210 | 262 | 195 | 217 | 202 | 229 | 219 | 224 | 356 |
| Medicaid/CHIP | 112 | 125 | 117 | 125 | 122 | 111 | 125 | 141 | 137 | 126 | 128 | 153 | 232 |
| Other ⁴ | 157 | 144 | 184 | 152 | 185 | 200 | 152 | 163 | 148 | 171 | 138 | 165 | 275 |
| No Coverage | 112 | 123 | 120 | 139 | 156 | 143 | 99 | 90 | 100 | 91 | 103 | 96 | 159 |

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019, and Quarters 1 and 4, 2020.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions).

NOTE: Respondents with unknown suicide information were excluded.

Because of the use of the 2013 Rural-Urban Continuum Codes in the creation of the county type variables, these 2008 to 2014 estimates may differ from estimates published prior to the 2015 NSDUH.

² Respondents aged 18 to 22 who were living in a college dormitory were excluded.

³ Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

⁴ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.

Table 10.32D Had Serious Thoughts of Suicide in Past Year: Among People Aged 18 or Older; by Geographic and Socioeconomic Characteristics, Standard Errors of Percentages, 2008-2020

| Geographic/Socioeconomic Characteristic | 2008 | 2009 | 2010 | 2011 | 2012 | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | 2020 |
|---|------|------|------|------|------|------|------|------|------|------|------|------|------|
| TOTAL | 0.13 | 0.13 | 0.14 | 0.13 | 0.13 | 0.14 | 0.12 | 0.12 | 0.11 | 0.12 | 0.11 | 0.12 | 0.20 |
| GEOGRAPHIC REGION | | | | | | | | | | | | | |
| Northeast | 0.30 | 0.27 | 0.29 | 0.30 | 0.30 | 0.30 | 0.25 | 0.30 | 0.26 | 0.26 | 0.24 | 0.31 | 0.41 |
| Midwest | 0.26 | 0.26 | 0.26 | 0.25 | 0.23 | 0.26 | 0.25 | 0.23 | 0.22 | 0.23 | 0.23 | 0.25 | 0.45 |
| South | 0.22 | 0.22 | 0.23 | 0.21 | 0.23 | 0.26 | 0.18 | 0.19 | 0.17 | 0.19 | 0.18 | 0.18 | 0.33 |
| West | 0.30 | 0.32 | 0.30 | 0.31 | 0.27 | 0.34 | 0.26 | 0.25 | 0.27 | 0.28 | 0.26 | 0.28 | 0.41 |
| COUNTY TYPE ¹ | | | | | | | | | | | | | |
| Large Metro | 0.19 | 0.19 | 0.19 | 0.18 | 0.18 | 0.21 | 0.16 | 0.16 | 0.15 | 0.16 | 0.16 | 0.16 | 0.26 |
| Small Metro | 0.25 | 0.22 | 0.23 | 0.23 | 0.23 | 0.24 | 0.21 | 0.20 | 0.21 | 0.21 | 0.20 | 0.22 | 0.36 |
| Nonmetro | 0.24 | 0.29 | 0.38 | 0.30 | 0.32 | 0.34 | 0.29 | 0.30 | 0.23 | 0.32 | 0.28 | 0.29 | 0.51 |
| Urbanized | 0.30 | 0.46 | 0.53 | 0.45 | 0.48 | 0.62 | 0.45 | 0.51 | 0.42 | 0.47 | 0.47 | 0.50 | 1.03 |
| Less Urbanized | 0.40 | 0.39 | 0.58 | 0.45 | 0.42 | 0.48 | 0.45 | 0.41 | 0.30 | 0.47 | 0.39 | 0.44 | 0.61 |
| Completely Rural | 0.73 | 0.84 | 1.21 | 1.08 | 1.34 | 0.68 | 0.77 | 0.68 | 0.75 | 1.11 | 0.85 | 0.49 | 0.70 |
| POVERTY LEVEL ² | | | | | | | | | | | | | |
| Less Than 100% | 0.41 | 0.47 | 0.41 | 0.46 | 0.35 | 0.41 | 0.33 | 0.33 | 0.38 | 0.35 | 0.36 | 0.37 | 0.55 |
| 100-199% | 0.33 | 0.30 | 0.29 | 0.28 | 0.37 | 0.31 | 0.30 | 0.28 | 0.28 | 0.27 | 0.25 | 0.32 | 0.56 |
| 200% or More | 0.14 | 0.15 | 0.16 | 0.15 | 0.14 | 0.18 | 0.13 | 0.13 | 0.12 | 0.14 | 0.13 | 0.13 | 0.21 |
| EDUCATION LEVEL | | | | | | | | | | | | | |
| < High School | nc | 0.35 | 0.35 | 0.31 | 0.31 | 0.33 | 0.65 |
| High School Graduate | nc | 0.22 | 0.20 | 0.24 | 0.23 | 0.24 | 0.40 |
| Some College/Associate's Degree | nc | 0.22 | 0.21 | 0.23 | 0.21 | 0.24 | 0.37 |
| College Graduate | nc | 0.19 | 0.18 | 0.19 | 0.18 | 0.20 | 0.28 |
| HEALTH INSURANCE ³ | | | | | | | | | | | | | |
| Private | 0.15 | 0.15 | 0.15 | 0.14 | 0.14 | 0.16 | 0.12 | 0.13 | 0.12 | 0.14 | 0.13 | 0.13 | 0.22 |
| Medicaid/CHIP | 0.59 | 0.63 | 0.54 | 0.52 | 0.52 | 0.45 | 0.41 | 0.40 | 0.38 | 0.34 | 0.35 | 0.41 | 0.57 |
| Other ⁴ | 0.30 | 0.27 | 0.34 | 0.27 | 0.31 | 0.33 | 0.24 | 0.25 | 0.22 | 0.25 | 0.20 | 0.24 | 0.37 |
| No Coverage | 0.32 | 0.33 | 0.31 | 0.35 | 0.39 | 0.38 | 0.33 | 0.34 | 0.41 | 0.39 | 0.42 | 0.36 | 0.58 |

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Respondents with unknown suicide information were excluded.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2008-2019, and Quarters 1 and 4, 2020.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Some 2008 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions).

Because of the use of the 2013 Rural-Urban Continuum Codes in the creation of the county type variables, these 2008 to 2014 estimates may differ from estimates published prior to the 2015 NSDUH.

² Respondents aged 18 to 22 who were living in a college dormitory were excluded.

³ Respondents could indicate multiple types of health insurance; thus, these response categories are not mutually exclusive.

⁴ Other Health Insurance is defined as having Medicare, military-related health care, or any other type of health insurance.