Table 8.24A (REVISED) Types and Locations of Mental Health Services Received in Past Year: Among People Aged 18 or Older; by Past Year Level of Mental Illness, Numbers in Thousands, 2019 and 2020

Type and Location of Mental Health Services	Any Mental Illness (2019)	Any Mental Illness (2020)	Serious Mental Illness (2019)	Serious Mental Illness (2020)	Any Mental Illness Excluding Serious Mental Illness (2019)	Any Mental Illness Excluding Serious Mental Illness (2020)	No Mental Illness ¹ (2019)	No Mental Illness ¹ (2020)
MENTAL HEALTH SERVICES ²	22,953	24,303	8,575	9,133	14,378	15,169	17,201	18,118
Outpatient	13,855	14,918	5,773	6,295	8,082	8,623	6,706	7,329
Outpatient Mental Health Clinic or Center Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or	3,990	4,659	2,033	2,358	1,958	2,301	1,172	1,369
Counselor - Not Part of a Clinic	8,240	9,253	3,398	3,789	4,843	5,465	4,188	4,535
Doctor's Office - Not Part of a Clinic Outpatient Medical Clinic Partial Day Hospital or Day	2,162 857	1,988 1,188	830 422	866 630	1,332 434	1,123 558	974 324	842 423
Treatment Program School or University	321	243	183	209	138	34	25	123
Setting/Clinic/Center ³	103	82	35	20	68	63	48	*
Some Other Place ⁴	312	269	180	200	132	69	223	213
Inpatient	1,674	1,545	974	929	699	616	697	719
Prescription Medication NONE OF THESE SERVICES	18,854 28,330	20,479 28,311	7,402 4,515	7,937 5,021	11,452 23,815	12,542 23,290	13,775 180,287	14,113 180,077

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020.

Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. Therefore, some estimates in this table could differ from corresponding estimates in other tables that did not undergo sample size adjustments. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Mental Health Services for adults includes inpatient treatment/counseling, outpatient treatment/counseling, or use of prescription medication for problems with emotions, nerves, or mental health. Respondents with unknown mental health service information were excluded. Questions on virtual services were added in Quarter 4 of 2020. Estimates in this table do not include data from these questions. Data on virtual services are included at the end of this section. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Mental İllness alığns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

¹ Respondents with no mental illness may still have received mental health treatment for a past disorder that is in partial or full remission or treatment for a disorder that was not included in the definition of mental illness or for other reasons.

² Respondents could indicate multiple service types; thus, these response categories are not mutually exclusive. Questions on virtual services were added in Quarter 4 of 2020. Estimates in this table do not include data from these questions. Data on virtual services are included at the end of this section. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

³ Respondents were permitted to specify other locations for receiving outpatient mental health services. This location was the most commonly reported other location for receiving outpatient mental health services.

⁴ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Services were excluded.

Table 8.24B (REVISED) Types and Locations of Mental Health Services Received in Past Year: Among People Aged 18 or Older; by Past Year Level of Mental Illness, Percentages, 2019 and 2020

Type and Location of Mental Health Services	Any Mental Illness (2019)	Any Mental Illness (2020)	Serious Mental Illness (2019)	Serious Mental Illness (2020)	Any Mental Illness Excluding Serious Mental Illness (2019)	Any Mental Illness Excluding Serious Mental Illness (2020)	No Mental Illness ¹ (2019)	No Mental Illness ¹ (2020)
MENTAL HEALTH SERVICES ²	44.8	46.2	65.5	64.5	37.6	39.4	8.7	9.1
Outpatient	27.1	28.5	44.2	45.1	21.2	22.5	3.4	3.7
Outpatient Mental Health Clinic or Center Office of a Private Therapist, Psychologist, Psychiatrist, Social Worker, or	7.8	8.9	15.6	16.9	5.1	6.0	0.6	0.7
Counselor - Not Part of a Clinic Doctor's Office - Not Part of a	16.1	17.7	26.1	27.2	12.7	14.3	2.1	2.3
Clinic	4.2	3.8	6.4	6.2	3.5	2.9	0.5	0.4
Outpatient Medical Clinic Partial Day Hospital or Day	1.7	2.3	3.2	4.5	1.1	1.5	0.2	0.2
Treatment Program School or University	0.6	0.5	1.4	1.5	0.4	0.1	0.0	0.1
Setting/Clinic/Center ³	0.2	0.2	0.3	0.1	0.2	0.2	0.0	*
Some Other Place ⁴	0.6	0.5	1.4	1.4	0.3	0.2	0.1	0.1
Inpatient	3.3	2.9	7.4	6.5	1.8	1.6	0.4	0.4
Prescription Medication	36.7	38.8	56.5	55.9	29.9	32.5	7.0	7.1
NONE OF THESE SERVICES	55.2	53.8	34.5	35.5	62.4	60.6	91.3	90.9

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020.

Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. Therefore, some estimates in this table could differ from corresponding estimates in other tables that did not undergo sample size adjustments. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Mental Health Services for adults includes inpatient treatment/counseling, outpatient treatment/counseling, or use of prescription medication for problems with emotions, nerves, or mental health. Respondents with unknown mental health service information were excluded. Questions on virtual services were added in Quarter 4 of 2020. Estimates in this table do not include data from these questions. Data on virtual services are included at the end of this section. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Mental İllness aligns with DSM-IV criteria and is defined as having a diagnosable mental, behavioral, or emotional disorder, other than a developmental or substance use disorder. Estimates of serious mental illness (SMI) are a subset of estimates of any mental illness (AMI) because SMI is limited to people with AMI that resulted in serious functional impairment. These mental illness estimates are based on a predictive model and are not direct measures of diagnostic status.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

¹ Respondents with no mental illness may still have received mental health treatment for a past disorder that is in partial or full remission or treatment for a disorder that was not included in the definition of mental illness or for other reasons.

² Respondents could indicate multiple service types; thus, these response categories are not mutually exclusive. Questions on virtual services were added in Quarter 4 of 2020. Estimates in this table do not include data from these questions. Data on virtual services are included at the end of this section. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

³ Respondents were permitted to specify other locations for receiving outpatient mental health services. This location was the most commonly reported other location for receiving outpatient mental health services.

⁴ Respondents with unknown or invalid responses to the other-specify question on Some Other Place Received Outpatient Mental Health Services were excluded.