Table 1.22A Any Use and Misuse of Prescription Psychotherapeutics in Past Year: Among People Aged 12 or Older; Misuse of Prescription Psychotherapeutics in Past Year: Among Past Year Any Users Aged 12 or Older; by Age Group, Numbers in Thousands, 2019 and 2020

Prescription Drug/Age Group	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users (2019)	Misuse in Past Year among Past Year Any Users (2020)
PSYCHOTHERAPEUTICS	108,037	101,602	16,304	16,073	16,304	16,073
12-17	5,612	4,723	1,062	688	1,062	688
18 or Older	102,425	96,879	15,242	15,385	15,242	15,385
18-25	12,187	10,529	3,868	3,193	3,868	3,193
26 or Older	90,238	86,350	11,374	12,192	11,374	12,192
PAIN RELIEVERS	82,559	75,198	9,724	9,254	9,724	9,254
12-17	3,980	3,079	567	396	567	396
18 or Older	78,579	72,118	9,158	8,858	9,158	8,858
18-25	8,357	6,858	1,754	1,361	1,754	1,361
26 or Older	70,222	65,261	7,404	7,498	7,404	7,498
STIMULANTS	18,290	18,332	4,929	5,092	4,929	5,092
12-17	1,837	1,593	430	288	430	288
18 or Older	16,453	16,739	4,499	4,804	4,499	4,804
18-25	4,320	3,773	1,951	1,616	1,951	1,616
26 or Older	12,133	12,966	2,548	3,188	2,548	3,188
TRANQUILIZERS OR SEDATIVES	44,051	43,001	5,895	6,205	5,895	6,205
12-17	1,370	936	436	226	436	226
18 or Older	42,682	42,065	5,459	5,979	5,459	5,979
18-25	3,995	3,805	1,409	1,237	1,409	1,237
26 or Older	38,687	38,261	4,050	4,742	4,050	4,742
TRANQUILIZERS	35,601	35,018	5,220	5,460	5,220	5,460
12-17	1,010	588	392	196	392	196
18 or Older	34,591	34,430	4,829	5,264	4,829	5,264
18-25	3,419	3,338	1,313	1,108	1,313	1,108
26 or Older	31,172	31,092	3,516	4,156	3,516	4,156
SEDATIVES	15,636	15,364	1,098	1,240	1,098	1,240
12-17	545	429	82	35	82	35
18 or Older	15,091	14,935	1,016	1,205	1,016	1,205
18-25	1,051	1,066	175	251	175	251
26 or Older	14,040	13,869	841	954	841	954
BENZODIAZEPINES	29,408	27,026	4,841	4,779	4,841	4,779
12-17	749	418	381	157	381	*
18 or Older	28,659	26,607	4,460	4,622	4,460	4,622
18-25	2,983	2,861	1,266	1,097	1,266	1,097
26 or Older	25,676	23,747	3,194	3,526	3,194	3,526

<sup>\* =</sup> low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health:

Methodological Summary and Definitions for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

Table 1.22B Any Use and Misuse of Prescription Psychotherapeutics in Past Year: Among People Aged 12 or Older; Misuse of Prescription Psychotherapeutics in Past Year: Among Past Year Any Users Aged 12 or Older; by Age Group, Percentages, 2019 and 2020

Prescription Drug/Age Group	Any Use in Past Year (2019)	Any Use in Past Year (2020)	Misuse in Past Year among Total Population (2019)	Misuse in Past Year among Total Population (2020)	Misuse in Past Year among Past Year Any Users (2019)	Misuse in Past Year among Past Year Any Users (2020)
PSYCHOTHERAPEUTICS	39.3	36.7	5.9	5.8	15.1	15.8
12-17	22.5	18.9	4.3	2.8	18.9	14.6
18 or Older	40.9	38.5	6.1	6.1	14.9	15.9
18-25	36.1	31.4	11.5	9.5	31.7	30.3
26 or Older	41.7	39.5	5.3	5.6	12.6	14.1
PAIN RELIEVERS	30.0	27.2	3.5	3.3	11.8	12.3
12-17	16.0	12.3	2.3	1.6	14.2	12.9
18 or Older	31.4	28.6	3.7	3.5	11.7	12.3
18-25	24.8	20.5	5.2	4.1	21.0	19.8
26 or Older	32.4	29.9	3.4	3.4	10.5	11.5
STIMULANTS	6.6	6.6	1.8	1.8	26.9	27.8
12-17	7.4	6.4	1.7	1.2	23.4	18.1
18 or Older	6.6	6.6	1.8	1.9	27.3	28.7
18-25	12.8	11.3	5.8	4.8	45.2	42.8
26 or Older	5.6	5.9	1.2	1.5	21.0	24.6
TRANQUILIZERS OR SEDATIVES	16.0	15.5	2.1	2.2	13.4	14.4
12-17	5.5	3.7	1.8	0.9	31.8	24.1
18 or Older	17.1	16.7	2.2	2.4	12.8	14.2
18-25	11.8	11.4	4.2	3.7	35.3	32.5
26 or Older	17.9	17.5	1.9	2.2	10.5	12.4
TRANQUILIZERS	12.9	12.6	1.9	2.0	14.7	15.6
12-17	4.1	2.4	1.6	0.8	38.8	33.3
18 or Older	13.8	13.7	1.9	2.1	14.0	15.3
18-25	10.1	10.0	3.9	3.3	38.4	33.2
26 or Older	14.4	14.2	1.6	1.9	11.3	13.4
SEDATIVES	5.7	5.5	0.4	0.4	7.0	8.1
12-17	2.2	1.7	0.3	0.1	15.1	8.1
18 or Older	6.0	5.9	0.4	0.5	6.7	8.1
18-25	3.1	3.2	0.5	0.7	16.6	23.5
26 or Older	6.5	6.3	0.4	0.4	6.0	6.9
BENZODIAZEPINES	10.7	9.8	1.8	1.7	16.5	17.7
12-17	3.0	1.7	1.5	0.6	50.9	*
18 or Older	11.4	10.6	1.8	1.8	15.6	17.4
18-25	8.8	8.5	3.8	3.3	42.4	38.3
26 or Older	11.9	10.9	1.5	1.6	12.4	14.8

<sup>\* =</sup> low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health:

Methodological Summary and Definitions for details.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.