Table 7.16C Alcohol Use in Lifetime, Past Year, and Past Month: Among People Aged 12 to 20; by Gender, Standard Errors of Numbers in Thousands, 2002-2020

Gender/Alcohol Use	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
TOTAL																			
Lifetime	286	286	272	286	282	312	288	302	291	332	324	282	276	256	284	300	262	317	429
Past Year	250	253	246	268	259	293	264	273	266	307	308	255	261	237	261	280	246	295	398
Past Month	188	196	191	208	208	231	201	212	204	239	244	194	211	186	199	210	189	219	300
Binge Alcohol Use <sup>1</sup>	nc	152	146	159	141	162	232												
Heavy Alcohol Use	nc	77	64	69	63	56	93												
MALE																			
Lifetime	179	183	177	184	185	192	184	203	197	188	200	177	181	175	171	176	170	181	276
Past Year	158	166	162	171	170	179	165	187	180	173	187	162	169	154	155	164	159	167	251
Past Month	124	134	130	137	137	142	131	151	141	131	154	124	138	117	117	130	115	127	192
Binge Alcohol Use <sup>1</sup>	107	114	115	123	117	122	109	128	118	107	121	102	114	98	91	98	88	96	143
Heavy Alcohol Use	59	59	70	67	73	72	66	76	66	59	66	53	56	52	45	47	43	38	76
FEMALE																			
Lifetime	180	170	170	180	169	190	179	175	170	213	189	174	184	167	189	194	172	199	285
Past Year	160	159	155	171	156	181	169	157	156	199	179	159	172	160	175	182	161	190	273
Past Month	118	123	119	128	125	141	124	124	118	159	140	122	140	123	137	129	128	139	198
Binge Alcohol Use <sup>1</sup>	nc	97	101	101	93	103	158												
Heavy Alcohol Use	nc	46	46	44	38	38	52												

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

<sup>\* =</sup> low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health:

Methodological Summary and Definitions for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions).

<sup>&</sup>lt;sup>1</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.

Table 7.16D Alcohol Use in Lifetime, Past Year, and Past Month: Among People Aged 12 to 20; by Gender, Standard Errors of Percentages, 2002-2020

Gender/Alcohol Use	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
TOTAL																			
Lifetime	0.40	0.43	0.41	0.41	0.42	0.45	0.43	0.45	0.44	0.49	0.48	0.47	0.48	0.48	0.52	0.54	0.48	0.53	0.92
Past Year	0.41	0.44	0.41	0.43	0.43	0.47	0.43	0.44	0.44	0.50	0.51	0.46	0.49	0.48	0.51	0.54	0.48	0.53	0.90
Past Month	0.39	0.41	0.39	0.41	0.42	0.46	0.40	0.43	0.41	0.47	0.48	0.40	0.46	0.42	0.45	0.47	0.42	0.45	0.72
Binge Alcohol Use <sup>1</sup>	nc	0.36	0.35	0.37	0.33	0.36	0.59												
Heavy Alcohol Use	nc	0.20	0.17	0.18	0.16	0.14	0.25												
MALE																			
Lifetime	0.55	0.53	0.56	0.55	0.54	0.59	0.55	0.57	0.60	0.61	0.61	0.58	0.66	0.64	0.65	0.64	0.63	0.64	1.20
Past Year	0.56	0.57	0.56	0.56	0.57	0.60	0.54	0.58	0.60	0.60	0.62	0.58	0.66	0.62	0.63	0.64	0.62	0.64	1.14
Past Month	0.52	0.53	0.55	0.53	0.54	0.57	0.53	0.58	0.55	0.54	0.61	0.51	0.60	0.53	0.53	0.57	0.50	0.54	0.91
Binge Alcohol Use <sup>1</sup>	0.48	0.49	0.52	0.51	0.49	0.51	0.47	0.53	0.50	0.47	0.52	0.45	0.53	0.46	0.43	0.46	0.42	0.43	0.71
Heavy Alcohol Use	0.29	0.29	0.34	0.32	0.35	0.34	0.31	0.36	0.32	0.29	0.31	0.25	0.28	0.26	0.22	0.24	0.22	0.19	0.39
FEMALE																			
Lifetime	0.51	0.59	0.55	0.56	0.58	0.61	0.58	0.60	0.59	0.67	0.63	0.62	0.65	0.64	0.70	0.73	0.65	0.72	1.29
Past Year	0.53	0.62	0.55	0.58	0.58	0.64	0.61	0.59	0.58	0.68	0.64	0.61	0.65	0.65	0.69	0.74	0.66	0.72	1.28
Past Month	0.50	0.57	0.51	0.54	0.56	0.59	0.53	0.55	0.53	0.65	0.59	0.54	0.61	0.57	0.63	0.59	0.59	0.61	0.99
Binge Alcohol Use <sup>1</sup>	nc	0.48	0.49	0.49	0.45	0.49	0.82												
Heavy Alcohol Use	nc	0.24	0.25	0.23	0.20	0.20	0.29												

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

<sup>\* =</sup> low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health:

Methodological Summary and Definitions for details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions).

<sup>&</sup>lt;sup>1</sup> Estimates of binge alcohol use include use by those who were heavy alcohol users.