Table 1.102A Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime: Among People Aged 12 or Older; by Age Group, Numbers in Thousands, 2019 and 2020

Drug or Method of Administration	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
HALLUCINOGENS	44,087	43,949	615	563	43,472	43,386	5,597	5,269	37,876	38,117
PCP	5,516	6,141	41	7	5,475	6,134	249	167	5,226	5,967
LSD	27,528	28,123	392	317	27,137	27,806	3,368	3,284	23,769	24,521
Peyote	5,670	5,040	49	24	5,621	5,016	208	111	5,413	4,905
Mescaline	7,571	6,913	21	3	7,550	6,911	176	98	7,373	6,813
Psilocybin/Mushrooms	25,254	26,111	285	204	24,968	25,908	2,990	2,936	21,979	22,971
Ecstasy	20,097	20,478	225	183	19,872	20,295	3,500	2,946	16,372	17,348
Ketamine	3,833	3,955	32	38	3,801	3,917	539	600	3,262	3,317
DMT, AMT, or 5-MeO-DIPT/Foxy	3,345	3,192	51	42	3,294	3,150	697	552	2,597	2,598
Salvia divinorum	5,188	5,106	52	28	5,136	5,079	699	423	4,437	4,655
INHALANTS	25,090	26,749	2,202	2,327	22,889	24,422	3,099	2,929	19,790	21,493
Amyl Nitrite, Poppers, Locker Room	•									
Odorizers, or Rush	7,122	8,168	126	155	6,995	8,013	541	549	6,454	7,464
Correction Fluid, Degreaser, or Cleaning Fluid	1,298	1,685	163	217	1,135	1,467	184	63	950	1,405
Gasoline or Lighter Fluid	2,814	3,132	436	472	2,378	2,660	321	334	2,057	2,326
Glue, Shoe Polish, or Toluene	2,723	3,489	525	692	2,198	2,797	297	330	1,901	2,467
Halothane, Ether, or Other Anesthetics	609	1,230	29	12	581	1,218	54	81	527	1,137
Lacquer Thinner or Other Paint Solvents	1,452	1,557	130	185	1,322	1,372	170	125	1,152	1,247
Lighter Gases	806	733	149	135	656	598	99	68	557	529
Nitrous Oxide or Whippits	12,640	12,891	139	125	12,500	12,766	1,138	1,031	11,363	11,735
Felt-Tip Pens or Felt-Tip/Magic Markers	6,595	7,640	1,599	1,875	4,996	5,765	1,412	1,336	3,584	4,428
Spray Paints	1,800	1,864	407	439	1,394	1,425	268	282	1,126	1,143
Computer Cleaner/Air Duster	2,885	2,685	188	116	2,697	2,569	551	285	2,146	2,283
Other Aerosol Sprays	1,368	1,372	131	112	1,237	1,260	193	134	1,044	1,126
NEEDLE USE ¹	4,592	5,417	19	2	4,572	5,415	280	187	4,292	5,228
Heroin Needle Use ¹	2,710	2,896	5	*	2,705	2,896	172	133	2,533	2,763
Cocaine Needle Use ¹	2,380	2,698	2	*	2,378	2,698	89	21	2,289	2,677
Methamphetamine Needle Use ¹	2,141	2,724	*	*	2,141	2,724	142	75	1,999	2,649
HEROIN	5,696	6,252	21	*	5,675	6,251	389	226	5,287	6,024
Smoke Heroin ¹	2,567	3,028	13	*	2,554	3,027	193	154	2,361	2,873
Sniff or Snort Heroin ¹	3,746	4,378	4	*	3,741	4,378	285	170	3,457	4,208

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

¹ These estimates may be inconsistent with estimates presented in other detailed tables because NSDUH data are not edited for consistency across discrete sections of the interview, except with regard to skip logic. Definitions: Measures and terms are defined in Appendix A.

Table 1.102B Specific Hallucinogen, Inhalant, Needle, and Heroin Use in Lifetime: Among People Aged 12 or Older; by Age Group, Percentages, 2019 and 2020

Drug or Method of Administration	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
HALLUCINOGENS	16.0	15.9	2.5	2.3	17.4	17.2	16.6	15.7	17.5	17.5
PCP	2.0	2.2	0.2	0.0	2.2	2.4	0.7	0.5	2.4	2.7
LSD	10.0	10.2	1.6	1.3	10.8	11.0	10.0	9.8	11.0	11.2
Peyote	2.1	1.8	0.2	0.1	2.2	2.0	0.6	0.3	2.5	2.2
Mescaline	2.8	2.5	0.1	0.0	3.0	2.7	0.5	0.3	3.4	3.1
Psilocybin/Mushrooms	9.2	9.4	1.1	0.8	10.0	10.3	8.9	8.8	10.1	10.5
Ecstasy	7.3	7.4	0.9	0.7	7.9	8.1	10.4	8.8	7.6	7.9
Ketamine	1.4	1.4	0.1	0.2	1.5	1.6	1.6	1.8	1.5	1.5
DMT, AMT, or 5-MeO-DIPT/Foxy	1.2	1.2	0.2	0.2	1.3	1.3	2.1	1.6	1.2	1.2
Salvia divinorum	1.9	1.8	0.2	0.1	2.1	2.0	2.1	1.3	2.0	2.1
INHALANTS	9.1	9.7	8.8	9.3	9.1	9.7	9.2	8.7	9.1	9.8
Amyl Nitrite, Poppers, Locker Room										
Odorizers, or Rush	2.6	2.9	0.5	0.6	2.8	3.2	1.6	1.6	3.0	3.4
Correction Fluid, Degreaser, or Cleaning Fluid	0.5	0.6	0.7	0.9	0.5	0.6	0.5	0.2	0.4	0.6
Gasoline or Lighter Fluid	1.0	1.1	1.7	1.9	1.0	1.1	1.0	1.0	0.9	1.1
Glue, Shoe Polish, or Toluene	1.0	1.3	2.1	2.8	0.9	1.1	0.9	1.0	0.9	1.1
Halothane, Ether, or Other Anesthetics	0.2	0.4	0.1	0.0	0.2	0.5	0.2	0.2	0.2	0.5
Lacquer Thinner or Other Paint Solvents	0.5	0.6	0.5	0.7	0.5	0.5	0.5	0.4	0.5	0.6
Lighter Gases	0.3	0.3	0.6	0.5	0.3	0.2	0.3	0.2	0.3	0.2
Nitrous Oxide or Whippits	4.6	4.7	0.6	0.5	5.0	5.1	3.4	3.1	5.2	5.4
Felt-Tip Pens or Felt-Tip/Magic Markers	2.4	2.8	6.4	7.5	2.0	2.3	4.2	4.0	1.7	2.0
Spray Paints	0.7	0.7	1.6	1.8	0.6	0.6	0.8	0.8	0.5	0.5
Computer Cleaner/Air Duster	1.0	1.0	0.8	0.5	1.1	1.0	1.6	0.9	1.0	1.0
Other Aerosol Sprays	0.5	0.5	0.5	0.4	0.5	0.5	0.6	0.4	0.5	0.5
NEEDLE USE ¹	1.7	2.0	0.1	0.0	1.8	2.1	0.8	0.6	2.0	2.4
Heroin Needle Use ¹	1.0	1.0	0.0	*	1.1	1.1	0.5	0.4	1.2	1.3
Cocaine Needle Use ¹	0.9	1.0	0.0	*	1.0	1.1	0.3	0.1	1.1	1.2
Methamphetamine Needle Use ¹	0.8	1.0	*	*	0.9	1.1	0.4	0.2	0.9	1.2
HEROIN	2.1	2.3	0.1	*	2.3	2.5	1.2	0.7	2.4	2.8
Smoke Heroin ¹	0.9	1.1	0.1	*	1.0	1.2	0.6	0.5	1.1	1.3
Sniff or Snort Heroin ¹	1.4	1.6	0.0	*	1.5	1.7	0.8	0.5	1.6	1.9

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

¹ These estimates may be inconsistent with estimates presented in other detailed tables because NSDUH data are not edited for consistency across discrete sections of the interview, except with regard to skip logic. Definitions: Measures and terms are defined in Appendix A.