Substance Use Disorder Status in Past Year: Among People Aged 12 to 17; by Past Year Major Depressive Episode (MDE) and MDE with Severe Table 9.9C Impairment Status and Demographic Characteristics, Standard Errors of Numbers in Thousands, 2019 and 2020

	SUBSTANCE USE DISORDER STATUS													
Major Depressive	Substance Use Disorder													
Episode Status/ Demographic	No Substance Use Disorder		Illicit Drugs		Marijuana		Opioids		Alcohol		Both Illicit Drugs and Alcohol		Illicit Drugs or Alcohol	
Characteristic	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)
$TOTAL^1$	nc	123	nc	104	nc	95	nc	24	nc	86	nc	56	nc	123
MDE	nc	212	nc	70	nc	58	nc	16	nc	57	nc	40	nc	81
Age Group														
12-13	nc	126	nc	15	nc	5	nc	5	nc	5	nc	1	nc	16
14-15	nc	111	nc	51	nc	43	nc	6	nc	36	nc	32	nc	53
16-17	nc	117	nc	42	nc	34	nc	14	nc	43	nc	24	nc	55
Gender														
Male	nc	110	nc	21	nc	19	nc	*	nc	18	nc	12	nc	25
Female	nc	166	nc	66	nc	55	nc	13	nc	54	nc	38	nc	76
MDE with Severe Impairment	nc	168	nc	50	nc	46	nc	15	nc	50	nc	35	nc	61
Age Group														
12-13	nc	101	nc	13	nc	4	nc	4	nc	5	nc	1	nc	14
14-15	nc	87	nc	41	nc	40	nc	5	nc	36	nc	32	nc	44
16-17	nc	95	nc	24	nc	20	nc	13	nc	35	nc	14	nc	39
Gender														
Male	nc	92	nc	17	nc	17	nc	*	nc	15	nc	11	nc	19
Female	nc	138	nc	46	nc	42	nc	11	nc	47	nc	33	nc	57
NO MDE	nc	532	nc	83	nc	80	nc	18	nc	68	nc	41	nc	99
Age Group														
12-13	nc	273	nc	10	nc	4	nc	1	nc	*	nc	1	nc	41
14-15	nc	278	nc	38	nc	38	nc	10	nc	36	nc	21	nc	49
16-17	nc	270	nc	73	nc	71	nc	15	nc	40	nc	36	nc	74
Gender														
Male	nc	357	nc	63	nc	60	nc	18	nc	60	nc	39	nc	78
Female	nc	331	nc	52	nc	52	nc	1	nc	28	nc	11	nc	58

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

^{*=} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

¹ The Total row includes respondents with unknown past year MDE information.

Substance Use Disorder Status in Past Year: Among People Aged 12 to 17; by Past Year Major Depressive Episode (MDE) and MDE with Severe Table 9.9D Impairment Status and Demographic Characteristics, Standard Errors of Percentages, 2019 and 2020

Major Depressive Episode Status/ Demographic Characteristic	SUBSTANCE USE DISORDER STATUS													
	Substance Use Disorder													
	No Substance Use Disorder		Illicit Drugs		Marijuana		Opioids		Alcohol		Both Illicit Drugs and Alcohol		Illicit Drugs or Alcohol	
	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)	(2019)	(2020)
$TOTAL^1$	nc	0.49	nc	0.42	nc	0.38	nc	0.10	nc	0.34	nc	0.22	nc	0.49
MDE	nc	1.79	nc	1.57	nc	1.31	nc	0.38	nc	1.30	nc	0.91	nc	1.79
Age Group														
12-13	nc	1.83	nc	1.72	nc	0.51	nc	0.51	nc	0.62	nc	0.16	nc	1.83
14-15	nc	3.11	nc	3.01	nc	2.61	nc	0.42	nc	2.19	nc	1.99	nc	3.11
16-17	nc	2.92	nc	2.32	nc	1.90	nc	0.80	nc	2.39	nc	1.36	nc	2.92
Gender														
Male	nc	2.21	nc	1.84	nc	1.74	nc	*	nc	1.61	nc	1.02	nc	2.21
Female	nc	2.31	nc	2.05	nc	1.72	nc	0.42	nc	1.70	nc	1.21	nc	2.31
MDE with Severe Impairment	nc	1.94	nc	1.61	nc	1.48	nc	0.50	nc	1.61	nc	1.14	nc	1.94
Age Group														
12-13	nc	2.22	nc	2.04	nc	0.70	nc	0.70	nc	0.87	nc	0.22	nc	2.22
14-15	nc	3.50	nc	3.33	nc	3.31	nc	0.47	nc	2.96	nc	2.71	nc	3.50
16-17	nc	3.11	nc	2.04	nc	1.71	nc	1.08	nc	2.74	nc	1.17	nc	3.11
Gender														
Male	nc	2.69	nc	2.34	nc	2.33	nc	*	nc	2.06	nc	1.55	nc	2.69
Female	nc	2.37	nc	1.94	nc	1.82	nc	0.51	nc	2.02	nc	1.43	nc	2.37
NO MDE	nc	0.48	nc	0.40	nc	0.39	nc	0.09	nc	0.33	nc	0.20	nc	0.48
Age Group														
12-13	nc	0.57	nc	0.14	nc	0.06	nc	0.02	nc	*	nc	0.02	nc	0.57
14-15	nc	0.69	nc	0.54	nc	0.54	nc	0.15	nc	0.52	nc	0.30	nc	0.69
16-17	nc	1.17	nc	1.15	nc	1.12	nc	0.24	nc	0.65	nc	0.58	nc	1.17
Gender														
Male	nc	0.67	nc	0.55	nc	0.52	nc	0.16	nc	0.53	nc	0.34	nc	0.67
Female	nc	0.65	nc	0.59	nc	0.59	nc	0.01	nc	0.31	nc	0.13	nc	0.65

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

^{*=} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Substance use disorder (SUD) estimates in 2020 are based on criteria from the Diagnostic and Statistical Manual of Mental Disorders, 5th edition (DSM-5). SUD and related estimates are not comparable between 2020 and prior years of NSDUH, because prior years' estimates were based on DSM-IV criteria. The 2020 estimates reflect additional methodological changes for the 2020 NSDUH. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details on these changes.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

¹ The Total row includes respondents with unknown past year MDE information.