

**Table 6.6A Source Where Friend or Relative Obtained Prescription Pain Relievers: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Pain Relievers from a Friend or Relative for Free in Past Year; by Age Group, Numbers in Thousands, 2019 and 2020**

Source Where Friend or Relative Obtained Pain Relievers	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	2,735	2,477	*	*	2,662	2,366	474	361	2,188	2,005
Prescription from One Doctor	2,664	2,390	*	*	2,598	2,285	457	358	2,141	1,927
Prescriptions from More Than One Doctor	54	26	*	*	49	24	13	*	36	22
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	18	*	*	*	15	*	4	*	11	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	306	285	*	*	289	276	60	84	228	*
From Friend or Relative for Free	193	192	*	*	191	191	43	*	148	*
Bought from Friend or Relative	70	77	*	*	67	72	11	*	56	*
Took from Friend or Relative without Asking	43	15	*	*	30	13	6	*	24	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	83	69	*	*	62	60	16	11	47	49
<b>SOME OTHER WAY<sup>1</sup></b>	53	49	*	*	43	49	12	8	32	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.6B Source Where Friend or Relative Obtained Prescription Pain Relievers: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Pain Relievers from a Friend or Relative for Free in Past Year; by Age Group, Percentages, 2019 and 2020**

Source Where Friend or Relative Obtained Pain Relievers	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	86.1	86.0	*	*	87.1	86.0	84.4	77.9	87.7	87.7
Prescription from One Doctor	83.8	83.0	*	*	85.0	83.1	81.3	77.3	85.8	84.3
Prescriptions from More Than One Doctor	1.7	0.9	*	*	1.6	0.9	2.4	*	1.4	0.9
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.6	*	*	*	0.5	*	0.7	*	0.4	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	9.6	9.9	*	*	9.4	10.0	10.7	18.0	9.2	*
From Friend or Relative for Free	6.1	6.7	*	*	6.3	7.0	7.7	*	5.9	*
Bought from Friend or Relative	2.2	2.7	*	*	2.2	2.6	2.0	*	2.3	*
Took from Friend or Relative without Asking	1.3	0.5	*	*	1.0	0.5	1.1	*	1.0	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	2.6	2.4	*	*	2.0	2.2	2.8	2.4	1.9	2.1
<b>SOME OTHER WAY<sup>1</sup></b>	1.7	1.7	*	*	1.4	1.8	2.1	1.7	1.3	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.8A Source Where Friend or Relative Obtained Prescription Stimulants: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Stimulants from a Friend or Relative for Free in Past Year; by Age Group, Numbers in Thousands, 2019 and 2020**

Source Where Friend or Relative Obtained Stimulants	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	1,981	<i>2,103</i>	*	*	1,843	<i>2,029</i>	736	<i>624</i>	1,107	<i>1,405</i>
Prescription from One Doctor	1,931	<i>2,059</i>	*	*	1,812	<i>1,996</i>	710	<i>613</i>	1,101	<i>1,383</i>
Prescriptions from More Than One Doctor	50	<i>41</i>	*	*	32	<i>29</i>	26	<i>8</i>	6	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	<i>3</i>	*	*	*	<i>3</i>	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	303	<i>254</i>	*	*	278	<i>236</i>	103	<i>90</i>	175	<i>145</i>
From Friend or Relative for Free	201	<i>127</i>	*	*	184	<i>112</i>	61	<i>34</i>	123	*
Bought from Friend or Relative	101	<i>125</i>	*	*	94	<i>120</i>	42	<i>56</i>	52	<i>64</i>
Took from Friend or Relative without Asking	0	<i>3</i>	*	*	*	<i>3</i>	*	<i>0</i>	*	<i>3</i>
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	40	<i>66</i>	*	*	27	<i>41</i>	5	*	22	<i>33</i>
<b>SOME OTHER WAY<sup>1</sup></b>	26	<i>9</i>	*	*	25	<i>1</i>	8	<i>1</i>	17	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.8B Source Where Friend or Relative Obtained Prescription Stimulants: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Stimulants from a Friend or Relative for Free in Past Year; by Age Group, Percentages, 2019 and 2020**

Source Where Friend or Relative Obtained Stimulants	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	84.3	86.5	*	*	84.8	87.9	86.4	86.2	83.7	88.7
Prescription from One Doctor	82.2	84.6	*	*	83.3	86.5	83.3	84.6	83.3	87.4
Prescriptions from More Than One Doctor	2.1	1.7	*	*	1.5	1.3	3.0	1.1	0.4	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	0.1	*	*	*	0.1	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	12.9	10.5	*	*	12.8	10.2	12.1	12.5	13.3	9.2
From Friend or Relative for Free	8.6	5.2	*	*	8.5	4.9	7.1	4.7	9.3	*
Bought from Friend or Relative	4.3	5.1	*	*	4.3	5.2	5.0	7.7	3.9	4.1
Took from Friend or Relative without Asking	0.0	0.1	*	*	*	0.1	*	0.1	*	0.2
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	1.7	2.7	*	*	1.2	1.8	0.6	*	1.7	2.1
<b>SOME OTHER WAY<sup>1</sup></b>	1.1	0.4	*	*	1.2	0.1	1.0	0.2	1.3	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.10A Source Where Friend or Relative Obtained Prescription Tranquilizers: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Tranquilizers from a Friend or Relative for Free in Past Year; by Age Group, Numbers in Thousands, 2019 and 2020**

Source Where Friend or Relative Obtained Tranquilizers	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	1,762	2,245	*	*	1,703	2,224	385	*	1,318	1,823
Prescription from One Doctor	1,707	2,222	*	*	1,652	2,201	375	*	1,277	1,803
Prescriptions from More Than One Doctor	40	23	*	*	37	23	5	*	32	20
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	15	*	*	*	15	*	5	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	240	218	*	*	202	206	100	*	103	128
From Friend or Relative for Free	98	167	*	*	85	165	45	*	40	120
Bought from Friend or Relative	134	29	*	*	112	22	50	14	62	9
Took from Friend or Relative without Asking	9	21	*	*	5	19	*	*	*	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	93	97	*	*	72	69	48	*	23	21
<b>SOME OTHER WAY<sup>1</sup></b>	32	41	*	*	29	41	1	0	28	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.10B Source Where Friend or Relative Obtained Prescription Tranquilizers: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Tranquilizers from a Friend or Relative for Free in Past Year; by Age Group, Percentages, 2019 and 2020**

Source Where Friend or Relative Obtained Tranquilizers	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	82.8	86.3	*	*	84.9	87.5	72.1	*	89.6	90.6
Prescription from One Doctor	80.2	85.4	*	*	82.3	86.6	70.2	*	86.7	89.6
Prescriptions from More Than One Doctor	1.9	0.9	*	*	1.9	0.9	0.9	*	2.2	1.0
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	0.7	*	*	*	0.7	*	1.0	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	11.3	8.4	*	*	10.1	8.1	18.7	*	7.0	6.4
From Friend or Relative for Free	4.6	6.4	*	*	4.2	6.5	8.4	*	2.7	5.9
Bought from Friend or Relative	6.3	1.1	*	*	5.6	0.9	9.3	2.6	4.2	0.4
Took from Friend or Relative without Asking	0.4	0.8	*	*	0.2	0.7	*	*	*	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	4.4	3.7	*	*	3.6	2.7	9.0	*	1.6	1.0
<b>SOME OTHER WAY<sup>1</sup></b>	1.5	1.6	*	*	1.4	1.6	0.2	0.0	1.9	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.12A Source Where Friend or Relative Obtained Prescription Sedatives: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Sedatives from a Friend or Relative for Free in Past Year; by Age Group, Numbers in Thousands, 2019 and 2020**

Source Where Friend or Relative Obtained Sedatives	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	430	*	*	*	*	*	*	*	*	*
Prescription from One Doctor	422	*	*	*	413	*	*	*	*	*
Prescriptions from More Than One Doctor	7	*	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	49	*	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*	*	*
Bought from Friend or Relative	11	*	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	5	*	*	*	*	*	*	*	*	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0	*	*	*	0	*	*	*	*	*
<b>SOME OTHER WAY<sup>1</sup></b>	*	*	*	*	*	*	*	*	*	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.

**Table 6.12B Source Where Friend or Relative Obtained Prescription Sedatives: Among Past Year Misusers Aged 12 or Older Who Obtained Most Recently Misused Prescription Sedatives from a Friend or Relative for Free in Past Year; by Age Group, Percentages, 2019 and 2020**

Source Where Friend or Relative Obtained Sedatives	Aged 12+ (2019)	Aged 12+ (2020)	Aged 12-17 (2019)	Aged 12-17 (2020)	Aged 18+ (2019)	Aged 18+ (2020)	Aged 18-25 (2019)	Aged 18-25 (2020)	Aged 26+ (2019)	Aged 26+ (2020)
<b>GOT THROUGH PRESCRIPTION(S) OR STOLE FROM A HEALTH CARE PROVIDER</b>	89.7	*	*	*	*	*	*	*	*	*
Prescription from One Doctor	88.2	*	*	*	89.5	*	*	*	*	*
Prescriptions from More Than One Doctor	1.5	*	*	*	*	*	*	*	*	*
Stole from Doctor's Office, Clinic, Hospital, or Pharmacy	*	*	*	*	*	*	*	*	*	*
<b>GIVEN BY, BOUGHT FROM, OR TOOK FROM A FRIEND OR RELATIVE</b>	10.3	*	*	*	*	*	*	*	*	*
From Friend or Relative for Free	*	*	*	*	*	*	*	*	*	*
Bought from Friend or Relative	2.4	*	*	*	*	*	*	*	*	*
Took from Friend or Relative without Asking	1.1	*	*	*	*	*	*	*	*	*
<b>BOUGHT FROM DRUG DEALER OR OTHER STRANGER</b>	0.0	*	*	*	0.0	*	*	*	*	*
<b>SOME OTHER WAY<sup>1</sup></b>	*	*	*	*	*	*	*	*	*	*

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Respondents were asked to choose only one source. Respondents with unknown data on Source for Most Recent Misuse and respondents with unknown or invalid responses to the corresponding other-specify questions were excluded from the analysis.

<sup>1</sup> Some Other Way includes write-in responses not already listed in this table or responses with insufficient information that could allow them to be placed in another category.

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2019 and Quarters 1 and 4, 2020.