Table 8.36A Major Depressive Episode in Past Year: Among People Aged 18 or Older; by Gender and Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
TOTAL	19,357	20,976	7,125	7,440	12,233	13,536
18-25	5,045	5,610	1,841	1,829	3,204	3,781
18-20	2,050	2,002	736	665	1,314	1,338
21-25	2,995	3,607	1,105	1,164	1,890	2,443
26 or Older	14,312	15,366	5,284	5,612	9,028	9,755
26-49	8,914	9,168	3,271	3,649	5,643	5,518
26-29	2,088	1,805	852	717	1,236	1,088
30-34	2,259	2,126	845	832	1,414	1,293
35-39	1,705	1,838	487	752	1,218	1,086
40-44	1,455	1,412	532	601	922	811
45-49	1,407	1,987	555	747	852	1,240
50 or Older	5,399	6,199	2,013	1,962	3,386	4,236
50-54	1,364	1,338	562	458	801	880
55-59	1,270	1,654	452	431	818	1,223
60-64	1,056	1,139	307	344	749	795
65 or Older	1,709	2,067	693	729	1,017	1,338

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Table 8.36B Major Depressive Episode in Past Year: Among People Aged 18 or Older; by Gender and Detailed Age Category, Percentages, 2019 and 2020

Age Category	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
TOTAL	7.8	8.4	6.0	6.2	9.6	10.5
18-25	15.2	17.0	11.1	11.1	19.4	22.9
18-20	15.8	16.3	11.1	10.5	20.8	22.5
21-25	14.8	17.4	11.1	11.4	18.5	23.2
26 or Older	6.7	7.1	5.1	5.4	8.1	8.7
26-49	8.9	9.1	6.6	7.3	11.1	10.8
26-29	11.7	10.3	9.5	8.2	14.0	12.3
30-34	10.2	9.4	7.8	7.3	12.6	11.5
35-39	8.0	8.4	4.7	7.2	11.1	9.5
40-44	7.6	7.3	5.7	6.3	9.5	8.3
45-49	7.0	10.2	5.5	7.6	8.5	12.8
50 or Older	4.7	5.4	3.8	3.6	5.6	6.9
50-54	6.7	6.8	5.6	5.2	7.8	8.2
55-59	6.1	7.7	4.5	4.1	7.5	11.2
60-64	5.2	5.5	3.2	3.3	6.9	7.8
65 or Older	3.3	3.8	3.0	3.0	3.5	4.5

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Table 8.37A Major Depressive Episode with Severe Impairment in Past Year: Among People Aged 18 or Older; by Gender and Detailed Age Category, Numbers in Thousands, 2019 and 2020

Age Category	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
TOTAL	13,112	14,845	4,646	5,387	8,466	9,458
18-25	3,407	3,978	1,236	1,300	2,170	2,678
18-20	1,341	1,335	482	469	859	866
21-25	2,065	2,644	754	832	1,311	1,812
26 or Older	9,705	10,867	3,409	4,087	6,296	6,780
26-49	6,101	6,510	2,169	2,584	3,932	3,926
26-29	1,427	1,243	558	447	869	796
30-34	1,468	1,530	511	547	957	983
35-39	1,192	1,374	331	626	860	747
40-44	1,039	920	401	398	638	522
45-49	976	1,443	367	565	609	878
50 or Older	3,604	4,357	1,240	1,503	2,364	2,854
50-54	887	1,051	341	373	546	678
55-59	886	1,091	314	368	572	723
60-64	737	830	174	277	563	553
65 or Older	1,094	1,385	411	485	684	900

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.

Table 8.37B Major Depressive Episode with Severe Impairment in Past Year: Among People Aged 18 or Older; by Gender and Detailed Age Category, Percentages, 2019 and 2020

Age Category	Total (2019)	Total (2020)	Male (2019)	Male (2020)	Female (2019)	Female (2020)
TOTAL	5.3	6.0	3.9	4.5	6.6	7.3
18-25	10.3	12.1	7.4	7.9	13.1	16.3
18-20	10.3	10.9	7.2	7.4	13.6	14.6
21-25	10.2	12.8	7.6	8.2	12.8	17.2
26 or Older	4.5	5.0	3.3	3.9	5.7	6.0
26-49	6.1	6.5	4.4	5.2	7.8	7.7
26-29	8.0	7.1	6.2	5.1	9.9	9.0
30-34	6.7	6.8	4.7	4.8	8.5	8.7
35-39	5.6	6.3	3.2	6.0	7.9	6.6
40-44	5.5	4.8	4.3	4.2	6.6	5.4
45-49	4.9	7.4	3.7	5.8	6.0	9.1
50 or Older	3.2	3.8	2.3	2.8	3.9	4.7
50-54	4.4	5.4	3.4	4.2	5.3	6.3
55-59	4.2	5.1	3.1	3.5	5.3	6.6
60-64	3.6	4.0	1.8	2.6	5.2	5.5
65 or Older	2.1	2.6	1.8	2.0	2.4	3.0

^{* =} low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.

NOTE: Estimates in this table for 2020 exclude a subset of respondents who did not complete the questionnaire. The analysis weights and estimates were adjusted for the reduced sample size. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.

NOTE: Respondents with unknown past year Major Depressive Episode (MDE) data were excluded.

NOTE: Respondents with unknown impairment data were excluded.

Definitions: Measures and terms are defined in Appendix A.