

Table 7.24C Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 12 to 20 and among Past Year Users of Specific Substances Aged 12 to 20; Standard Errors of Numbers in Thousands, 2002-2020

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION																			
Marijuana	50	49	43	53	50	47	51	50	52	52	66	58	62	61	56	62	60	60	<i>107</i>
Cocaine	5	6	3	5	6	6	4	6	2	6	7	2	2	6	4	5	2	4	<i>20</i>
Crack	3	3	2	3	1	3	2	*	2	2	3	*	*	4	*	*	*	4	*
Heroin	2	4	2	2	*	3	3	3	8	6	5	2	7	4	3	3	3	4	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	3	2	2	3	*	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	3	5	10	3	4	2
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	6	7	7	5	6	*
Alcohol	27	28	24	22	22	25	25	23	24	23	18	23	25	27	18	26	21	20	22
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS																			
Marijuana	50	49	43	53	50	47	51	50	52	52	66	58	62	61	56	62	60	60	<i>107</i>
Cocaine	5	6	3	5	6	6	4	6	2	6	7	2	2	6	4	5	2	4	*
Crack	3	*	*	*	1	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	2	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	3	2	2	3	1	<i>0</i>
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	3	5	10	3	4	2
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	*
Alcohol	27	28	24	22	22	25	25	23	24	23	18	23	25	27	18	26	21	20	22

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions* for details.

NOTE: Some 2002 to 2019 standard errors of numbers in thousands may differ from previously published estimates because of a change to the standard error estimation method. See the Introduction to these tables for more details.

NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the *2020 National Survey on Drug Use and Health: Methodological Summary and Definitions*).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.

Table 7.24D Daily or Almost Daily Use of Specific Substances in Past Year: Among All People Aged 12 to 20 and among Past Year Users of Specific Substances Aged 12 to 20; Standard Errors of Percentages, 2002-2020

Substance	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG TOTAL POPULATION																			
Marijuana	0.13	0.13	0.11	0.13	0.13	0.12	0.13	0.13	0.13	0.13	0.16	0.15	0.16	0.16	0.15	0.16	0.15	0.15	0.28
Cocaine	0.01	0.02	0.01	0.01	0.02	0.01	0.01	0.02	0.01	0.01	0.02	0.00	0.01	0.02	0.01	0.01	0.01	0.01	0.05
Crack	0.01	0.01	0.01	0.01	0.00	0.01	0.01	*	0.00	0.01	0.01	*	*	0.01	*	*	*	0.01	*
Heroin	0.00	0.01	0.01	0.01	*	0.01	0.01	0.01	0.02	0.01	0.01	0.00	0.02	0.01	0.01	0.01	0.01	0.01	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.01	0.01	0.01	*	*
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.01	0.01	0.03	0.01	0.01	0.01
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.02	0.02	0.02	0.01	0.02	*
Alcohol	0.07	0.07	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.06	0.05	0.06	0.06	0.07	0.05	0.07	0.05	0.05	0.06
DAILY/ALMOST DAILY USE IN PAST YEAR AMONG PAST YEAR USERS																			
Marijuana	0.57	0.57	0.52	0.64	0.61	0.59	0.63	0.57	0.60	0.59	0.70	0.66	0.74	0.73	0.71	0.76	0.69	0.67	1.53
Cocaine	0.39	0.50	0.28	0.39	0.45	0.45	0.39	0.65	0.29	0.68	0.89	0.31	0.35	0.79	0.60	0.73	0.31	0.79	*
Crack	1.30	*	*	*	0.64	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Heroin	1.49	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Hallucinogens	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.17	0.16	0.18	0.24	0.05	0.03
Inhalants	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	0.29	0.68	1.25	0.32	0.40	0.29
Methamphetamine	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	nc	*	*	*	*	*	*
Alcohol	0.16	0.16	0.13	0.12	0.13	0.14	0.14	0.14	0.15	0.14	0.12	0.15	0.17	0.19	0.13	0.20	0.16	0.15	0.20

\* = low precision; -- = not available; da = does not apply; nc = not comparable due to methodological changes; nr = not reported due to measurement issues.  
NOTE: Estimates in the 2020 column are italicized to indicate caution should be used when comparing estimates between 2020 and prior years because of methodological changes for 2020. Due to these changes, significance testing between 2020 and prior years was not performed. See the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions for details.  
NOTE: Some 2006 to 2010 estimates may differ from previously published estimates due to updates (see Chapter 3 of the 2020 National Survey on Drug Use and Health: Methodological Summary and Definitions).

Definitions: Measures and terms are defined in Appendix A.

Source: SAMHSA, Center for Behavioral Health Statistics and Quality, National Survey on Drug Use and Health, 2002-2019, and Quarters 1 and 4, 2020.