

Distributed Algorithms 2022/2023

(practical exercise)

Leader election

- 1 – Find the expected value for the random variable $X \sim Geo(p)$.
- 2 – Find the variance of the random variable $X \sim Geo(p)$.
- 3 – Let $p \in [0, 1]$ and $n \geq k \geq 1, n, k \in \mathbb{N}$. For what value of the argument a function f takes the maximum value?
 - a. $f(p) = np(1-p)^{n-1}$,
 - b. $f(n) = np(1-p)^{n-1}$,
 - c. $f(k) = \binom{n}{k} p^k (1-p)^{n-k}$.
- 4 – Prove that $(1+x)^r \geq 1+rx$ for $x \geq -1, r \geq 1$.
- 5 – Prove that $(1+x)^r \leq 1+rx$ for $x \geq -1, r \in (0, 1)$.
- 6 – Prove that $1+x \leq e^x$ for $x \in \mathbb{R}$.
- 7 – Prove that $\frac{x}{e^x} < \frac{1.5}{x^2}$ for $x > 0$.
- 8 – Let $f_i(n) = n^{\frac{1}{2^i}} \left(1 - \frac{1}{2^i}\right)^{n-1}$. Prove that functions $f_i(2^{i-1}), f_{i+1}(2^{i-1})$ are decreasing and function $f_{i-1}(2^i)$ is increasing for $i \geq 2$.
- 9 – Present the definition of the [Lambert W](#) functions family and draw its real branches.
- 10 – Use Lambert W function to analytically [determine](#) real solutions to the equation

$$3^x = x^3.$$

What is Lambert's W function called in Mathematica?

- 11 – Have a look at [this paper](#) and show that for $x \geq e$

$$\ln x - \ln \ln x < W_0(x) \leq \ln x - \frac{1}{2} \ln \ln x.$$

- 12 – Completion of the lecture proof. Check that if $K \geq 1, f > 1, u \geq 2$ and

$$3e(K+1)u^{\frac{-1}{2(K+1)}} \geq 1 - \frac{1}{f} \quad \text{then} \quad K \geq \frac{\ln u}{2W_0(\frac{3e}{2} \frac{f}{f-1} \ln u)} - 1.$$

Data stream analysis: approximate counting

13 – For continuous and independent random variables X_1, X_2, \dots, X_n with the same distribution given by the density function $f(x)$ and the cumulative distribution function $F(x)$ show that k -th order statistic $X_{k:n}$ has a distribution described by the density function

$$f_k(x) = \frac{F^{k-1}(x) [1 - F(x)]^{n-k} f(x)}{B(k, n - k + 1)},$$

where $B(\alpha, \beta)$ denotes [beta function](#). Hint: see [this notes](#).

14 – For n independent random variables U_1, U_2, \dots, U_n with the uniform distribution: $U_i \sim \mathcal{U}(0, 1)$, show that k -th order statistic has distribution $Beta(k, n - k + 1)$ and an expected value equal to $k/(n + 1)$.

Hint. In this and in the next exercise use different representations of the beta function: given by the definition and [by factorial](#) for arguments that are natural numbers.

15 – Let $U_{k:n}$ denote k th order statistic for n independent uniformly distributed random variables with distribution $\mathcal{U}(0, 1)$. Show that for the estimator $\hat{n}_k = \frac{k-1}{U_{k:n}}$ and $k \geq 2$ we have $\mathbb{E}(\hat{n}_k) = n$ and that for $k \geq 3$ we have

$$\text{Var}(\hat{n}_k) = \frac{n(n - k + 1)}{k - 2}.$$

16 – (Markov's inequality) Let X denote the random variable that takes only non-negative values. Then for all $a > 0$

$$\mathbb{P}(X \geq a) \leq \frac{\mathbb{E}(X)}{a}.$$

17 – (Chebyshev's inequality) Let X denote a random variable with a finite expected value and a finite, non-zero variance. Show that for any $a > 0$ the following inequality holds:

$$\mathbb{P}(|X - \mathbb{E}(X)| < a) > 1 - \frac{\text{Var}(X)}{a^2}.$$

Hint: use Markov's inequality.

18 – ([Chernoff inequality for sum of Bernoulli trials](#)) Let X_1, \dots, X_n be independent Bernoulli trials such that $\mathbb{P}(X_i = 1) = p_i$. Let $X = \sum_{i=1}^n X_i$ and $\mu = \mathbb{E}(X)$. Show that

a) for any $\delta > 0$

$$\mathbb{P}(X \geq (1 + \delta)\mu) \leq \left(\frac{e^\delta}{(1 + \delta)^{(1 + \delta)}} \right)^\mu,$$

b) for any $0 < \rho \leq 1$

$$\mathbb{P}(X \leq (1 - \rho)\mu) \leq \left(\frac{e^{-\rho}}{(1 - \rho)^{(1 - \rho)}} \right)^\mu.$$

Hint: see chapter 4.2. in [this book](#).

19 – Using the notations and the inequalities obtained in the previous task, show that for any $0 < \delta < 1$

$$\mathbb{P}(|X - \mu| \geq \delta\mu) \leq 2e^{-\mu\delta^2/3}.$$

20 – Let S_n be the number of heads in n flips of a symmetrical coin. Show that

a) using Chebyshev's inequality we have

$$\mathbb{P} \left(\left| S_n - \frac{n}{2} \right| \geq \frac{n}{4} \right) \leq \frac{4}{n},$$

b) using Chernoff's inequality from the previous task we have

$$\mathbb{P} \left(\left| S_n - \frac{n}{2} \right| \geq \frac{n}{4} \right) \leq 2e^{-n/24}.$$

21 – Consider the following algorithm, from which the idea of the HyperLogLog algorithm is derived.

Probabilistic Counter

1: Initialization: $C \leftarrow 1$

Upon event:

2: **if** random() $\leq 2^{-C}$ **then** \triangleright random returns a random number in a range $[0, 1)$

3: $C \leftarrow C + 1$

4: **end if**

In other words, when an event occurs, we toss a coin C times, and if each time we get heads, we increment the C counter by one. Otherwise, we do nothing. Let C_n be the value stored in the counter C after observing n events. Show that $\mathbb{E}(2^{C_n}) = n+2$ and $\mathbb{Var}(2^{C_n}) = \frac{1}{2}n(n+1)$. Based on C_n , define an unbiased estimator of n and calculate its variance.

Good luck,
J.L.