

Images

 Writeup

Case #1

(/call-prep/msk/casestacks
radiographs/pelviship/si
xr-pelviship-case1)
hip-og-se2)/pelvist

Demographics: 59 years old, Male

Indication: Motorcycle accident

Case #1

FREE | View all ↗ (/cases-index?free=1)

Findings

▶ Read



- [Diastasis of the pubic symphysis \(/call-prep/msk-radiographs/pelviship/classic/msk-xr-pelviship-case1?annotation=406\)](/call-prep/msk-radiographs/pelviship/classic/msk-xr-pelviship-case1?annotation=406) measuring up to 3 cm
- Mild [widening of the right sacroiliac joint \(/call-prep/msk-radiographs/pelviship/classic/msk-xr-pelviship-case1?annotation=412\)](/call-prep/msk-radiographs/pelviship/classic/msk-xr-pelviship-case1?annotation=412)
- Mild degenerative changes of the hips, sacroiliac joints, and of the partially visualized lower lumbar spine
- Arterial and vas deferens calcifications

Diagnosis

▶ Read



Unstable pelvic ring injury (APC II)

Sample Report

Unstable pelvic ring injury with diastasis of the pubic symphysis measuring up to 3 cm and mild widening of the right sacroiliac joint.

Mild degenerative changes of the hips, sacroiliac joints, and of the partially visualized lower lumbar spine.

Arterial and vas deferens calcifications.

Discussion

▶ Read



- It is important to be able to recognize unstable pelvic injuries because this impacts immediate patient management (pelvic binders are typically placed before the patient even goes for CT imaging)
- The Young Burgess classification is the most commonly used system:
 - **Anterior-posterior compression (APC)** - the pubic symphysis is the weak link, so it is affected first

- APC I: widening of the pubic symphysis < 2.5 cm
 - Stable
- APC II: widening of the pubic symphysis > 2.5 cm, anterior sacroiliac joint widening
 - Vertically stable, rotationally unstable
- APC III: widening of the pubic symphysis > 5 cm, widening of anterior and posterior sacroiliacs joints
 - Globally unstable
- **Lateral compression** - the obturator ring is the weak link, so it is affected first
 - LC I: obturator ring fractures and sacral compression fractures
 - Stable
 - LC II: LC I + iliac crescent fracture
 - Vertically stable, rotationally unstable
 - LC III: LC I/II with a contralateral APC injury
 - Globally unstable
- **Vertical shear** - Vertical hemipelvis translation with offset of the pubic symphysis and sacroiliac joint
 - Globally unstable

Annotated Images & Illustrations



APC II unstable pelvic ring...



Postoperative radiograph in..

References 

Leave Feedback