



**POLITECNICO**  
MILANO 1863


# Coupled Markov chains with applications to Approximate Bayesian Computation for model based clustering

E. Bertoni, M. Caldarini, F. Di Filippo, G. Gabrielli, E. Musiani  
10 January 2022


# Introduction

## A complex problem

1/14



Unbiased Markov chain  
Monte Carlo methods with  
couplings




likelihood

intractable



Approximate Bayesian  
Computation




# Unbiased Markov chain Monte Carlo methods with couplings

Faster MCMC  $\implies$  Parallelization

Faster MCMC  $\implies$  Parallelization  $\iff$  **Unbiased estimator**

Faster MCMC  $\implies$  Parallelization  $\iff$  **Unbiased estimator**

Exact estimations algorithms  
using **coupling of Markov  
chain.**



The goal is to estimate

$$\mathbb{E}_{\pi}[h(X)] = \int h(x)\pi(dx).$$

The estimator we are going to construct is based on a coupled pair of Markov chains,  $(X_t)_{t \geq 0}$  and  $(Y_t)_{t \geq 0}$ , which marginally start from  $\pi_0$  and evolve accordingly to  $P$ .

We consider some assumptions:

① as  $t \rightarrow \infty$ ,

$$\mathbb{E}[h(X_t)] \rightarrow \mathbb{E}_\pi[h(X)];$$

and there exists  $\eta > 0$  and  $D < \infty$  such that  $\mathbb{E}[|h(X_t)|^{2+\eta}] \leq D$  for all  $t \geq 0$ ;



We consider some assumptions:

- ① as  $t \rightarrow \infty$ ,

$$\mathbb{E}[h(X_t)] \rightarrow \mathbb{E}_\pi[h(X)];$$

and there exists  $\eta > 0$  and  $D < \infty$  such that  $\mathbb{E}[|h(X_t)|^{2+\eta}] \leq D$  for all  $t \geq 0$ ;

- ② the chains are such that the meeting time

$$\tau = \inf\{t \geq 1 : X_t = Y_{t-1}\}$$

satisfies  $\mathbb{P}(\tau > t) \leq C\delta^t$  for all  $t \geq 0$ , for some constants  $C < \infty$  and  $\delta \in (0, 1)$ ;

We consider some assumptions:

- ① as  $t \rightarrow \infty$ ,

$$\mathbb{E}[h(X_t)] \rightarrow \mathbb{E}_\pi[h(X)];$$

and there exists  $\eta > 0$  and  $D < \infty$  such that  $\mathbb{E}[|h(X_t)|^{2+\eta}] \leq D$  for all  $t \geq 0$ ;

- ② the chains are such that the meeting time

$$\tau = \inf\{t \geq 1 : X_t = Y_{t-1}\}$$

satisfies  $\mathbb{P}(\tau > t) \leq C\delta^t$  for all  $t \geq 0$ , for some constants  $C < \infty$  and  $\delta \in (0, 1)$ ;

- ③ the chains stay together after meeting:

$$X_t = Y_{t-1} \text{ for all } t \geq \tau.$$

Thanks to the previous assumptions we can prove that:

$$\mathbb{E}_{\pi}[h(X)] = \mathbb{E}[h(X_k) + \sum_{t=k+1}^{\tau-1} \{h(X_t) - h(Y_{t-1})\}];$$

Thanks to the previous assumptions we can prove that:

$$\mathbb{E}_\pi[h(X)] = \mathbb{E}[h(X_k) + \sum_{t=k+1}^{\tau-1} \{h(X_t) - h(Y_{t-1})\}];$$

and we define the Rhee–Glynn estimator as:

$$H_k(X, Y) = h(X_k) + \sum_{t=k+1}^{\tau-1} \{h(X_t) - h(Y_{t-1})\}$$

which is **unbiased** by construction.

$$H_{k:m}(X, Y) = \frac{1}{m - k + 1} \sum_{l=k}^m h(X_l) + \sum_{l=k+1}^{\tau-1} \min(1, \frac{l - k}{m - k + 1}) \{h(X_l) - h(Y_{l-1})\}$$

$$H_{k:m}(X, Y) = \underbrace{\frac{1}{m-k+1} \sum_{l=k}^m h(X_l)}_{MCMC_{k:m}} + \sum_{l=k+1}^{\tau-1} \min(1, \frac{l-k}{m-k+1}) \{h(X_l) - h(Y_{l-1})\}$$

- $MCMC_{k:m}$  is the standard MCMC average;

$$H_{k:m}(X, Y) = \underbrace{\frac{1}{m-k+1} \sum_{l=k}^m h(X_l)}_{MCMC_{k:m}} + \underbrace{\sum_{l=k+1}^{\tau-1} \min(1, \frac{l-k}{m-k+1}) \{h(X_l) - h(Y_{l-1})\}}_{BC_{k:m}}$$

- $MCMC_{k:m}$  is the standard MCMC average;
- $BC_{k:m}$  is the bias correction;

## Time-averaged estimator II

- ① draw  $X_0$  and  $Y_0$  from an initial distribution  $\pi_0$  and draw  $X_1 \sim P(X_0, \cdot)$ ;
- ② set  $t = 1$ : while  $t < \max\{m, \tau\}$  and:
  - a draw  $(X_{t+1}, Y_t) \sim \bar{P}\{(X_t, Y_{t-1}), \cdot\}$ ;
  - b set  $t \leftarrow t + 1$ ;
- ③ compute the time-averaged estimator:

$$H_{k:m}(X, Y) = \frac{1}{m - k + 1} \sum_{l=k}^m h(X_l) + \sum_{l=k+1}^{\tau-1} \min(1, \frac{l - k}{m - k + 1}) \{h(X_l) - h(Y_{l-1})\}.$$



## Time-averaged estimator II

- ① draw  $X_0$  and  $Y_0$  from an initial distribution  $\pi_0$  and draw  $X_1 \sim P(X_0, \cdot)$ ;
- ② set  $t = 1$ : while  $t < \max\{m, \tau\}$  and:
  - a draw  $(X_{t+1}, Y_t) \sim \bar{P}\{(X_t, Y_{t-1}), \cdot\}$ ;  $\bar{P}$  must be evaluated before!
  - b set  $t \leftarrow t + 1$ ;
- ③ compute the time-averaged estimator:

$$H_{k:m}(X, Y) = \frac{1}{m - k + 1} \sum_{l=k}^m h(X_l) + \sum_{l=k+1}^{\tau-1} \min(1, \frac{l - k}{m - k + 1}) \{h(X_l) - h(Y_{l-1})\}.$$

Metropolis–Hasting algorithm allow us to calculate the coupled kernel  $\bar{P}\{(X_t, Y_{t-1}), \cdot\}$ :

- 1 sample  $(X^*, Y^*)|(X_t, Y_{t-1})$  from a maximal coupling of  $q(X_t, \cdot)$  and  $q(Y_{t-1}, \cdot)$ ;
- 2 sample  $U \sim \mathcal{U}([0, 1])$ ;

3 if

$$U \leq \min \left\{ 1, \frac{\pi(X^*)q(X^*, X_t)}{\pi(X_t)q(X_t, X^*)} \right\}$$

then  $X_{t+1} = X^*$ ; otherwise  $X_t = X_{t-1}$ ;

4 if

$$U \leq \min \left\{ 1, \frac{\pi(Y^*)q(Y^*, Y_t)}{\pi(Y_t)q(Y_t, Y^*)} \right\}$$

then  $Y_{t+1} = Y^*$ ; otherwise  $Y_t = Y_{t-1}$ .

Metropolis–Hasting algorithm allow us to calculate the coupled kernel  $\bar{P}\{(X_t, Y_{t-1}), \cdot\}$ :

- 1 sample  $(X^*, Y^*)|(X_t, Y_{t-1})$  from a maximal coupling of  $q(X_t, \cdot)$  and  $q(Y_{t-1}, \cdot)$ ;
- 2 sample  $U \sim \mathcal{U}([0, 1])$ ;

- 3 if

$$U \leq \min \left\{ 1, \frac{\pi(X^*)q(X^*, X_t)}{\pi(X_t)q(X_t, X^*)} \right\}$$

then  $X_{t+1} = X^*$ ; otherwise  $X_t = X_{t-1}$ ;

- 4 if

$$U \leq \min \left\{ 1, \frac{\pi(Y^*)q(Y^*, Y_t)}{\pi(Y_t)q(Y_t, Y^*)} \right\}$$

then  $Y_{t+1} = Y^*$ ; otherwise  $Y_t = Y_{t-1}$ .



# Approximate Bayesian Computation

*Inputs:*

- a target posterior density  $\pi(\theta|y_{obs}) \propto p(y_{obs}|\theta)\pi(\theta)$ , consisting of a prior distribution  $\pi(\theta)$  and a procedure of generating data under the model  $p(y_{obs}|\theta)$ ;
- a proposal density  $g(\theta)$ , with  $g(\theta) > 0$  if  $\pi(\theta|y_{obs}) > 0$ ;
- an integer  $N > 0$ ;
- a kernel function  $K_h(u)$  and a scale parameter  $h > 0$ ;
- a low dimensional vector of summary statistics  $s = S(y)$ .

*Sampling* for  $i = 1, \dots, N$ :

- ① generate  $\theta^{(i)} \sim g(\theta)$  from sampling density  $g$ ;
- ② generate  $y \sim p(y|\theta^{(i)})$  from the likelihood;
- ③ compute summary statistic  $s = S(y)$ ;
- ④ accept  $\theta^{(i)}$  with probability  $\frac{K_h(\|s - s_{obs}\|)\pi(\theta^{(i)})}{Kg(\theta^{(i)})}$ , where  
 $K \geq K_h(0) \max_{\theta} \frac{\pi(\theta)}{g(\theta)}$ ; else go to 1.

*Output:*

- a set of parameter vectors  $\theta^{(1)}, \dots, \theta^{(N)} \sim \pi_{ABC}(\theta|S_{obs})$ .











# Conclusions

The next step will be the conclusion of the **separate multivariate implementation** of both solution to be tested on simulated data and the **parallelized multivariate implementation**.

Further steps will be testing on more complex data.

Pierre Jacob, John O'Leary, and Yves Atchadé.

Unbiased markov chain monte carlo with couplings.

*Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 82, 08 2017.

Peter W. Glynn and Chang han Rhee.

Exact estimation for markov chain equilibrium expectations, 2014.

Jeffrey S. Rosenthal.

Faithful couplings of markov chains: Now equals forever.

*Advances in Applied Mathematics*, 18(3):372–381, 1997.

Dylan Cordaro.

Markov chain and coupling from the past.

2017.

Jinming Zhang.

Markov chains, mixing times and coupling methods with an application in social learning.

2020.

S. A. Sisson, Y. Fan, and M. A. Beaumont.

Overview of approximate bayesian computation, 2018.

Y. Fan and S. A. Sisson.

Abc samplers, 2018.

Dennis Prangle.

Summary statistics in approximate bayesian computation, 2015.