




POLITECNICO
MILANO 1863


Coupled Markov chains with applications to Approximate Bayesian Computation for model based clustering

E. Bertoni, M. Caldarini, F. Di Filippo, G. Gabrielli, E. Musiari

10 January 2022



Unbiased Markov chain Monte Carlo methods with couplings



likelihood

intractable



Approximate Bayesian Computation



Unbiased Markov chain Monte Carlo methods with couplings

Time-averaged estimator

- ① draw X_0 and Y_0 from an initial distribution π_0 and draw $X_1 \sim P(X_0, \cdot)$;
- ② set $t = 1$: while $t < \max\{m, \tau\}$ and:
 - a draw $(X_{t+1}, Y_t) \sim \bar{P}\{(X_t, Y_{t-1}), \cdot\}$;
 - b set $t \leftarrow t + 1$;
- ③ compute the time-averaged estimator:

$$H_{k:m}(X, Y) = \frac{1}{m - k + 1} \sum_{l=k}^m h(X_l) + \sum_{l=k+1}^{\tau-1} \min(1, \frac{l - k}{m - k + 1}) \{h(X_l) - h(Y_{l-1})\}.$$

Metropolis–Hasting algorithm for a coupled kernel

Metropolis–Hasting algorithm allow us to calculate the coupled kernel $\bar{P}\{(X_t, Y_{t-1}), \cdot\}$:

- 1 sample $(X^*, Y^*) | (X_t, Y_{t-1})$ from a maximal coupling of $q(X_t, \cdot)$ and $q(Y_{t-1}, \cdot)$;
- 2 sample $U \sim \mathcal{U}([0, 1])$;

- 3 if

$$U \leq \min \left\{ 1, \frac{\pi(X^*)q(X^*, X_t)}{\pi(X_t)q(X_t, X^*)} \right\}$$

then $X_{t+1} = X^*$; otherwise $X_t = X_{t-1}$;

- 4 if

$$U \leq \min \left\{ 1, \frac{\pi(Y^*)q(Y^*, Y_t)}{\pi(Y_t)q(Y_t, Y^*)} \right\}$$

then $Y_{t+1} = Y^*$; otherwise $Y_t = Y_{t-1}$.

Maximal coupling

The algorithm:

Set $p = \mathcal{N}(X_{t-1}, 1)$ and $q = \mathcal{N}(Y_{t-1}, 1)$,

- ① sample $X_t \sim p$;
- ② sample $W|X_t \sim \mathcal{U}\{[0, p(X_t)]\}$;
- ③ if $W \leq q(X_t)$ then output (X_t, X_t) , otherwise:
 - ① sample $Y_t \sim q$;
 - ② sample $W^*|Y_t \sim \mathcal{U}\{[0, q(Y_t)]\}$ until $W^* > p(Y_t)$ and output (X_t, Y_t) .

Study case

Model

$$Y_i | \mu \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma_{obs}^2)$$

$$\mu \sim \mathcal{N}(\mu_0, \sigma_0^2)$$

$$\mu_0 = 38, \quad \sigma_0^2 = 4$$

Dataset

1000 samples generated from a Gaussian distribution:

$$Y_{obs} \sim \mathcal{N}(\mu_{obs}, \sigma_{obs}^2)$$

$$\mu_{obs} = 43, \quad \sigma_{obs}^2 = 5$$

Implementation and parallelization

Results

7/14

$$\mathcal{N}(\mu_n, \sigma_n^2)$$

$$\mu_n = y = \text{tfd.Normal}(1/(1/\sigma_0^2 * 2 + n/\sigma_{obs}^2)(\mu_0/\sigma_0^2 * 2 + (np.sum(y)/\sigma_{obs}^2)), 1/(1/\sigma_0^2 + n/\sigma_{obs}^2 * 2))$$

- posterior
- risultato del time averaged
- grafico catene che si incontrano
- istogramma dei samplings (sotto alla posterior)



Approximate Bayesian Computation

NO, UTILIZZARE ALGORITMO DI PAGINA 103 *Inputs:*

- a target posterior density $\pi(\theta|y_{obs}) \propto p(y_{obs}|\theta)\pi(\theta)$, consisting of a prior distribution $\pi(\theta)$ and a procedure of generating data under the model $p(y_{obs}|\theta)$;
- a proposal density $g(\theta)$, with $g(\theta) > 0$ if $\pi(\theta|y_{obs}) > 0$;
- an integer $N > 0$;
- a kernel function $K_h(u)$ and a scale parameter $h > 0$;
- a low dimensional vector of summary statistics $s = S(y)$.

ABC rejection sampling algorithm

NO, UTILIZZARE ALGORITMO DI PAGINA 103 *Sampling* for $i = 1, \dots, N$:

- ① generate $\theta^{(i)} \sim g(\theta)$ from sampling density g ;
- ② generate $y \sim p(y|\theta^{(i)})$ from the likelihood;
- ③ compute summary statistic $s = S(y)$;
- ④ accept $\theta^{(i)}$ with probability $\frac{K_h(\|s - s_{obs}\|)\pi(\theta^{(i)})}{Kg(\theta^{(i)})}$, where
 $K \geq K_h(0) \max_{\theta} \frac{\pi(\theta)}{g(\theta)}$; else go to 1.

Output:

- a set of parameter vectors $\theta^{(1)}, \dots, \theta^{(N)} \sim \pi_{ABC}(\theta|S_{obs})$.

summary statistics: media distanza: norma (modulo della differenza)
(mahalanobis nel caso multivariato) kernel:
 $1/(\text{np.sqrt}(2*\text{math.pi}))\text{np.exp}(-1/2*u^2)$

Model

$$Y_i | \mu \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma_{obs}^2)$$

$$\mu \sim \mathcal{N}(\mu_0, \sigma_0^2)$$

$$\mu_0 = 38, \quad \sigma_0^2 = 4$$

Dataset

1000 samples generated from a Gaussian distribution:

$$Y_{obs} \sim \mathcal{N}(\mu_{obs}, \sigma_{obs}^2)$$

$$\mu_{obs} = 43, \quad \sigma_{obs}^2 = 5$$



Conclusions

The next step will be the conclusion of the **separate multivariate implementation** of both solution to be tested on simulated data and the **parallelized multivariate implementation**.

Further steps will be testing on more complex data.

Pierre Jacob, John O'Leary, and Yves Atchadé.

Unbiased markov chain monte carlo with couplings.

Journal of the Royal Statistical Society: Series B (Statistical Methodology), 82, 08 2017.

Peter W. Glynn and Chang han Rhee.

Exact estimation for markov chain equilibrium expectations, 2014.

Jeffrey S. Rosenthal.

Faithful couplings of markov chains: Now equals forever.

Advances in Applied Mathematics, 18(3):372–381, 1997.

Dylan Cordaro.

Markov chain and coupling from the past.

2017.

Jinming Zhang.

Markov chains, mixing times and coupling methods with an application in social learning.

2020.

S. A. Sisson, Y. Fan, and M. A. Beaumont.

Overview of approximate bayesian computation, 2018.

Y. Fan and S. A. Sisson.

Abc samplers, 2018.

Dennis Prangle.

Summary statistics in approximate bayesian computation, 2015.