




**POLITECNICO**  
MILANO 1863


# **Coupled Markov chains with applications to Approximate Bayesian Computation for model based clustering**

**E. Bertoni, M. Caldarini, F. Di Filippo, G. Gabrielli, E. Musiari**

10 January 2022



**Unbiased Markov chain Monte Carlo methods with couplings**



likelihood

**intractable**



**Approximate Bayesian Computation**



# Unbiased Markov chain Monte Carlo methods with couplings

## Time-averaged estimator

- ① draw  $X_0$  and  $Y_0$  from an initial distribution  $\pi_0$  and draw  $X_1 \sim P(X_0, \cdot)$ ;
- ② set  $t = 1$ : while  $t < \max\{m, \tau\}$  and:
  - a draw  $(X_{t+1}, Y_t) \sim \bar{P}\{(X_t, Y_{t-1}), \cdot\}$ ;
  - b set  $t \leftarrow t + 1$ ;
- ③ compute the time-averaged estimator:

$$H_{k:m}(X, Y) = \frac{1}{m - k + 1} \sum_{l=k}^m h(X_l) + \sum_{l=k+1}^{\tau-1} \min(1, \frac{l - k}{m - k + 1}) \{h(X_l) - h(Y_{l-1})\}.$$

## Metropolis–Hasting algorithm for a coupled kernel

Metropolis–Hasting algorithm allow us to calculate the coupled kernel  $\bar{P}\{(X_t, Y_{t-1}), \cdot\}$ :

- 1 sample  $(X^*, Y^*) | (X_t, Y_{t-1})$  from a maximal coupling of  $q(X_t, \cdot)$  and  $q(Y_{t-1}, \cdot)$ ;
- 2 sample  $U \sim \mathcal{U}([0, 1])$ ;

- 3 if

$$U \leq \min \left\{ 1, \frac{\pi(X^*)q(X^*, X_t)}{\pi(X_t)q(X_t, X^*)} \right\}$$

then  $X_{t+1} = X^*$ ; otherwise  $X_t = X_{t-1}$ ;

- 4 if

$$U \leq \min \left\{ 1, \frac{\pi(Y^*)q(Y^*, Y_t)}{\pi(Y_t)q(Y_t, Y^*)} \right\}$$

then  $Y_{t+1} = Y^*$ ; otherwise  $Y_t = Y_{t-1}$ .

## Maximal coupling

The algorithm:

Set  $p = \mathcal{N}(X_{t-1}, 1)$  and  $q = \mathcal{N}(Y_{t-1}, 1)$ ,

- 1 sample  $X_t \sim p$ ;
- 2 sample  $W|X_t \sim \mathcal{U}\{[0, p(X_t)]\}$ ;
- 3 if  $W \leq q(X_t)$  then output  $(X_t, X_t)$ , otherwise:
  - 1 sample  $Y_t \sim q$ ;
  - 2 sample  $W^*|Y_t \sim \mathcal{U}\{[0, q(Y_t)]\}$  until  $W^* > p(Y_t)$  and output  $(X_t, Y_t)$ .

## Study case

## Model

$$Y_i | \mu \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma_{obs}^2)$$

$$\mu \sim \mathcal{N}(\mu_0, \sigma_0^2)$$

$$\mu_0 = 38, \quad \sigma_0^2 = 4$$

## Dataset


1000 samples generated from a Gaussian distribution:

$$Y_{obs} \sim \mathcal{N}(\mu_{obs}, \sigma_{obs}^2)$$

$$\mu_{obs} = 43, \quad \sigma_{obs}^2 = 5$$

# Implementation


6/14





# Parallelization


7/14




## Results

$$\mathcal{N}(\mu_n, \sigma_n^2), \quad \mu_n = \frac{1}{\frac{1}{\sigma_0^2} + \frac{n}{\sigma_{obs}^2}} \cdot \left( \frac{\mu_0}{\sigma_0^2} + \frac{\sum y_{obs}}{\sigma_{obs}^2} \right), \quad \sigma_n^2 = \frac{1}{\frac{1}{\sigma_0^2} + \frac{n}{\sigma_{obs}^2}}$$


Coupled chains



Complete sampling



Sampling histogram



Time Averaged Estimators mean:

$$\mathbb{E}[H_{k:m}(X, Y)] = 42.9498$$



# Approximate Bayesian Computation

*Inputs:*

- a target posterior density  $\pi(\theta|y_{obs}) \propto p(y_{obs}|\theta)\pi(\theta)$ , consisting of a prior distribution  $\pi(\theta)$  and a procedure of generating data under the model  $p(y_{obs}|\theta)$ ;
- a Markov proposal density  $g(\theta, \theta')=g(\theta'|\theta)$ ;
- an integer  $N > 0$ ;
- a kernel function  $K_h(u)$  and a scale parameter  $h > 0$ ;
- a low dimensional vector of summary statistics  $s = S(y)$ .

*Initialise:*

repeat:

- ① choose an initial parameter vector  $\theta^{(0)}$  from the support of  $\pi(\theta)$ ;
- ② generate  $y^0 \sim p(y|\theta^{(0)})$  from the model and compute summary statistics  $s^0 = S(y^{(0)})$ , until  $K_h(\|s^{(0)} - s_{obs}\|) > 0$ .

*Sampling for  $i = 1, \dots, N$ :*

- 1 generate candidate vector  $\theta' \sim g(\theta^{(i-1)}, \theta)$  from the proposal density  $g$ ;
- 2 generate  $y' \sim p(y|\theta')$  from the model and compute summary statistics  $s' = S(y')$ ;
- 3 with probability

$$\min\left\{1, \frac{K_h(\|s' - s_{obs}\|) \pi(\theta') g(\theta', \theta^{(i-1)})}{K_h(\|s^{(i-1)} - s_{obs}\|) \pi(\theta^{(i-1)}) g(\theta^{(i-1)}, \theta')}\right\}$$

set  $(\theta^{(i)}, s^{(i)}) = (\theta', s')$ . Otherwise set  $(\theta^{(i)}, s^{(i)}) = (\theta^{(i-1)}, s^{(i-1)})$ .

*Output:*

- a set of correlated parameter vectors  $\theta^{(1)}, \dots, \theta^{(N)}$  from a Markov chain with stationary distribution  $\pi_{ABC}(\theta|S_{obs})$ .

**Summary statistic:**

Mean

**Distance:**

Norm (absolute value) of the difference.

**Kernel:**

$$\frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}u^2}.$$

Same as previous:

### Model

$$Y_i | \mu \stackrel{iid}{\sim} \mathcal{N}(\mu, \sigma_{obs}^2)$$

$$\mu \sim \mathcal{N}(\mu_0, \sigma_0^2)$$

$$\mu_0 = 38, \quad \sigma_0^2 = 4$$

### Dataset


1000 samples generated from a Gaussian distribution:

$$Y_{obs} \sim \mathcal{N}(\mu_{obs}, \sigma_{obs}^2)$$

$$\mu_{obs} = 43, \quad \sigma_{obs}^2 = 5$$








## **The complete method: Monte Carlo Markov chain method with couplings and approximate bayesian computation**

# The complete method: Monte Carlo Markov chain method with couplings and approximation

## Implementation

12/14





## Multivariate study case



## Conclusions

The next step will be the conclusion of the **separate multivariate implementation** of both solution to be tested on simulated data and the **parallelized multivariate implementation**.

Further steps will be testing on more complex data.

Pierre Jacob, John O'Leary, and Yves Atchadé.

Unbiased markov chain monte carlo with couplings.

*Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 82, 08 2017.

Peter W. Glynn and Chang han Rhee.

Exact estimation for markov chain equilibrium expectations, 2014.

Jeffrey S. Rosenthal.

Faithful couplings of markov chains: Now equals forever.

*Advances in Applied Mathematics*, 18(3):372–381, 1997.

Dylan Cordaro.

Markov chain and coupling from the past.

2017.

Jinming Zhang.

Markov chains, mixing times and coupling methods with an application in social learning.

2020.

S. A. Sisson, Y. Fan, and M. A. Beaumont.

Overview of approximate bayesian computation, 2018.

Y. Fan and S. A. Sisson.

Abc samplers, 2018.

Dennis Prangle.

Summary statistics in approximate bayesian computation, 2015.