

Machine Learning from Data – IDC

HW5 – Support Vectors Machine

Theoretical questions:

1. Use Lagrange Multipliers to find the maximum and minimum values of the function subject to the given constraints:
 - a. $f(x, y) = e^{xy}$; constraint: $2x^2 + y^2 = 72$
 - b. $f(x, y) = x^2 + y^2$; constraint: $y - \cos 2x = 0$
- 2.

- a. Consider two kernels K_1 and K_2 , with the mappings φ_1 and φ_2 respectively. Show that $K = 7K_1 + 3K_2$ is also a kernel and find its corresponding φ .
- b. Consider a kernel K_1 and its corresponding mapping φ_1 that maps from the lower space R^n to a higher space R^m ($m > n$). We know that the data in the higher space R^m , is separable by a linear classifier with the weights vector w .

Given a different kernel K_2 and its corresponding mapping φ_2 , we create a kernel $K = 7K_1 + 3K_2$ as in section a above. Can you find a linear classifier in the higher space to which φ , the mapping corresponding to the kernel K , is mapping?

If YES, find the linear classifier weight vector.

If NO, prove why not.

- c. What is the dimension of the mapping function φ that corresponds to a polynomial kernel $K(x, y) = (\alpha x \cdot y + \beta)^d$, ($\alpha, \beta \neq 0$), where the lower dimension is n ?
- d. Consider the space $S = \{1, 2, \dots, N\}$ for some finite N (each instance in the space is a 1-dimension vector and the possible values are $1, 2, \dots, M$) and the function $f(x, y) = \min(x, y)$.

Find the mapping φ such that:

$$\varphi(x) \cdot \varphi(y) = 5 \min(x, y)$$

For example, if the instances are $x = 3, y = 5$, for some $N \geq 5$, then:

$$\varphi(x) \cdot \varphi(y) = \varphi(3) \cdot \varphi(5) = 5 \min(3, 5) = 15$$

- e. Consider the space $S = \{1, 2, \dots, N\}$ for some finite N (each instance in the space is a 1-dimension vector and the possible values are $1, 2, \dots, M$) and the function $f(x, y) = \max(x, y)$.

Prove that the function $\max(x, y)$ is not a kernel, i.e., there is no mapping φ such that:

$$\varphi(x) \cdot \varphi(y) = \max(x, y)$$

3. Find the Kernel function for the following mapping. Provide a representation of the form $\alpha K_1 + \beta K_2$, where both K_1 and K_2 are polynomial kernels and $\alpha, \beta > 0$:
 - a. $\varphi(x) = (x_1^3, x_2^3, \sqrt{3}x_1^2x_2, \sqrt{3}x_1x_2^2, 2\sqrt{3}x_1^2, 2\sqrt{3}x_2^2, 2\sqrt{6}x_1x_2, 4\sqrt{3}x_1, 4\sqrt{3}x_2, 8)$
 - b. $\varphi(x) = (\sqrt{10}x_1^2, \sqrt{10}x_2^2, \sqrt{20}x_1x_2, \sqrt{8}x_1, \sqrt{8}x_2, \sqrt{2})$
4. The purpose of this exercise is to demonstrate the usefulness of using kernels. Write a Python script that performs the following:
 - a. Draw 20,000 vectors with 20 dimensions.
 - b. Use the kernel $(x \cdot y + 1)^2$ to calculate the Gram matrix for these data. That is: each cell i, j in the matrix is the result of applying the kernel function on the vectors i and j ($i=j$ is also a valid input) : $K(x_i, x_j)$
 - c. Find the associated mapping function φ . Into which dimension?
 - d. Use φ to map the vectors to the higher dimension.
 - e. Calculate the matrix where each cell i, j is the result of the dot product of the mapping images of the vectors i and j : $\varphi(x_i) \cdot \varphi(x_j)$
 - f. Compare the matrices from sections b and e (use `np.isclose`) – they should be the same.
* NOTE: both matrices should be of the same size 20,000x20,000.
 - g. Compare the time it took your machine to calculate the two matrices. What do you observe?