BSJE JOINT EXAMINATION -2024-

Kenya Certificate of Secondary Education

233/3

Chemistry

Paper 3

July, 2024

TIME:	2	hours	15	minutes
-------	---	-------	-----------	---------

Name:	ission No:
School:Signature:	<u>233/3 - Chemistry</u>
	Wednesday, 17 TH July, 2024
	8.00am-10.15am
INSTRUCTIONS TO CANDIDATES	(1 ST session)

- a) Write your name and Index number in the spaces provided.
- b) Sign and write the date of examination in the spaces provided.
- c) Answer all the questions in the spaces provided in the question paper.
- d) You are not allowed to start working with the apparatus for the first 15 minutes of the 2 \(^{1}\sqrt{4}\) hours allowed for this paper. This time is to enable you to read through the question paper and make sure you have all the chemicals and apparatus that you may need.
- e) This paper consists of 10 printed pages
- f) Candidates should check the question paper to ascertain that all the pages are printed as indicated and that no questions are missing.
- g) Candidates should answer the questions in **English**.

FOR EXAMINER'S USE

Question	Maximum score	Candidate's Score
1	19	
2	7	
3	14	

Download More Resources: Enovate KCSE Revision App Highschool.co.ke

TOTAL SCORE	40	

- 1. You are provided with;
- \triangleright 4.5g of solid **T** in a boiling tube.
- > Solution N, 0.06M acidified potassium manganate (VII).

You are required to determine;

- \checkmark The solubility of of solid **T** at different temperatures.
- ✓ The number of moles of water of crystalisation in solid **T**.

PROCEDURE I

- i. Using a burette, add 4cm³ of distilled water to solid **T** in the boiling tube. Heat the mixture while stirring with the thermometer to about 85°C for the solid to dissolve; allow the solution the solution to cool while stirring. Note the temperature at which crystals of solid **T** first appear. Record the temperature in **table 1**.
- ii. Using a burette ,add 2cm³ of distilled water to the contents of the boiling tube . warm the mixture while stirring with the thermometer until all the solid dissolve. Allow the mixture to cool while stirring. Note and record the temperature at which crystals of solid **T** first appear.
- iii. Repeat procedure (ii) two more times and record the temperature in table 1.

Retain the contents of the boiling tube for use in procedure II.

Table 1.

Total volume of water in in the boiling tube (cm ³)	4	6	8	10
Temperature at which crystals of T first appear (°C)				
Solubility of solid T (g/100g of water)				

(a) Complete the table 1 by calculating solubility of solid **T** at different temperatures. (3mks)

Download More Resources: Enovate KCSE Revision App

(2mks)

(b) On the grid provided, plot a graph of solubility against temperature.

-			 	11		1	+		+	+	-	++	+			-		+	++-	 	11	+	-	+	++	$\dagger \dagger$			-	-	++		+	-	++	tt	+++	Ħ
				П	‡	П							11					1							Ш						11		İ		1		П	#
-	\vdash		┼┼	++	+-	╁	+		++		+	++	++			+		+	┼┼	╁┼┼	++	+	+-	+	++	╁┼	+			╟┼	++		+	+	++	╁┼	₩	+
					İ	Ш					i	Ш	Ш																					士			ш	I
-		4	₩.	1		1	44		44	-		1	44			4			 	ļļ	1	- -	-		44	44	-		ļļ	Щ.	44		-	4	₩.		₩.	+i
+	\vdash		₩	++	+-	₩	+		++	+	-	++	++		\vdash	+	+++	+	+		+	+	+	+	++	++	+			╁┼	++	++	1	+	++	┼┼	₩	$\pm \pm$
										1	1		11					T.	tt.					1							11				<u>tt</u>		LL	± 1
		4			<u></u>	1	44		11	4		11	44				44	4	<u> </u>	Щ.		1	4	4	1	11					44	44	1		44	Ш	1	1
-			┢╌┝╌	\vdash	- -	-	+		+-+		<u>i</u>	╁	++						╁┼┼		╁	+-			+-+-	++	-				++		+		╫		-	₩i
						ΙÌ							ΙÌ						Ħ												ΙÌ		1		tt			
		4	Ш	Н	+	Ш	44	+	Ш	_	4	Ш	Н	\perp		Ш	ш	_	1	ш	Н	+	_	_	ш	Н	Ш			Н	ш	-		+	₩	₩	4	+i
			┼┼	++	<u>-</u>	₩	+		┿┽		-i-	┼┼	+-+						┼┼	╢	┼┼	++			+-+-	++	+				++		i		++	┼┼	┼┼-	⊹ i
		1		Ħ		ΙT	\top		Ħ	T	1	tt	$\dagger \dagger$					T	tt	ttt	TT			1	tt	TT					TT			1	tt	H	tt	
		1			1	П	11			1		Ш	11				Ш		Ш	Ш	П			_	Щ	\prod				П	ŢŢ			4	Ш	Ш	Ш	
-+-	┝╼┾╼┾		++	++	+-	 -+-	-4-4	-+-	4-4	-+	-+-	+-+-	-+-+	-+-+	-+-+	-+-		-+-	+-+-	┾┿╸	++	++	++	-+	++	 - -	+-+		}- +	 - -	+-+	-+-+			+-+-		++	
			叶	Ħ		tt	力	_	17	_	÷	±Τ	#					J	廿	Ħ	11		1		ΞŤ	Ιt	\pm			_+	廿			\pm	廾	Ħ	±٢	
		1			1	Ш	П				-	Ш	П					I	Ш					Ţ	Ш					П	П		1		Ш		Щ	
			-	₩	-	₩	+		+			₩	+						-		₩	+		-	++	₩	-				++		-		+		++-	+-
+		†		H	-	H	H	+	H	+		H	Ħ	+		Ħ	+	+	+	H	H		Н	+	Ħ	Ħ				+	$\dagger \dagger$		1 1	+	+	+	H	\pm
					1	Ш	11						11											1	П	\Box				П	11				\Box		Ш	
			┝┿	₩	+	₩	+-		+			++	+-+					-		⊢⊢	+-+				+-	++			-		++		H		++			+
+		+-	H	+ +	+-	H	+		+	+	+	++	++	+		\pm	+	+	++	+	+		+	+	++	++	+	-	-	H	+ +	++	 -	+	++	H	+	+
					7	П	11		\square			Ш	Ш									T		1	11	П					Ш	II	1		Ш	Ш	II	
				-	Ŧ	₩			4-4		÷	₩							 	├ ┼	-	\blacksquare					-		ļļ				H		₩.		} }-	4
-			-	\vdash	+-	H	+-		++	+	+	tt	++					+	++-	t-t-	+	++	+-	+	 	++					++		+		++	tt	tt	
					T						-																										ш	
-	\vdash		-	\vdash	+	-	+	-	++	+	-i-	₩	++			+i		+		┼┼┼	++	-	+	-		₩	+			-	++		1	-	++	₩		+-
+			H	$^{++}$	+-	H	+		$^{++}$	+	+	tt	++			+		+	++-	 -	$^{++}$	++	+	+	$^{++}$	tt	+			1	++	++	+	+	++	H	tt	+
						П							\Box													Ш					\blacksquare							
				++	.	-+-	-4-4		4-4	-+		 -+∙	-+-+		+-	-+-i			┾-┾-	┝╼┝╺	++	++		-4	-+-	-	4-4				-+-+	-+-+			┾┿╸		┾┿.	
+			++	11		1	+		++	+	+	++	+					+	+++	t	11	+	+	+	++	††	+		-	-	++	+++	1	-	++	tt	 	
					1	П					1		\Box													\Box					\Box				ш		Ш	
				-	 	₩			+			₩									-					+-+	-						÷		₩			
+		+	H	H	÷	Н	+	+	Н	+	+	H	H	+		\pm	\Box	+	+	+	H	+	Н	+	+	H	Н			Н	+	-		+	+	\vdash	H	\pm
					1	Ш					1		Ш						Ш					1		П					Ш			1	Ш	Ш	Ш	
	\vdash		-	++	+-	₩	+		++			++	++			-	\vdash				++	-			++	₩	-				+	++	+-		++		₩	+
+			tt	$\dagger\dagger$	- 	tt	+		++	+	+	++	++	+		+	+	+	++	+	++	+	+	+	$^{++}$	$\dagger \dagger$		-	-	1-1-	++	++	\vdots	+	++	t†	++	+
7		T		П	1	П	11	1	11	1	1	Ш	11	11		1		7			[]	I	T	7	П	Ш					11	II		4	ፗፗ	[]	II.	
-			-	₩	-	₩	+		+		+	₩	++								₩	+	-	-	+-	₩	-				++		-		+	 	₩-	
+	\vdash		tt	$\dagger\dagger$		H	+		+	+	+	tt	+ +	+			+	+	$^{++}$	+	††	+	+	+	$^{++}$	††	+			1-1-	+	++	+	-	++	tt	tt	-
		T		П	1	П		#	П	T		П	\Box					T	П	П				I	П	П					П			#	П	П	П	
				₩		₩			+			₩	+-+							₩.	+	+-			₩	+-+	-				++		÷	-	+	₩.	₩.	-
+			H	+	+-	H	+		++	+		++	++	+		+	+	+	++-	++	++	+	++	+	++	+	+		-	++	+ +	++	: 1	+	++-	H	++	17
					-	Ш							11													П					11		H		#		Ш	
-+-	┝╼┾╼┾	+-	++	++	÷-	₽4-	-4-4	-4-	4-4	-4	-‡-	 - -	-+-+	-+-+	-+-	-+-			+-+-	++-	++	++	+-	-4	++	 - -		-4	- ∔	 - -	-+-+	-+-+	+-+		+-+-		++	+-
+			++	++		+	+		++	+	+	++	++	+		+	+	+	++-	++	++	+	++	+	++	++	+			-	++	++		+	++-	t+	++	
					+						1		\Box												П						\blacksquare				Ш		Ш	
			-	\vdash	÷	₩	+		+		į.	₩	++					-		₩.	++	+	-	-	+-	₩	-				++		+	-	+	┿	₩.	+
+		+	H	H		H	+	+	H	+	+	H	H	+		\pm	Н	+	+	H	H	+	Н	+	H	H	н	+	H	+	+	+		+	+	+	H	H
					Ť	II	##						##					1						#		茸					##				ፗ		ፗ	
-			-	₩	+	₩	+		+		-i-	₩	+	+		-		-	-	-	++	-	-	-	++	₩	-		-		$+\!+$			-	+-	-	₩	+i
	$\vdash \vdash \vdash$			++		₩	-++		+-+			+-+-			\vdash				 		+-+				+-+-	++-			├}	├					+	├├- -	₩.	+-1

solid T is dissolved in 100cm ³ of distilled water at	68°C then the solution is allowed to
cool.	(1mk)

Procedure 2

Transfer all the contents of the boiling tube into a 250ml volumetric flask. Rinse the boiling tube with distilled water and transfer the rinses to the volumetric flask. Add more distilled water to the flask to top it to the mark. **Label this as solution T**. Pipette 25cm³ of solution **T** into a 250ml conical flask. Fill the burette with solution **N**. Titrate solution **T** with solution **N** from the burette until the contents of the conical flask first become pink. Repeat the titration two more times and complete the table 2 below.

(5mks)

Table 2

	Ι	II	III
Final burette reading (cm ³)			
Initial burette reading (cm ³)			
Volume of solution N (cm ³)			

Calculate;	
(a) The average volume of solution N used.	(1mk)
(b) The number of moles of N that reacted.	(1mk)
(c) The number of moles of T that reacted with solution N . (mole ratio of N : T is	2:5)(1mk)

Download More Resources: Enovate KCSE Revision App

(d	I) The number of moles of T in 10cm ³ of t	he original solution (in procedure 1). (2	
(e	e) If solid T was $C_2H_2O_4$.X H_2O , determine		2mks)
2. Yo	u are provided with substance S . Use it to	carry out the tests that follow. Record vo	
	vations and inferences in the spaces provide	•	, di
(a) D	escribe the appearance of substance S .	(1/2)	nark)
(b)Sc	oop a little of substance S using a clean m		
	oop a little of substance S using a clean m		
(b)Sc	oop a little of substance S using a clean m	etallic spatula and ignite it in a Bunsen b	
(b)Sc	oop a little of substance S using a clean m	etallic spatula and ignite it in a Bunsen b	

(c)Pla	ace about 2cm ³ of substance S in a clean ter	st tube and add solid P , sodium hydrogen
carbo	onate provided.	
	Observations	Inferences
	(1mark)	(½ mark)
(b	b) Place about 2cm³ of substance S in a test dichromate(VI) and warm the mixture.	t tube and add 2cm ³ of acidified potassium
	Observations	Inferences
	(1mark)	(1mark)
(e)Pla	ace about 2cm ³ of substance S and add 2cm	n ³ of distilled water.
	Observations	Inferences
	(1mark)	(1mark)
	(IIIIaik)	(IIIaik)

- 3. You are provided with solid **Q**. Carry out the following tests and write your observations and inferences in the spaces provided.
 - (a) Place about a half of solid Q in a test tube and heat strongly. Test any gases produced with both red and blue litmus papers.

Observations	Inferences
(2mlsa)	(1 mouls)
(2mks)	(1mark)

(b) Place the rest of solid \mathbf{Q} in a boiling tube and add about 10cm^3 of distilled water. Shake well.

To a 2cm³ portion of the solution, add about 1cm³ of hydrogen peroxide and shake well. To the resulting mixture, add aqueous sodium hydroxide drop wise until in excess.

Observations	Inferences
(1 ½ marks)	(½ mark)

- (c) You are provided with solution \mathbf{R} . Carry out the following tests and write your observations in the spaces provided. Divide solution \mathbf{R} into two portions.
- (i) To the first portion, of solution \mathbf{R} , add three drops of barium nitrate solution. Retain the mixture for use in test (ii) below.

Download More Resources: Enovate KCSE Revision App

Observations	Inferences
(1mark)	(1mark)

(ii) To the mixture obtained in (c(i) above, add about 2cm³ of 2M nitric V acid.

Observations	Inferences
(1mark)	(1mark)

(iii) To the second portion of solution \mathbf{R} , add two drops of acidified potassium dichromate(VI) and warm the mixture.

Observations	Inferences
(1mark)	(1mark)

Download More Resources: <u>Enovate KCSE Revision App</u>

d). To 2cm^3 of solution $\mathbf{R_1}$ in a test –tube add 5 drops of aqueous sodium hydroxide. Add a piece of aluminium foil provided to the mixture and shake. Warm the mixture and test for any gas produced with both red and blue litmus papers.

Observation	Inference
(2mks)	(1mk)