

MATH 4441 Homework 3

September 12, 2018

4.1

Assume $\alpha(t)$ has unit speed, then $\kappa = \|a''\|$. If $\kappa = 0$ then $a'' = 0$,

$$\alpha'(t) = v_0 + \int_0^t \alpha''(s) ds = v_0$$

$$\alpha(t) = x_0 + \int_0^t \alpha'(s) ds = x_0 + v_0 t$$

which is the equation of a straight line.

4.2

By definition $\|T\| = \langle T(t), T(t) \rangle = 1$. Differentiate and then scale both sides:

$$\langle T(t), T(t) \rangle' = 2\langle T(t), T(t)' \rangle = 0$$

$$\left\langle T(t), \frac{T(t)'}{\|T'(t)\|} \right\rangle = N(t) = 0$$

4.4

For a given t , the osculating circle γ can be parameterized by

$$\gamma(s) = p(t) + \frac{1}{\kappa(t)} T(t) \cos s + \frac{1}{\kappa(t)} N(t) \sin s$$

where

$$p(t) := \alpha(t) + \frac{1}{\kappa(t)} N(t)$$

And we can compute the tantrix for the osculating circle:

$$\gamma'(s) = -\frac{1}{\kappa(t)} T(t) \sin s + \frac{1}{\kappa(t)} N(t) \cos s$$

$$\|\gamma'(s)\| = \langle \gamma'(s), \gamma'(s) \rangle^{1/2} = \frac{1}{\kappa(t)}$$

$$\bar{T}(s) = \frac{\gamma'(s)}{\|\gamma'(s)\|} = -T(t) \sin s + N(t) \cos s$$

$$\bar{T}'(s) = -T(t) \cos s - N(t) \sin s$$

$$\|\bar{T}'(s)\| = \langle \bar{T}'(s), \bar{T}'(s) \rangle^{1/2} = 1$$

Circle γ touches α at $s = -\pi/2$, its tantrix at which point is

$$\bar{T}\left(-\frac{\pi}{2}\right) = -T(t) \sin\left(-\frac{\pi}{2}\right) + N(t) \cos\left(-\frac{\pi}{2}\right) = T(t)$$

so it is indeed tangent to $\alpha(t)$. And its curvature is

$$\kappa_\gamma(s) = \frac{\|\bar{T}'(s)\|}{\|\gamma'(s)\|} = \frac{1}{1/\kappa(t)} = \kappa(t)$$

4.5

i

When α is parameterized by arclength, $\|\alpha'\| = 1$,

$$\begin{aligned} p'(t) &= \alpha'(t) + \frac{1}{\kappa(t)} N'(t) \\ &= \alpha'(t) + \frac{1}{\kappa(t)} (-\kappa(t) T(t)) \\ &= \alpha'(t) - T(t) = \alpha'(t) - \alpha'(t) = 0 \end{aligned}$$

so $p(t)$ is a fixed point.

ii

$$\begin{aligned} \alpha(t) - p(t) &= -\frac{1}{\kappa(t)} N(t) = -\frac{1}{c} N(t) \\ \|\alpha(t) - p(t)\| &= \frac{1}{c} \end{aligned}$$

4.8

$$\begin{aligned} T &= \frac{\gamma'}{\|\gamma'\|} = \frac{\gamma'}{\langle \gamma', \gamma' \rangle^{1/2}} \\ T' &= \frac{\gamma'' \langle \gamma', \gamma' \rangle + \gamma' \langle \gamma', \gamma'' \rangle}{\|\gamma'\|^3} \end{aligned}$$

$$\begin{aligned}
\langle T', iT \rangle &= \left\langle \frac{\gamma'' \langle \gamma', \gamma' \rangle + \gamma' \langle \gamma', \gamma'' \rangle}{\|\gamma'\|^3}, \frac{i\gamma'}{\|\gamma'\|} \right\rangle \\
&= \frac{\langle \gamma'', i\gamma' \rangle \langle \gamma', \gamma' \rangle + \langle \gamma', i\gamma' \rangle \langle \gamma', \gamma'' \rangle}{\|\gamma'\|^4} \\
&= \frac{\langle \gamma'', i\gamma' \rangle \langle \gamma', \gamma' \rangle}{\|\gamma'\|^4} \\
&= \frac{\langle \gamma'', i\gamma' \rangle}{\|\gamma'\|^2}
\end{aligned}$$

Suppose $\gamma = (\gamma_1, \gamma_2)$, then

$$\gamma' = (\gamma'_1, \gamma'_2) = \gamma'_1 + i\gamma'_2, \quad i\gamma' = -\gamma'_2 + i\gamma'_1 = (-\gamma'_2, \gamma'_1)$$

$$\gamma'' = (\gamma''_1, \gamma''_2)$$

$$\langle \gamma'', i\gamma' \rangle = \gamma'_1 \gamma''_2 - \gamma'_2 \gamma''_1 = (\gamma' \times \gamma'')_z = \langle \gamma' \times \gamma'', (0, 0, 1) \rangle$$

Substitute it back into the formula:

$$\begin{aligned}
\langle T', iT \rangle &= \frac{\langle \gamma' \times \gamma'', (0, 0, 1) \rangle}{\|\gamma'\|^2} \\
\bar{\kappa} &= \frac{\langle T', iT \rangle}{\|\gamma'\|} = \frac{\langle \gamma' \times \gamma'', (0, 0, 1) \rangle}{\|\gamma'\|^3}
\end{aligned}$$

4.12

i

A unit circle oriented clockwise can be parameterized as

$$\alpha(t) = (\cos(-t), \sin(-t))$$

for $0 \leq t \leq 2\pi$.

$$\alpha'(t) = (\sin(-t), -\cos(-t))$$

$$\alpha''(t) = (-\cos(-t), -\sin(-t))$$

$$\langle \alpha' \times \alpha'', (0, 0, 1) \rangle = -\sin^2(-t) - \cos^2(-t) = -1$$

$$\bar{\kappa} = \frac{\langle \alpha' \times \alpha'', (0, 0, 1) \rangle}{\|\alpha'\|^3} = \frac{-1}{1} = -1$$

$$total \ \bar{\kappa}[\alpha] = \int_0^{2\pi} \bar{\kappa} dt = -2\pi$$

$$rot[\alpha] = \frac{total \ \bar{\kappa}[\alpha]}{2\pi} = -1$$

ii

A unit circle oriented counter-clockwise can be parameterized as

$$\alpha(t) = (\cos(t), \sin(t))$$

for $0 \leq t \leq 2\pi$.

$$\alpha'(t) = (-\sin(t), \cos(t))$$

$$\alpha''(t) = (-\cos(t), -\sin(t))$$

$$\langle \alpha' \times \alpha'', (0, 0, 1) \rangle = \sin^2(t) + \cos^2(t) = 1$$

$$\bar{\kappa} = \frac{\langle \alpha' \times \alpha'', (0, 0, 1) \rangle}{\|\alpha'\|^3} = \frac{1}{1} = 1$$

$$total \bar{\kappa}[\alpha] = \int_0^{2\pi} \bar{\kappa} dt = 2\pi$$

$$rot[\alpha] = \frac{total \bar{\kappa}[\alpha]}{2\pi} = 1$$

iii

$$\alpha(t) = (\cos t, \sin 2t)$$

$$\alpha'(t) = (-\sin t, 2 \cos 2t)$$

$$\alpha''(t) = (-\cos t, -4 \sin 2t)$$

$$\langle \alpha' \times \alpha'', (0, 0, 1) \rangle = 4 \sin t \sin 2t + 2 \cos t \cos 2t$$

$$\bar{\kappa}(t) = \frac{\langle \alpha' \times \alpha'', (0, 0, 1) \rangle}{\|\alpha'\|^3} = \frac{4 \sin t \sin 2t + 2 \cos t \cos 2t}{\sqrt{\sin^2 t + 4 \cos^2 2t}^3}$$

$$total \bar{\kappa}[\alpha] = \int_0^{2\pi} \bar{\kappa} dt = \int_0^{2\pi} \frac{4 \sin t \sin 2t + 2 \cos t \cos 2t}{\sqrt{\sin^2 t + 4 \cos^2 2t}^3} dt = 0$$

$$rot[\alpha] = \frac{total \bar{\kappa}[\alpha]}{2\pi} = 0$$