

PHYS 7125 Homework 4

Wenqi He

March 13, 2019

1

For a timelike vector t^μ at any point p it is always possible to construct an orthonormal frame $\underline{\mathbf{e}}_i^\mu$ where $\underline{\mathbf{e}}_0^\mu = t^\mu$. (Without loss of generality, t^μ can be assumed to have unit length (-1). In the general case, the results only differ by a positive factor.) In such coordinates, $t^\mu = (1, 0, 0, 0)$, the metric is locally $\eta_{\mu\nu}$, and the components of the electromagnetic energy-momentum tensor is

$$T_{\mu\nu} = \frac{1}{\mu_0} \left[F_\mu{}^\alpha F_{\nu\alpha} - \frac{1}{4} \eta_{\mu\nu} F_{\alpha\beta} F^{\alpha\beta} \right]$$

Considering t^μ is only non-zero in its time component and $T_{\mu\nu}$ is symmetric, we only need to compute

$$T_{0\nu} = \frac{1}{\mu_0} \left[F_0{}^\alpha F_{\nu\alpha} - \frac{1}{4} \eta_{0\nu} F_{\alpha\beta} F^{\alpha\beta} \right] =: A + B$$

The first term evaluates to: (Define $\mathbf{S} := \frac{1}{\mu_0} \mathbf{E} \times \mathbf{B}$)

$$A = \frac{1}{\mu_0} \begin{pmatrix} 0 & E_x/c & E_y/c & E_z/c \\ -E_x/c & 0 & -B_z & B_y \\ -E_y/c & B_z & 0 & -B_x \\ -E_z/c & -B_y & B_x & 0 \end{pmatrix} \begin{pmatrix} 0 \\ E_x/c \\ E_y/c \\ E_z/c \end{pmatrix} = \begin{pmatrix} \epsilon_0 E^2 \\ -\frac{1}{\mu_0} (E_y B_z - E_z B_y)/c \\ -\frac{1}{\mu_0} (E_z B_x - E_x B_z)/c \\ -\frac{1}{\mu_0} (E_x B_y - E_y B_x)/c \end{pmatrix} = \begin{pmatrix} \epsilon_0 E^2 \\ -S_x/c \\ -S_y/c \\ -S_z/c \end{pmatrix}$$

The second term:

$$B = -\frac{1}{4\mu_0} (-2E^2/c^2 + 2B^2) \begin{pmatrix} -1 \\ 0 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} \frac{1}{2} (-\epsilon_0 E^2 + \frac{1}{\mu_0} B^2) \\ 0 \\ 0 \\ 0 \end{pmatrix} \Rightarrow T_{0\nu} = \begin{pmatrix} \frac{1}{2} (\epsilon_0 E^2 + \frac{1}{\mu_0} B^2) \\ -S_x/c \\ -S_y/c \\ -S_z/c \end{pmatrix}$$

In matrix form, the relevant components of the energy-momentum tensor are:

$$T_{\mu\nu} = \begin{pmatrix} \frac{1}{2} (\epsilon_0 E^2 + \frac{1}{\mu_0} B^2) & -S_x/c & -S_y/c & -S_z/c \\ -S_x/c & \cdots & \cdots & \cdots \\ -S_y/c & \cdots & \cdots & \cdots \\ -S_z/c & \cdots & \cdots & \cdots \end{pmatrix}$$

(i) The weak energy condition is obviously satisfied:

$$T_{\mu\nu} t^\mu t^\nu = T_{00} = \frac{1}{2} (\epsilon_0 E^2 + \frac{1}{\mu_0} B^2) \geq 0$$

(ii)

$$T_{\mu\nu} t^\mu = T_{0\nu} = \left(\frac{1}{2} (\epsilon_0 E^2 + \frac{1}{\mu_0} B^2), -S_x/c, -S_y/c, -S_z/c \right)$$

$$T^\nu{}_\alpha t^\alpha = T^\nu{}_0 = g^{\nu\nu} \cdot T_{\nu 0} = \left(-\frac{1}{2} \left(\epsilon_0 E^2 + \frac{1}{\mu_0} B^2 \right), -S_x/c, -S_y/c, -S_z/c \right)$$

Using Lagrange's identity for cross products,

$$\begin{aligned} (T_{\mu\nu} t^\mu)(T^\nu{}_\alpha t^\alpha) &= -\frac{1}{4} \left(\epsilon_0 E^2 + \frac{1}{\mu_0} B^2 \right)^2 + \frac{\|\mathbf{S}\|^2}{c^2} \\ &= -\frac{1}{4} \left(\epsilon_0 E^2 + \frac{1}{\mu_0} B^2 \right)^2 + \frac{\epsilon_0}{\mu_0} \|\mathbf{E} \times \mathbf{B}\|^2 \\ &= -\frac{1}{4} \left(\epsilon_0 E^2 + \frac{1}{\mu_0} B^2 \right)^2 + \frac{\epsilon_0}{\mu_0} (E^2 B^2 - (\mathbf{E} \cdot \mathbf{B})^2) \\ &= -\frac{1}{4} \left(\epsilon_0 E^2 - \frac{1}{\mu_0} B^2 \right)^2 - \frac{\epsilon_0}{\mu_0} (\mathbf{E} \cdot \mathbf{B})^2 \leq 0 \end{aligned}$$

Thus the dominant energy condition is also satisfied.

2

a

The only non-vanishing components of $\Gamma_{0\nu}^\mu$ are

$$\Gamma_{0i}^0 = \frac{Mx^i}{(1-2M/r)r^3}, \quad \Gamma_{00}^i = \frac{Mx^i}{(1+2M/r)r^3} \quad (i = 1, 2, 3)$$

The geodesic equation can be simplified as

$$\begin{aligned} \frac{dp_0}{d\lambda} &= \sum_i \left(\frac{Mx^i}{(1-2M/r)r^3} p_0 p^i + \frac{Mx^i}{(1+2M/r)r^3} p^0 p_i \right) \\ &= \sum_i \left(g_{00} \frac{Mx^i}{(1-2M/r)r^3} + g_{ii} \frac{Mx^i}{(1+2M/r)r^3} \right) p^0 p^i \\ &= \sum_i \left(-\frac{Mx^i}{r^3} + \frac{Mx^i}{r^3} \right) p^0 p^i = 0 \end{aligned}$$

b

No. $p^0 = g^{0\nu} p_\nu = g^{00} p_0 = -(1-2M/r)^{-1} p_0$, which is not constant unless r is constant.

c

For the atom at rest on the surface of the sun, $dx^i = 0$, $r = R$,

$$d\tau^2 = -ds^2 = (1-2M/R)dt^2 \quad \Rightarrow \quad u^0 = \frac{dt}{d\tau} = \frac{1}{\sqrt{1-2M/R}}$$

d

For both the atom and the distant observer, $dx^i = 0$,

$$\begin{aligned} d\tau^2 &= -ds^2 = (1-2M/r)dt^2 \\ \Rightarrow \quad u^\mu &= \frac{dx^\mu}{d\tau} = (dt/d\tau, 0, 0, 0) = ((1-2M/r)^{-1/2}, 0, 0, 0) \end{aligned}$$

The photon energy observed at both locations can be expressed as

$$E = g_{\mu\nu} p^\mu u^\nu = (1 - 2M/r)^{-1/2} g_{\mu\nu} p^\mu K^\nu$$

where $K^\nu = (1, 0, 0, 0)$ is a Killing vector as the metric has no time dependence; therefore $g_{\mu\nu} p^\mu K^\nu$ is conserved along the photon's world line, which is a geodesic as stated in the problem. Then,

$$\frac{\lambda_r}{\lambda_e} = \frac{hc/\lambda_e}{hc/\lambda_r} = \frac{E_e}{E_r} = \frac{(1 - 2M/R)^{-1/2}}{\lim_{r \rightarrow \infty} (1 - 2M/r)^{-1/2}} = 1 + \frac{M}{R} + \mathcal{O}\left(\frac{M^2}{R^2}\right)$$

$$z = \frac{\lambda_r - \lambda_e}{\lambda_e} = \frac{\lambda_r}{\lambda_e} - 1 = \frac{M}{R} + \mathcal{O}\left(\frac{M^2}{R^2}\right)$$