

Equilibrium chemistry down to 100 K

Impact of silicates and phyllosilicates on carbon/oxygen ratio

P. Woitke^{1,2}, Ch. Helling^{1,2,5}, G. H. Hunter^{1,2}, J. D. Millard^{1,2}, G. E. Turner^{1,2}, M. Worters^{1,2}, J. Blečić³, J. W. Stock⁴

¹ SUPA School of Physics & Astronomy, University of St Andrews, North Haugh, KY16 9SS, St Andrews, UK

² Centre for Exoplanet Science, University of St Andrews, St Andrews, UK

³ New York University Abu Dhabi, Abu Dhabi, United Arab Emirates

⁴ Department of Chemistry and Environmental Science, Medgar Evers College-City University of New York, 1650 Bedford Avenue, Brooklyn, NY 11235, US

⁵ Anton Pannekoek Institute for Astronomy, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, NL

Received Oct. 27th, 2017; accepted Dec. 2nd, 2017

ABSTRACT

We introduce a fast and versatile computer code, GGchem, to determine the chemical composition of gases in thermo-chemical equilibrium down to 100 K, with or without equilibrium condensation. We review the data for molecular equilibrium constants, $k_p(T)$, from several sources and discuss which functional fits are most suitable for low temperatures. We benchmark our results against another chemical equilibrium code. We collect Gibbs free energies, ΔG_f° , for about 200 solid and liquid species from the NIST-JANAF database and the geophysical database SUPCRTBL. We discuss the condensation sequence of the elements with solar abundances in phase equilibrium down to 100 K. Once the major magnesium silicates $\text{Mg}_2\text{SiO}_4[\text{s}]$ and $\text{MgSiO}_3[\text{s}]$ have formed, the dust/gas mass ratio jumps to a value of about 0.0045 which is significantly lower than the often assumed value of 0.01. Silicate condensation is found to increase the carbon/oxygen ratio (C/O) in the gas from its solar value of ~ 0.55 up to ~ 0.71 , and, by the additional intake of water and hydroxyl into the solid matrix, the formation of phyllosilicates at temperatures below ~ 400 K increases the gaseous C/O further to about 0.83. Metallic tungsten (W) is the first condensate found to become thermodynamically stable around 1600 – 2200 K (depending on pressure), several hundreds of Kelvin before subsequent materials like zirconium dioxide (ZrO_2) or corundum (Al_2O_3) can condense. We briefly discuss whether tungsten, despite its low abundance of $\sim 2 \times 10^{-7}$ times the silicon abundance, could provide the first seed particles for astrophysical dust formation. The GGchem code is publicly available at <https://github.com/pw31/GGchem>.

Key words. astrochemistry – planets and satellites: atmospheres – planets and satellites: composition – stars: winds, outflows – molecular data – methods: numerical

1. Introduction

Deriving the chemical composition of gases in the atmospheres of stars, brown dwarfs and planets is one of the most fundamental modelling tasks in astronomy. All hydrodynamical, evolutionary and radiative processes depend on the thermodynamics of the gas. In particular, the interpretation of spectral observations requires detailed knowledge about the gas composition. Only then, the inverse problem of inferring the internal physico-chemical structure of an object in space from spectroscopic observations can be successfully solved.

Instrument development has progressed tremendously over the past decades, allowing us to use a large range of wavelengths to analyse e.g. the chemical composition of brown dwarfs (see Allard et al. 1997; Helling & Casewell 2014 and references in these reviews) and exoplanets (e.g. Désert et al. 2008; Grillmair et al. 2008; de Kok et al. 2013; Kreidberg et al. 2014; Fraine et al. 2014; Birkby et al. 2017). We can also revisit the long-standing question of what drives the mass loss of oxygen-rich Asymptotic Giant Branch (AGB) stars (Gail & Sedlmayr 1998; Woitke 2006; Höfner & Andersen 2007; Golriz et al. 2014; Gail et al. 2016; Decin et al. 2017), including the problem of seed particle formation and the search for the first condensate in space (see e.g. Patzer et al. 1995; Patzer 2007). High-resolution spec-

troscopy of molecular lines from CO and H_2O are also used, for example, to derive winds properties of extrasolar planets (Brogi et al. 2016). Infrared instruments on board of the JAMES-WEBB SPACE TELESCOPE are expected to revolutionise our understanding of cool objects like exoplanets, brown dwarfs and cool stars (e.g. Marley & Leggett 2009; Beichman et al. 2014). High-resolution ground-based spectroscopy with e.g. CARMENES (e.g. Sanchez-Lopez et al. 2017) aim at a better understanding of late-type main-sequence stars and their low-mass planets. In the more distant future, PLATO (Rauer et al. 2016) will combine astroseismology for solar-like stars with the search for habitable, Earth-like planets. Exploring such high-end quality data requires substantial modelling efforts of which deriving the chemical gas composition is one of the key issues, which needs to be fast and sufficiently accurate over a wide range of temperatures and pressures. The predictions of the molecular composition of the observed gases depends critically on the availability and precision of thermo-chemical data.

In the first approximation, the gas is assumed to be in Local Thermodynamical Equilibrium (LTE), so the gas composition can be calculated from the thermodynamical principle of minimisation of the system Gibbs free energy (e.g. White et al. 1958; Eriksson 1971). Thermo-chemical equilibrium is a part of the set of LTE assumptions used in particular to compute

the abundances of molecules. Pioneering work was done here for the atmospheres of cool stars where molecules are the main opacity carriers (e.g. [Tsuji 1965](#); [Auer & Mihalas 1968](#); [Gustafsson 1971](#)). To date, applications of thermo-chemical equilibrium are widespread in astronomy, for example in gas giant planet atmospheres, brown dwarfs and low-mass dwarf stars ([Allard & Hauschildt 1995](#); [Tsuji et al. 1996](#); [Lodders & Fegley 2002](#); [Visscher et al. 2010](#); [Marley & Robinson 2015](#)).

Deviations from thermo-chemical equilibrium are generally expected to occur in diluted gases, in particular due to UV-fields, X-rays or cosmic rays as, for example, in protoplanetary discs (e.g. [Semenov & Wiebe 2011](#); [Woitke et al. 2016](#)), in the upper layers of planetary atmospheres (e.g. [Moses 2014](#); [Rimmer & Helling 2016](#)), or in case of fast hydrodynamical processes where the dynamical timescale becomes smaller than the chemical timescale, for example in interstellar shocks (e.g. [Hollenbach & McKee 1989](#)).

[Griffith & Yelle \(1999\)](#) and [Cooper & Showman \(2006\)](#) argue for strong chemical dis-equilibrium effects and higher CO abundances in BD and exoplanet atmospheres. Concerning the atmospheres of cool giant planets, [Moses \(2014\)](#) have shown that thermo-chemical equilibrium only prevails in the deepest layers ($p > \text{a few bar}$), whereas transport-induced quenching and photochemistry are likely to be important in the upper layers. However, recent kinetic chemical models ([Visscher et al. 2006, 2010](#); [Zahnle et al. 2009](#); [Line et al. 2010](#); [Moses et al. 2011](#); [Kopparapu et al. 2012](#); [Venot et al. 2012](#)) have shown that in hot exoplanet atmospheres ($T \gtrsim 1200 \text{ K}$), the chemical timescales are in fact quite short, and hence thermo-chemical equilibrium prevails. For the hottest planets, [Line & Yung \(2013\)](#) and [Oreshenko et al. \(2017\)](#) have shown that at a pressure of 100 mbar CH_4 , CO , H_2O and H_2 should be close to chemical equilibrium even when fast atmospheric mixing is assumed. [Line & Yung \(2013\)](#) have pointed out that spectral observations probe layers in exoplanet atmospheres where quenching by atmospheric mixing can be relevant, but not photochemistry.

At low temperatures ($\lesssim 2000 \text{ K}$), the formation of solids and liquids becomes an essential part of the problem of finding the equilibrium composition. The condensates selectively consume most of certain elements (e.g. Al, Si, Mg, Fe), whereas others remain in the gas phase down to substantially lower temperatures (e.g. S, Cl). The abundant elements H, He, C, N, O remain mostly gaseous before even they, eventually, will condense e.g. in form of water, ammonia and methane ices like on Titan. The remaining gas phase element abundances can differ by orders of magnitude from the initial, undepleted (e.g. solar) abundances before condensation (e.g. [Woitke & Helling 2004](#); [Juncher et al. 2017](#)), which is essential to understand the spectroscopic appearance of cool and cold objects like AGB stars and their winds ([Gail et al. 1984](#); [Sedlmayr & Dominik 1995](#); [Helling et al. 1996](#); [Woitke 2001](#); [Jeong et al. 2003](#)), or the formation of clouds in brown dwarfs and planetary atmospheres ([Lunine et al. 1986](#); [Tsuji et al. 1996](#); [Marley et al. 1999](#); [Allard et al. 2001](#); [Burrows et al. 2002](#); [Tsuji 2005](#); [Marley et al. 2002](#); [Helling et al. 2008](#)).

Since dust grains can easily de-couple from the gas phase (for example rain-out in planetary atmospheres, gravitational settling in protoplanetary discs, acceleration by radiation pressure in AGB star winds), the condensed elements may be carried away, gather in other places (like oceans) or re-evaporate in other places (e.g. comet tails). These effects are very likely to cause most peculiar local element abundances, with large effects on gas phase abundances and spectroscopic appearance. Such effects are rarely studied in detail (see, however, [Booth et al. 2017](#)). Instead, researchers tend to rely on known sets of element abun-

dances, in particular the solar element abundances (e.g. [Asplund et al. 2009](#)) and those derived from Earth rock analysis.

Deviations of element abundances from these known sets have mostly been treated by reducing all metal abundances by the same factor, i.e. by changing the metallicity $[\text{M}/\text{H}]$, (e.g. [Helling & Lucas 2009](#); [Witte et al. 2009](#); [Crossfield et al. 2013](#); [Hu & Seager 2014](#); [Morley et al. 2017](#)) or by changing the carbon-to-oxygen ratio (e.g. [Madhusudhan et al. 2011](#); [Madhusudhan 2012](#); [Helling et al. 2017](#); [Van Eck et al. 2017](#)). [Gaidos \(2000\)](#) was one of the earliest works suggesting the importance of C/O in an exoplanet context. [Mollière et al. \(2015\)](#) have studied the impact of C/O on the (p, T) -structure and emergent spectra of irradiated planets. Planetary atmospheres with very different element abundances have been studied by ([Mbarek & Kempton 2016](#); [Mahapatra et al. 2017](#)), considering an atmosphere made of evaporated rock.

In order to derive the composition of a gas in thermo-chemical equilibrium, different approaches are used by different groups working on stellar, brown dwarf and planetary atmospheres. Three major schools are: (i) use of law of mass action and molecular equilibrium constants based on Gibbs free energy data ([Tsuji 1973](#); [Stock 2008](#); [Bilger et al. 2013](#)), (ii) use of law of mass action and molecular equilibrium constants based on partition functions ([Allard & Hauschildt 1995](#); [Gustafsson et al. 2008](#); [Barklem & Collet 2016](#); [Van Eck et al. 2017](#)), and (iii) minimisation of the system Gibbs free energy ([Lodders 2003](#); [Ackerman & Marley 2001](#); [Miller-Ricci et al. 2009](#); [Mbarek & Kempton 2016](#); [Blecic et al. 2016](#)). [Helling et al. \(2008\)](#) provide a summary of approaches used in brown dwarf and planetary atmospheres.

Approaches (i) and (ii) usually lead to a system of N algebraic equations for N unknowns, which can be solved by any root-finding algorithms, for example by the Newton-Raphson method. In contrast, the Gibbs free energy minimisation technique (iii) requires special numerical minimisation algorithms ([Spang 1962](#)), for example the Dantzig-simplex method, the Hessian-conjugate gradient method, or the Lagrangian steepest-descent method ([Blecic et al. 2016](#)). The Gibbs free energy minimisation approach can easily be generalised to include equilibrium condensation, which is the reason why this approach was introduced originally. However, finding the root of a coupled non-linear equation system $\mathbf{F}(\mathbf{x}) = 0$ can be done numerically in much more efficient ways than finding the minimum of $F = \sum F_i^2(\mathbf{x}) \rightarrow \min$, because the vector \mathbf{F} has much more information than the scalar F . Therefore, Gibbs free energy minimisation codes tend to be considerably slower.

Beyond thermo-chemical equilibrium models for gas phase and condensates, hybrid methods have been developed where the condensation is treated time-dependently, but the concentrations of gaseous molecules are calculated in thermo-chemical equilibrium (e.g. [Gail et al. 1984](#); [Sedlmayr & Dominik 1995](#); [Woitke & Helling 2004](#); [Woitke 2006](#); [Höfner & Andersen 2007](#); [Höfner et al. 2016](#); [Helling & Fomins 2013](#)), following the idea that dust formation is by far the slowest process, causing the first deviations from LTE.

The probably most common approach used in the retrieval community, however, is not to solve any equations at all, but to treat the molecular concentrations as free parameters which are then adjusted to match observations (see e.g. Table 1 in [Helling et al. 2008](#)). This is partly done for reasons of feasibility, as the retrieval technique requires to run tens of thousands of models, but also to allow the method to retrieve non-equilibrium effects whereas enforcing equilibrium would rather be considered as a limitation. [Benneke \(2015\)](#), [Line et al. \(2016\)](#) and [Oreshenko](#)

et al. (2017) have shown that it is feasible to include equilibrium chemistry in exoplanet retrieval methods.

This paper summarises in Sect. 2 the theoretical background of our approach to compute the abundances of molecules and condensates in thermo-chemical equilibrium. Section 3 gives an overview of the thermo-chemical data available for molecules and condensates. We assess the level of (dis-)agreement between various data sources. Section 4 introduces our thermo-chemical equilibrium code. Section 5 presents our results concerning the spectroscopically most active molecules, a benchmark test, and the condensation sequence of elements. We demonstrate that equilibrium condensation leads to a significant increase of the C/O ratio in the gas phase where phyllosilicates play a particular role. Section 6 re-addresses the question of the first condensate in space, which is of particular interest for dust formation in AGB star winds. Section 7 contains our summary and conclusions.

2. Chemical and phase equilibrium

2.1. Molecular equilibrium

Let us consider a molecule $A_aB_bC_c$ made of three elements A, B, C, where a, b, c are the stoichiometric factors. Guldberg's law of mass action (e.g. Berline & Bricker 1969) is given by

$$\frac{p_{A_aB_bC_c}}{p^\circ} = \left(\frac{p_A}{p^\circ}\right)^a \left(\frac{p_B}{p^\circ}\right)^b \left(\frac{p_C}{p^\circ}\right)^c \exp\left(-\frac{\Delta G_f^\circ}{RT}\right), \quad (1)$$

where the $p_i = n_i kT$ are the partial pressures [dyn/cm²], n_i the particle densities [cm⁻³] and p° is a standard pressure. ΔG_f° is the Gibbs free energy of formation [J/mol] of the molecule at standard pressure from neutral atoms at the same temperature

$$\Delta G_f^\circ = G^\circ(A_aB_bC_c, T) - a G^\circ(A, T) - b G^\circ(B, T) - c G^\circ(C, T). \quad (2)$$

The equilibrium constants k_p are introduced as

$$p_{A_aB_bC_c} = k_p(A_aB_bC_c, T) p_A^a p_B^b p_C^c, \quad (3)$$

where

$$k_p(A_aB_bC_c, T) = (p^\circ)^{1-a-b-c} \exp\left(-\frac{\Delta G_f^\circ}{RT}\right). \quad (4)$$

As we are using cgs-units in this paper, the k_p have units (dyn/cm²)^{1-a-b-c}. T is the gas temperature [K], R the ideal gas constant [J/mol/K] and k the Boltzmann constant [erg/K].

To determine the chemical composition of the gas, we solve the element and charge conservation equations as

$$\epsilon_k n_{(H)} = \sum_i s_{i,k} n_i, \quad (5)$$

where ϵ_k are the element abundances normalised to hydrogen ($\epsilon_H = 1$), $n_{(H)}$ is the total hydrogen nuclei density $n_{(H)} = \sum_i s_{i,H} n_i$. n_i denote all gas particle densities including free electrons, neutral and charged atoms, and neutral and charged molecules. $s_{i,k}$ is the stoichiometric factor of element k in gas particle i . We assume charge neutrality by the inclusion of the charge as an additional element 'el' with zero abundance ($\epsilon_{el} = 0$) where $s_{i,el} = 0$ for neutrals, $s_{i,el} = +1$ for ions and $s_{i,el} = -1$ for cations. The free electron has $s_{el,el} = -1$. The gas density ρ [g/cm³] is given by

$$\rho = \sum_i m_i n_i = n_{(H)} \sum_k m_k \epsilon_k, \quad (6)$$

where the second part of Eq. (6) follows from Eq. (5), and the total gas pressure is given by

$$p = \sum_i n_i kT = n(\rho, T) kT = \frac{\rho kT}{\mu(\rho, T)}. \quad (7)$$

m_i are the gas particle masses and m_k are the masses of the elements. The total particle density $n = \sum_i n_i$ and the mean molecular weight $\mu = \sum_i m_i n_i / n$ are results of the computations and hence depend on density and temperature. The total hydrogen nuclei particle density $n_{(H)}$ is always proportional to the mass density ρ , whereas n is not, causing the gas to deviate from an ideal gas.

After elimination of all molecular particle densities from Eq. (5) by using the $k_p(T)$ -data (see Eq. 3), Eq. (5) becomes a system of non-linear algebraic equations with K unknowns, namely the atomic partial pressures and the electron partial pressure, for example $\{p_k | k = H, \dots, W, el\}$ where H is hydrogen and W is tungsten. All results depend solely on $n_{(H)}$, T and ϵ_k .

If a solution is requested for a given pressure and temperature, an iteration is performed where the mean molecular weight μ is initially guessed, Eqs. (7) and (6) are used to compute $n_{(H)}$ from p and T , the equilibrium chemistry is solved with $n_{(H)}$ and T , and finally μ is re-calculated. We find this iteration to converge after 1-5 calls of the equilibrium chemistry.

For high temperatures, $T \gtrsim 1000$ K, the solution is easy to find numerically as all particle concentrations stay within 10^{-a} a few ten. However, for $T \rightarrow 500$ K, the results become increasingly more extreme, for example the electron concentration approaches 10^{-30} , and for $T \rightarrow 100$ K the problem turns into a numerical nightmare with some $k_p > 10^{+several\ 1000}$ and some atom and electron concentrations $< 10^{-several\ hundred}$. We overcome these problems by switching to quadruple precision arithmetics at low temperatures and by applying an iterative procedure according to the hierarchical order of elements to provide extremely good initial guesses of the unknowns for the final Newton-Raphson iteration. We explain the details of the numerical approach of the new GGChem-code in App. A.

2.2. Condensed phases

The supersaturation ratio of a condensate j is given by

$$S_j = \frac{p_j}{p_j^{\text{vap}}(T)}, \quad (8)$$

where $p_j^{\text{vap}}(T)$ is the vapour pressure

$$p_j^{\text{vap}}(T) = p^\circ \exp\left(\frac{G^\circ(j[\text{cond}], T) - G^\circ(j, T)}{RT}\right). \quad (9)$$

In phase equilibrium we have

$$S_j \begin{cases} < 1 & \text{condensate is unstable and not present,} \\ = 1 & \text{condensate is stable and present,} \end{cases} \quad (10)$$

whereas $S_j > 1$ (supersaturation) is not possible in phase equilibrium. Equation (10) states that the partial pressure of any molecule is limited in phase equilibrium, if a corresponding condensed phase exists. Once p_j exceeds that threshold (its vapour pressure), that molecule condenses¹ and leaves the gas phase,

¹ The actual process of condensation is not considered in the frame of chemical and phase equilibrium.

which reduces a few gas phase element abundances ϵ_k . This reduction causes not only p_j to drop, but also affects on all other partial pressures in the gas (usually reduces them). This process continues until phase equilibrium is established, see appendix B in (Helling et al. 2008).

For condensates which have no corresponding molecule in the gas phase (which is true for most minerals), Eq. (8) is not applicable. However, we can consider the fictive nominal molecule of that condensate, for example $j = A_a B_b C_c$ with unknown $G^\circ(A_a B_b C_c, T)$, and apply Eqs. (8) and (9). Using also Eqs. (3) and (4), we find the generalised supersaturation ratio to be

$$S_{A_a B_b C_c} = \left(\frac{p_A}{p^\circ}\right)^a \left(\frac{p_B}{p^\circ}\right)^b \left(\frac{p_C}{p^\circ}\right)^c \exp\left(-\frac{\Delta G_f^\circ}{RT}\right) \quad (11)$$

where

$$\Delta G_f^\circ = G^\circ(A_a B_b C_c[\text{cond}], T) - a G^\circ(A, T) - b G^\circ(B, T) - c G^\circ(C, T) \quad (12)$$

is the Gibbs free energy of formation of the condensed phase $A_a B_b C_c$ at standard pressure from atoms in the gas phase at temperature T . The unknown Gibbs free energy $G^\circ(A_a B_b C_c, T)$ of the fictive molecule cancels out.

The addition of Eq. (10) to chemical equilibrium models (\rightarrow *equilibrium condensation* or *phase equilibrium* models) leads to a considerable complication of the mathematical structure of the problem to solve. The case differentiation in Eq. (10) means that we do not know a-priori which condensates to consider as only those which finally result to be present contribute to the number of equations to be solved (in form of $S_j = 1$), whereas the supersaturation ratios of all other condensates do not matter as long as they stay < 1 . Clearly, at very high temperatures, where no condensates are stable at all, the problem falls back to the pure gas phase equilibrium discussed in the previous section.

Our numerical method in GGChem is based on a Newton-Raphson iteration with nested calls of the equilibrium chemistry with reduced gas phase element abundances ϵ_k due to condensation, see App. B. This reduction becomes extremely large at low temperatures – up to hundreds of orders of magnitudes. The additional unknowns to be solved for are the reduced gas phase element abundances as affected by the condensation of the selected solid and liquid phases. This selection may change during the iteration. Albeit being somewhat sophisticated, we find that this method is fast and very accurate down to 100 K when using quadruple precision arithmetics.

3. Thermo-chemical data

3.1. Molecular equilibrium constants

The k_p -data of the various ions and molecules must usually be fitted as function of temperature in some way, before they can be used in models. An alternative approach was recently proposed by Blečić et al. (2016), who use all available NIST-JANAF ΔG_f° data-points directly by means of internal spline-fits. Various fit-functions have been proposed as summarised below. From Eq. (4) we have the general relation between ΔG_f° and k_p as

$$\ln k_p = (1 - n) \ln p^\circ - \frac{\Delta G_f^\circ}{RT} \quad (13)$$

where $n = a + b + c$ is the sum of stoichiometric coefficients in the molecule, for example $n = 3$ for H_2^- and $n = 1$ for H_2^+ .

Tsuji (1973) has used the following fit-function

$$\log_{10} k_p^{\text{Tsu}}(T) = -a_0 - a_1 \theta - a_2 \theta^2 - a_3 \theta^3 - a_4 \theta^4 \quad (14)$$

for 335 molecules, where $\theta = 5040/T$ and a_i are the fit coefficients. All k_p formulae listed here are valid in cgs units.

Gail & Sedlmayr (1986) have used

$$\ln k_p^{\text{Gail}}(T) = a_0 + a_1 \theta + a_2 \theta^2 + a_3 \theta^3 + a_4 \theta^4 \quad (15)$$

which is the functional form that was used in the old GGChem code until recently (e.g. Helling et al. 2006; Bilger et al. 2013; Helling et al. 2017). The old data collection had 205 molecules.

Sharp & Huebner (1990) have used

$$\ln k_p^{\text{S\&H}} = (1 - n) \ln p^\circ - \frac{a_0/T + a_1 + a_2 T + a_3 T^2 + a_4 T^3}{R_{\text{cal}} T}, \quad (16)$$

fitting directly the ΔG_f° [cal/mol] of 184 molecules. Here, the standard pressure is $p^\circ = 1$ atm and $R_{\text{cal}} = 1.987$ cal/mol/K.

A new approach was presented by Stock (2008)

$$\ln k_p^{\text{St}} = (1 - n) \ln p^\circ + \left(\frac{a_0}{T} + a_1 \ln T + a_2 + a_3 T + a_4 T^2\right) \quad (17)$$

who fitted the dimensionless quantity $-\Delta G_f^\circ/(RT)$ for 924 molecules, to be used with $p^\circ = 1$ bar. This functional form has a very smooth and stable behaviour towards low temperatures. It also gives automatically more weight to the data points at low temperatures, so we have adopted this functional form and most of their data in the new GGChem code.

Barklem & Collet (2016) have recently published partition functions for 291 diatomic, and charged diatomic molecules, from which k_p is derived, but without providing a fit formula. Note that their data is in SI units, and the definition of their k_p is different from ours, in particular for the charged molecules. We have done all necessary conversions (see App. C) and fitted the Barklem & Collet data with a Stock-function.

Figure 1 compares the fitted data for two example molecules found in all 5 datasets. The deviations are as large as 10 kJ/mol for HS and about 5 kJ/mol for CN. At $T = 300$ K, an uncertainty of ± 6 kJ/mol (0.06 eV) translates into an uncertainty in k_p of about one order of magnitude, same for ± 60 kJ/mol (0.6 eV) at $T = 3000$ K. The complete comparison catalogue (Worters et al. 2017) includes 2782 individual datasets for 1155 molecules, where similar plots as shown in Figure 1 can be found for all molecules for which we found at least two different data sources. We observe astonishingly large deviations in some cases. The reasons for these deviations are not entirely clear, but some factors could be

- different primary data sources,
- extrapolation errors,
- the “art of fitting”, and human errors.

Gail & Sedlmayr (1986), Sharp & Huebner (1990) and Stock (2008) have all based their fits on the JANAF database, but using different editions. Gail & Sedlmayr have used the 2^{ed} edition (Stull & Prophet 1971), Sharp & Huebner have used the 3rd edition (Chase et al. 1982; Chase 1986), and Stock has used the 4th edition (Chase 1998). All three works include some Tsuji-data for molecules missing in JANAF. Stock (2008) has refitted the Tsuji-data for $T > 1000$ K with his fit-function that is more reliable concerning extrapolation towards lower temperatures. Despite using more or less the same original data sources, some molecules show considerable deviations even at medium temperatures (see Worters et al. 2017). We conclude that errors of about




Fig. 1. Comparison of $k_p(T)$ data for molecules HS (mercapto) and CN (cyanogen) used by several groups. The second column of smaller plots shows the deviations of $\log_{10} k_p$ from the mean values (computed without the Tsuji data). On the right side, the data has been converted to ΔG_f° [kJ/mol] (see Eq. 13), before computing the deviations from the mean values. The large deviations at small temperatures for the (Tsuji 1973) and (Sharp & Huebner 1990) data are due to extrapolation errors beyond the valid fit-range. HS is classified as belonging to the group “data disagrees”, and CN to “data disagrees at low temperatures” in our comparison catalogue (Worters et al. 2017) since even without the Tsuji data, there are relatively large differences to the new (Barklem & Collet 2016) data.

half an order of magnitude for k_p can easily occur, especially at low temperatures, depending on which fit-formula is chosen and how the fit was obtained in detail. Tsuji (1973) and Barklem & Collet (2016) have used other, independent data sources. Some obvious differences occur in particular when different molecular dissociation energies are assumed. We also find a few outliers, where one data source is off by many orders of magnitude with respect to all others. Although we cannot exclude human errors on our side, our comparisons have been made in a highly automated way, so these outliers are quite puzzling.

Tsuji (1973) and Sharp & Huebner (1990) recommend to use their fits only for $T \gtrsim 1000$ K, where the Sharp & Huebner fit-function behaves somewhat more robustly in extrapolations towards low temperatures. This is because $\Delta G_f^\circ(T)$ is an approximately linear function of T as shown in Fig. 1. The fit term $\Delta G_f^\circ \sim 1/T$ causes the extrapolation problems at low T which is avoided in the (Stock 2008)-fits, which are therefore much more reliable for extrapolations towards low T . The old GGchem fits (Gail & Sedlmayr 1986) should not be applied for $T \lesssim 500$ K, and behave in a similar way as the Tsuji fits, namely with strong extrapolation errors toward low temperatures, because of a $\Delta G_f^\circ \sim 1/T^3$ term.

Appendix C highlights a few more details of this comparison and Table D.1 lists our selected set of $k_p(T)$ functions for 568 molecules composed of 24 elements, which can be safely applied to temperatures 100 – 6000 K. We could use more elements and molecules, but the data listed in Table D.1 have been carefully

checked and benchmarked against the TEA code (Blecic et al. 2016), see Sect. 5.1.

3.2. Condensed phase thermo-chemical data

We have used two main sources for thermo-chemical data of condensed phases in this paper, the NIST-JANAF database (Chase et al. 1982; Chase 1986)², and the geophysical SUPCRTBL database (Zimmer et al. 2016; Johnson et al. 1992). When extracting Gibbs free energy data $\Delta G_f^\circ(T)$ and/or vapour pressure data $p^{\text{vap}}(T)$ from these sources, it is important to understand the reference states of the elements with respect to which the data is presented, see more details in App. D. NIST-JANAF uses temperature-dependent reference states. When subtracting the $\Delta G_{f,\text{NIST}}^\circ$ data, the reference state Gibbs free energies cancel out, so we can use

$$\Delta G_f^\circ = \Delta G_{f,\text{NIST}}^\circ(A_a B_b C_c[\text{cond}], T) \quad (18)$$

$$-a \Delta G_{f,\text{NIST}}^\circ(A, T) - b \Delta G_{f,\text{NIST}}^\circ(B, T) - c \Delta G_{f,\text{NIST}}^\circ(C, T),$$

or, respectively,

$$p_{A_a B_b C_c}^{\text{vap}}(T) = p^\circ \exp\left(\frac{1}{RT} \left[\Delta G_{f,\text{NIST}}^\circ(A_a B_b C_c[\text{cond}], T) - \Delta G_{f,\text{NIST}}^\circ(A_a B_b C_c, T) \right] \right) \quad (19)$$

with $p^\circ = 1$ bar. The $\Delta G_{f,\text{NIST}}^\circ(T)$ data is available in the 7th column of the NIST-JANAF files at temperature points mostly sepa-

² <http://kinetics.nist.gov/janaf>




Fig. 2. Comparison of condensed phase Gibbs free energy data from (Sharp & Huebner 1990, orange), fits to the NIST-JANAF database done by the authors of this paper for GGchem (green), and fits to the SUPCRTBL data (Zimmer et al. 2016, blue). The SUPCRTBL data points have been generated for temperatures 100 K to 2500 K according to the equations and parameters given in Zimmer et al. (black dashed curve). The lower plot shows the deviations [kJ/mol] with respect to the mean of all three data sources.

rated by 100 K. The Gibbs free energy data can be directly compared to Sharp & Huebner (1990), and the vapour pressure data can be compared to e.g. Yaws (1999), Weast (1971) and Ackerman & Marley (2001).

The geophysical SUPCRTBL database provides condensed phase Gibbs free energy data $\Delta G_{f,SU}^{\circ}(T, p)$ via formulas with tabulated coefficients which partly have a direct physical meaning. In this database, the Gibbs free energy of formation is defined in a different way, namely with respect to the standard states of the elements. These standard states are the most stable forms of the pure elements at standard pressure $p^{\circ} = 1$ bar and room temperature $T_{\text{ref}} = 298.15$ K. These are atomic gases for the noble gases He, Ne, Ar, ...; diatomic gases $\frac{1}{2}\text{H}_2$, $\frac{1}{2}\text{N}_2$, $\frac{1}{2}\text{O}_2$, $\frac{1}{2}\text{F}_2$, $\frac{1}{2}\text{Cl}_2$ for H, N, O, F, Cl; liquids for Br and Hg; and crystalline solids for all other elements. In order to use these data for our purpose, we need to subtract the Gibbs free energy of formation of the free atoms at temperature T from the standard states at reference temperature T_{ref} . The respective thermo-chemical data of the atoms are not available in SUPCRTBL, however we can use the NIST-JANAF database to provide this link, see further details in App. D. To fit the resulting $\Delta G_f^{\circ}(T)$ functions, we have generated 100 ΔG_f° data points at log-equidistant temperature points between 100 K and 2500 K.

While fitting the data points from either the NIST-JANAF or the SUPCRTBL database by smooth functions, we have tried various options and carefully selected those fits which perform best concerning precision and possible interpolation and extrapolation issues, see App. D. If the corresponding molecule of a

condensate is known to exist as a free molecule in the gas, it is generally advantageous to use $p^{\text{vap}}(T)$ fits, because the values to be fitted only correspond to a few 100 kJ/mol. However, if no such molecule exists (which is true for most minerals) we need to fit the ΔG_f° directly, with values of several 1000 kJ/mol, where the relative precision of the fits becomes crucial. A particular challenge is to properly fit pairs of solid and liquid phase data, which are quite similar to each other, yet their intersection point should match the melting point, see Fig. D.1. Table D.2 lists our fit formula and fit coefficients for 97 condensates, mostly fitted by us to the NIST-JANAF database, with a few additions from other sources. The fits done by ourselves generally have precisions < 1 kJ/mol, much better in most cases. The melting points are matched to within < 20 K (or $< 1\%$), much better in most cases, see D.3. Table D.4 lists the ΔG_f° data for 160 condensates extracted from the SUPCRTBL database.

Figure 2 shows a comparison between ΔG_f° -fits for MgSiO_3 (enstatite) obtained from three sources. The reader is welcome to study an auxiliary document (Woitke et al. 2017) which contains additional figures like Fig. 2 for all 121 condensates extracted from the SUPCRTBL database. The different data generally agree well with each other, within about 5 kJ/mol, although there are a few exceptions, remarkably $\text{CaAl}_2\text{Si}_2\text{O}_8$ (anorthite) which seems off by about 100 kJ/mol in (Sharp & Huebner 1990). The Sharp & Huebner fits can safely be extrapolated to about 500 K, but may have extrapolation artefacts below. All new fits to the NIST-JANAF data can be used from about 3500 K down to 100 K. This does require some extrapolations for a small number of condensates where no such NIST-JANAF data exist, but we have carefully checked that our fits continue smoothly and at least do not produce any artefacts. The behaviour of the various fit functions is similar to the molecular k_p -fits as described in Sect. 3.1, the Stock-fit function is most robust towards low temperatures, but slightly less accurate at high temperatures.

4. The GGchem code

GGchem is an abbreviation for the German word “Gleich-Gewichts-Chemie” which means equilibrium chemistry. The code has been originally developed by H.-P. Gail in the 1970s (e.g. Gail & Sedlmayr 1986). Further works on the code have been done by C. Dominik, Ch. Helling and P. Woitke in Berlin until 2005. For this paper, we have completely re-written the code and given it a modern FORTRAN-90 code architecture. We have updated the thermo-chemical input data (Sect. 3.1), and have added a quadruple precision version and a new pre-iteration scheme (App. A) to improve code stability and range of applicability down to 100 K. We have added the equilibrium condensation part (App. B), and have added and revised the condensed phase thermo-chemical data (Sect. 3.2). The code is publicly available including all thermo-chemical data³.

GGchem allows the user to select elements, molecules and condensates via versatile input files. All elements from hydrogen to zirconium (atomic number 40) are supported, with additional options to include tungsten (atomic number 74) and charges. The element abundances are currently pre-compiled with options for solar abundances (Asplund et al. 2009), meteorite abundances or Earth crust abundances. The default choice is to select all molecules and condensates which are composed of the selected elements from provided data-files, but the user is welcome to develop their own input files, e.g. a customised small model optimised for speed, with just a few elements and hand-selected

³ <https://github.com/pw31/GGchem>




Fig. 3. Benchmark test against TEA (Blecic et al. 2016) at constant $p = 1$ bar and $T = 100 \dots 6000$ K, showing mixing ratios n_i/n . The coloured full and dashed curves are the GGchem results. The TEA results are overplotted with narrow grey lines. Only the most important molecules are shown, i.e. those which reach a certain threshold concentration, depending on element. The TEA results stop at 400 K as TEA has difficulties converging at lower temperatures, even using 2000 iterations. The following molecules with notable concentrations are missing in TEA: SiH_2 , SiH_3 ; CaH ; TiS , TiH , TiN , TiC ; CrH , CrS ; MnH , MnS , MnCl , MnF , MnO ; FeH ; NiH , NiF , NiO , and the following molecules have been omitted by us in the TEA model to improve its convergence: $\text{Si}(\text{CH}_3)_4$, SiCH_3Cl_3 and $\text{Ni}(\text{CO})_4$. See continuation on next page.

species. There are options to just solve the pure gas phase chemical equilibrium or the combined problem with equilibrium condensation. The gas mass density ρ or the total gas pressure p can be chosen as input parameter, besides the temperature T .

5. Results

5.1. TEA benchmark test

To benchmark our results, we use the public Thermo-chemical Equilibrium Abundances (TEA) code (Blecic et al. 2016)⁴. TEA follows the Gibbs free energy minimisation method as described by White et al. (1958) and Eriksson (1971), using a Lagrangian optimisation scheme with Lambda correction algorithm. The Gibbs free energies of the molecules are computed from the thermo-chemical data in the NIST-JANAF tables (Chase 1998), for most molecules given between 100 K and 6000 K, separated between 100 K, provided by Dr. Thomas C. Allison in October

2012 through private communication. The program uses internal spline interpolations to compute the Gibbs free energy between these temperature points. The NIST-JANAF data is available for 600 gaseous molecular species (84 elements).

TEA has been developed in particular to determine the abundances of the major, most-abundant and spectroscopically most active gaseous species expected to be present in hot, giant exoplanetary atmospheres, as these species have a dominant influence on the planetary spectra in the optical and infrared (e.g. Seager & Deming 2010; Burrows 2014). TEA has been benchmarked against CEA (Chemical Equilibrium with Applications), see (Gordon & McBride 1994; McBride & Gordon 1996), as well as against several analytical codes (Burrows & Sharp 1999; Heng & Tsai 2016; Heng & Lyons 2016). These tests typically involve a few elements with a few dozens of molecules between 500 K and 4000 K, where TEA is found to converge robustly within reasonable times (2-3 CPU-sec per (p, T) -point).

For our benchmark test between TEA and GGchem we have selected the following 24 elements of astrophysical interest: H,

⁴ <https://github.com/dzesmin/TEA>




Fig. 3. (continued): The differences in the selection/availability of molecular species explain all visible deviations between GGchem and TEA.

He, Li, C, N, O, F, Na, Mg, Al, Si, P, S, Cl, K, Ca, Ti, V, Cr, Mn, Fe, Ni, Zr and W with solar abundances (Asplund et al. 2009). The task was to compute the chemical composition of the gas at $p = 1$ bar from $T = 6000$ K down to 100 K (or as low as possible), with all available molecules.

Since the TEA-code can presently only handle neutral gas species, we have disregarded all charged species and switched off equilibrium condensation in GGChem for this test. The results are shown in Fig. 3. For practically all depicted molecules, the GGChem results (coloured lines) agree very well with the TEA results, to a precision better than the linewidth of about ~ 0.05 dex. All visible deviations can be explained by some molecules not available in NIST-JANAF, and are hence missing in TEA, or some very large molecules that we have de-selected on purpose in order to improve the convergence of TEA. The missing molecules are listed in the caption of Fig. 3. GGChem uses 421 molecular species plus 24 neutral atoms for this test, whereas TEA uses altogether 400 species including the 24 neutral atoms. The agreement between TEA and GGChem is very reassuring, however, TEA failed to converge within 2000 iterations for temperatures below 400 K.

GGChem needs about 0.004 CPU-sec per point on a 2.8 GHz Linux Laptop. The time consumption is mainly caused by the internal matrix operations which scale as K^3 where K is the number of elements. For the lower temperatures in this benchmark test, GGChem uses quadruple precision arithmetics. For temperatures > 1000 K, the code switches to double precision which takes only 0.0013 CPU-sec per point for 24 elements. To compute the 100 temperature points requested for this benchmark test, the total computational time consumption of GGChem is 0.71 CPU-sec, where about half of it is initialisation. This is in sharp contrast to TEA which needs about 45 CPU-min per point with 2000 iterations, resulting in altogether 74 CPU-hours to complete this benchmark test. Clearly, we have pushed TEA to its limits in this benchmark test, TEA should normally be run with about 100 iterations where it performs as stated above. However, using TEA with 2000 iterations in this benchmark test to obtain results for as low as possible temperatures, we conclude that GGChem is more than 10^5 times faster than TEA.

5.2. Abundance of major molecules

In a first application, we have used the simplified setup of Heng & Tsai (2016) with 4 elements (H, C, N, O) and 9 molecules (H_2 , H_2O , CO, CO_2 , N_2 , NH_3 , HCN, C_2H_2 , C_2H_4) with solar abundances (Asplund et al. 2009). The results shown in Fig. 4 can be directly compared to the upper part of figure 1 in (Heng & Tsai 2016). GGChem needs about 0.12 milli CPU-sec per point in this setup.

Figure 4 compares these results to two other GGChem models. The dashed lines show the results from the full model with 24 elements and 445 molecules as specified in Sect. 5.1, and the dotted lines show the results from a full model with equilibrium condensation as specified in Sect. 5.3. The comparison shows that, although the simplified 9-molecule model results in very reasonable approximations for the concentrations of the major molecules between 500 K and 3000 K, there are some notable deviations around 1000-2000 K which are likely to be relevant for the retrieval of atmospheric properties from spectroscopic observations. These deviations are due to the consumption of oxygen by (i) the formation of SiO , $\text{Mg}(\text{OH})_2$ and $\text{Fe}(\text{OH})_2$ molecules, and (ii) silicate and phyllosilicate condensation. Both effects reduce the concentrations of H_2O and CO_2 , and increase




Fig. 4. Concentrations of major molecules at 1 bar according to a simplified 9-molecule model in chemical equilibrium (full lines), in comparison to a model with 24 elements and 445 molecules (dashed) and a full model with equilibrium condensation (dotted).

C_2H_2 and C_2H_4 concentrations, by up to 0.5 dex around 1500 K. This effect is discussed further in Sect. 5.6.

5.3. The condensation of the elements

In the following, we study the results of a model with equilibrium condensation switched on, at constant pressure $p = 1$ bar, for 22 elements (H, He, Li, C, N, O, Na, Mg, Al, Si, S, Cl, K, Ca, Ti, V, Cr, Mn, Fe, Ni, Zr, W) and charged species, with solar abundances. We have omitted fluorine and phosphorus in this model because of our possibly incomplete data for fluorine and phosphorus condensates. Altogether 388 gas phase and 190 condensed species are taken into account in this model. Gas phase species include 365 molecules, 22 free atoms and the free electron. Condensed species include 38 liquids (Table D.3). The thermo-chemical data of the condensed species are mainly taken from the SUPCRTBL database (Table D.4), completed/replaced by our own fits to NIST-JANAF data of 92 condensed species (Table D.2), plus five further condensed species ($\text{CaTiO}_3[\text{s}]$, $\text{MnS}[\text{s}]$, $\text{NiS}[\text{s}]$, $\text{NiS}_2[\text{s}]$, $\text{Ni}_3\text{S}_2[\text{s}]$) taken from Sharp & Huebner, which are neither available in NIST-JANAF nor in SUPCRTBL.

Figure 5 shows the dust/gas mass ratio as function of temperature in phase equilibrium, calculated as

$$\text{dust/gas} = \frac{\rho_{\text{cond}}}{\rho} = \frac{1}{\rho} \sum_{j[\text{cond}]} n_{j[\text{cond}]} m_{j[\text{cond}]}, \quad (20)$$

where $j[\text{cond}]$ is the index of a condensed species, $n_{j[\text{cond}]}$ its particle density (i.e. number of condensed units per volume) and $m_{j[\text{cond}]}$ its mass. Tungsten establishes a dust/gas ratio of $\sim 10^{-9}$ at $T \lesssim 2250$ K, zirconium dioxide brings it to $\sim 10^{-7.5}$ at 2000 K, and aluminium increases it to $\sim 10^{-4}$ at 1900 K. Subsequently, calcium, iron, silicon and magnesium compounds build up a dust/gas ratio of $\sim 10^{-2.38} \approx 0.004$ at 1500 K, which then remains about constant toward much lower temperatures. The condensation of titanium, nickel, vanadium, chromium, manganese, sodium and potassium compounds make only minor contributions to dust/gas(T). The value of dust/gas ≈ 0.004 is significantly lower than the standard value of dust/gas $\approx 1/100$ (e.g. Beckwith et al. 1990; Chiang & Goldreich 1997; D'Alessio et al. 1998; Dullemond et al. 2002). At around 650 K, sulphur starts




Fig. 5. Upper part: The dust/gas mass ratio in phase equilibrium as function of temperature at constant $p = 1$ bar for solar element abundances. The arrows indicate where the elements start forming condensed phases in significant amounts. The elements marked in red have a significant influence on the shape of the dust/gas(T)-curve. The **lower part** shows the remaining element abundances in the gas phase.

to condense which increases dust/gas to about 0.0045 with further minor contributions by chlorine and lithium. Just below 500 K, several minerals start to become hydrated by the incorporation of either water or hydroxyl groups, forming so-called phyllosilicates. This additional intake of oxygen atoms increases the dust/gas ratio to about 0.0052. Around 240 K and 140 K, water ice and ammonia condense, respectively, which finally brings the dust/gas ratio to a value of almost exactly 1/100.

The lower part of Fig. 5 shows the depletion of the elements from the gas phase due to condensation. At 1000 K, for example, the gas consists mainly of S and Cl besides H, He, C, N and O, whereas all other elements, which are usually more abundant like Fe, Si and Mg, are already depleted by more than 8 orders of magnitude. The intake of oxygen in silicates and phyllosilicates leads to a substantial increase of the C/O ratio in the gas phase, from $C/O = 0.55$ at $T = 2500$ K to $C/O = 0.71$ at $T = 1000$ K (silicates) and $C/O = 0.83$ at $T = 300$ K (phyllosilicates). This increase of C/O may well have observational consequences in brown dwarf and giant gas planet spectra. Once water condenses, the C/O ratio becomes very large, for example $C/O > 10^6$ at 150 K, and the gas consists mainly of CH_4 and NH_3 (beside H_2




Fig. 6. The onset of condensation at $p = 1$ bar for solar abundances in phase equilibrium. The graph shows the concentration of the various condensed species with respect to hydrogen nuclei.

and noble gases), similar to the atmosphere of Titan (Fulchignoni et al. 2005)⁵. Remarkably, at $p = 1$ bar, not a single carbon atom has been incorporated into any condensed state in phase equilibrium down to 100 K for solar abundances.

The onset of condensation: Figure 6 visualises the first stages of condensation in phase equilibrium at $p = 1$ bar. W[s] (crystalline tungsten) is found to be the first stable condensate at 2216 K, followed by ZrO_2 [s] (baddeleyite) at 2027 K. Further phase transitions are as follows: Al-gas \rightarrow Al_2O_3 [s] (corundum) at 1957 K, Ti-gas \rightarrow $CaTiO_3$ [s] (perovskite) at 1913 K, Ca-gas \rightarrow $Ca_2Al_2SiO_7$ [s] (gehlenite) at 1880 K, Fe-gas \rightarrow Fe[l] (liquid iron) at 1841 K. These are all type-1 (gas \rightarrow condensed) phase transitions, featured by a smooth, rounded-off increase of the concentration of the new condensate as function of temperature. This gradual increase is mirrored by a gradual decrease of one element abundance in the gas phase, an element which was not affected by condensation before, to keep the partial pressure equal to the vapour pressure $p^{\text{vap}}(T)$ of the new condensate.

At 1820 K and 1777 K, the first type-2 transitions (condensed \rightarrow condensed) occur, namely Fe[l] \rightarrow Fe[s] (solid iron) and Al_2O_3 [s] \rightarrow $MgAl_2O_4$ [s] (spinel), respectively. These transitions transform one combination of condensates into another, without substantial changes of any gas phase abundances. Type-2 transitions occur suddenly, they do not change the number of elements affected by condensation, hence the number of independent variables in the equation system to be solved, and thus the total number of present condensates (see App. B). Type-2 transitions have a box-like appearance in Fig. 6.

The appearance of the major condensates: The condensation of the elements continues in Fig. 6 with a number of further type-1 transitions: Si-gas \rightarrow SiO [s] (silicon monoxide) at 1729 K at 1 bar, V-gas \rightarrow VO [s] (vanadium monoxide) at 1710 K, Ni-gas \rightarrow Ni[s] (crystalline nickel) at 1690 K, Mg-gas \rightarrow Mg_2SiO_4 [s] (fosterite) at 1661 K and Cr-gas \rightarrow Cr[s] (crystalline chromium) at 1513 K. At this stage, the build-up of the dust/gas ratio to a value of about 0.004 is mostly completed and there are 11 elements which are strongly reduced by condensation (see lower

⁵ Titan's atmosphere has little hydrogen, hence it consists mainly of N_2 with just some CH_4 , whereas oxygen is absent due to water condensation.

part of Fig. 5), which sets the number of simultaneously present condensates to 11 (as explained in App. B).

Increasing complexity 1: A number of further type-2 transitions occur in Fig. 6 towards lower temperatures, namely $\text{Ca}_2\text{Al}_2\text{Si}_2\text{O}_7[\text{s}] \rightarrow \text{Ca}_2\text{MgSi}_2\text{O}_7[\text{s}]$ (*akermanite*) at 1685 K, $\text{MgAl}_2\text{O}_4[\text{s}] \rightarrow \text{CaAl}_2\text{Si}_2\text{O}_8[\text{s}]$ (*anorthite*) at 1440 K, $\text{CaTiO}_3[\text{s}] \rightarrow \text{Ti}_4\text{O}_7[\text{s}]$ (*titanium oxide*) at 1420 K, $\text{Ca}_2\text{MgSi}_2\text{O}_7[\text{s}] \rightarrow \text{CaMgSi}_2\text{O}_6[\text{s}]$ (*diopside*) at 1420 K, $\text{Cr}[\text{s}] \rightarrow \text{MgCr}_2\text{O}_4[\text{s}]$ (*picromerite*) at 1336 K, $\text{ZrO}_2[\text{s}] \rightarrow \text{ZrSiO}_4[\text{s}]$ (*zircon*) at 1334 K, $\text{SiO}[\text{s}] \rightarrow \text{MgSiO}_3[\text{s}]$ (*enstatite*) at 1292 K, $\text{Ti}_4\text{O}_7[\text{s}] \rightarrow \text{MnTiO}_3[\text{s}]$ (*pyrophanite*) at 1211 K, and $\text{VO}[\text{s}] \rightarrow \text{V}_2\text{O}_3[\text{s}]$ (*karelianite*) at 1194 K. These type-2 transitions can have some side-effects on other more abundant condensates, so strictly speaking, each type-2 phase transition affects a combination of solids where one is formed, one is consumed and a number of other, more abundant condensates adjust their concentrations. This applies in particular to $\text{MgSiO}_3[\text{s}]$ and $\text{Mg}_2\text{SiO}_4[\text{s}]$ which are the main reservoir for condensed silicon and magnesium, which are affected by most type-2 transitions.

On the left side of Fig. 6, a few more type-1 transitions take place, namely Mn-gas $\rightarrow \text{MnS}[\text{s}]$ (*alabandite*) at $T = 1290$ K, Na-gas $\rightarrow \text{NaAlSi}_3\text{O}_8[\text{s}]$ (*albite*) at $T = 1231$ K and K-gas $\rightarrow \text{KAlSi}_3\text{O}_8[\text{s}]$ (*microcline*) at $T = 1154$ K, increasing the number of simultaneously present condensates to 14. At 1000 K, all three major components of *feldspar* are present ($\text{KAlSi}_3\text{O}_8[\text{s}] - \text{NaAlSi}_3\text{O}_8[\text{s}] - \text{CaAl}_2\text{Si}_2\text{O}_8[\text{s}]$) which is the most common mineral on Earth making up about 41% of its continental crust (Anderson & Anderson 2010). *Feldspar* is the major constituent of *basalt*, which is a key component of oceanic crust as well as the principal volcanic rock on Earth.

Increasing complexity 2: Figure 7 continues to show the model results down to 460 K. Most prominently, we see three more type-1 transitions: S-gas $\rightarrow \text{FeS}[\text{s}]$ (*troilite*) at 678 K, Cl-gas $\rightarrow \text{NaCl}[\text{s}]$ (*halite*) at 613 K and Li-gas $\rightarrow \text{LiCl}[\text{s}]$ (*lithium-chloride*) at 573 K, increasing the number of simultaneously present condensates to 17. Further type-2 transitions occurs as well, in particular $\text{MnTiO}_3[\text{s}] \rightarrow \text{CaTiSiO}_5[\text{s}]$ (*sphene*) at 814 K, $\text{MnS}[\text{s}] \rightarrow \text{Mn}_3\text{Al}_2\text{Si}_3\text{O}_{12}[\text{s}]$ (*spessartine*) at 670 K, $\text{KAlSi}_3\text{O}_8[\text{s}] \rightarrow \text{KMg}_3\text{AlSi}_3\text{O}_{12}\text{H}_2[\text{s}]$ (*phlogopite*) at 520 K, $\text{CaAl}_2\text{Si}_2\text{O}_8[\text{s}] \rightarrow \text{MgAl}_2\text{O}_4[\text{s}]$ (*spinel*) at 514 K, $\text{NaAlSi}_3\text{O}_8[\text{s}] \rightarrow \text{NaMg}_3\text{AlSi}_3\text{O}_{12}\text{H}_2[\text{s}]$ (*sodaphlogopite*) at 502 K, and $\text{CaTiSiO}_5[\text{s}] \rightarrow \text{FeTiO}_3[\text{s}]$ (*ilmenite*) at 486 K. $\text{KMg}_3\text{AlSi}_3\text{O}_{12}\text{H}_2$ and $\text{NaMg}_3\text{AlSi}_3\text{O}_{12}\text{H}_2$ are the first two phyllosilicates, with additional intake of hydrogen and oxygen.

The formation of phyllosilicates: Figure 8 shows the results down to 100 K. Again, a number of complex type-2 transformations occur, where most of the condensates are replaced by phyllosilicates, step by step. The most significant phyllosilicate is $\text{Mg}_3\text{Si}_2\text{O}_9\text{H}_4[\text{s}]$ (*lizardite*) which replaces $\text{Mg}_2\text{SiO}_4[\text{s}]$ at 345 K. At 270 K, the main condensates are $\text{Mg}_3\text{Si}_2\text{O}_9\text{H}_4[\text{s}]$, $\text{FeS}[\text{s}]$, $\text{Fe}_3\text{O}_4[\text{s}]$ (*magnetite*), $\text{NaMg}_3\text{AlSi}_3\text{O}_{12}\text{H}_2[\text{s}]$, $\text{Fe}_3\text{Al}_2\text{Si}_3\text{O}_{12}[\text{s}]$ (*almandine*), $\text{Ca}_3\text{Al}_2\text{Si}_3\text{O}_{12}[\text{s}]$ (*grossular*) and $\text{Ni}_3\text{S}_2[\text{s}]$ (*heazlewoodite*), whereas the more simple condensates commonly known from higher temperatures (fosterite, enstatite, solid Fe, etc.) are all gone.

The condensation of water and ammonia: The condensation of the elements is completed by two major type-1 transitions: O-gas $\rightarrow \text{H}_2\text{O}[\text{s}]$ (*water ice*) at 247 K and N-gas $\rightarrow \text{NH}_3[\text{s}]$ (*ammonia ice*) at 141 K. We do not observe the condensation of $\text{CO}[\text{s}]$ or $\text{CO}_2[\text{s}]$ (*dry ice*) in our equilibrium models with solar abundances, although at 1 bar, CO_2 is well-known to deposit to a solid at $-79^\circ\text{C} = 195$ K on Earth. This is because the Earth atmosphere is




Fig. 7. Concentration of condensed species at medium temperatures 460 K – 1000 K at $p = 1$ bar for solar abundances. Note the changed scaling of the y-axis in comparison to Fig. 6.




Fig. 8. Concentration of condensed species at temperatures 100 K – 460 K at $p = 1$ bar for solar abundances.

oxygen-rich and relatively hydrogen-poor. For solar abundances, all carbon is locked up in gaseous CH_4 . At 100 K, all elements except hydrogen, carbon and noble gases have condensed into solid phases, and the composition of the gas phase is extremely simple – just H_2 , He, CH_4 , and noble gases. The next major phase transition to occur would be C-gas $\rightarrow \text{CH}_4[\text{s}]$ (*methane ice*), but this transition occurs below 100 K and our models do not reach these temperatures. We note that we are currently lacking organic liquids/solids in our collection of condensed species, which might change some of these conclusions.

5.4. CO-blocking

Figure 9 demonstrates the well-known impact of the carbon-to-oxygen ratio $C/O = \epsilon_C/\epsilon_O$ on the composition of the gas phase, and on the condensation. The figure is designed to match the results of Mollière et al. (2015), although here (in contrast to Mollière et al.) we are plotting particle concentrations. At temperatures between about 850 K and 4000 K at 10 mbar, CO is the dominant gas species in the carbon-oxygen system with a dissociation energy of about 11.1 eV. Therefore, CO almost com-




Fig. 9. The impact of CO-blocking on the composition of the gas at constant pressure and temperature. In the upper plot, the full lines represent the results from a pure gas phase model without condensation (24 elements and 445 molecules, see Sect. 5.1), and the dashed lines show the results from the model including equilibrium condensation (Sect. 5.3). ϵ_0 is a fixed value here, whereas ϵ_C varies.

pletely consumes all carbon when $C/O \lesssim 1$ or all oxygen when $C/O \gtrsim 1$, an effect known as *CO-blocking* (see e.g. Sedlmayr & Dominik 1995; Sedlmayr 1997; Beck et al. 1992). The effect is particularly strong on hydro-carbon molecules like C_2H_2 (*acetylene*) which are suppressed by more than 6 orders of magnitude if $C/O < 1$.

Closer inspection of our results (Fig. 9) shows that the transition actually takes place around $C/O \approx 0.96$, which agrees well with the Mollière et al. results, who have been using the CEA code (Gordon & McBride 1994; McBride & Gordon 1996). Figure 1 in Madhusudhan (2012) actually shows the same effect, although Madhusudhan has used a log-scaling of the C/O -axis where deviations of order 5% are simply not visible. The reason for this asymmetry is the formation of the SiO molecule with a dissociation energy of 8.2 eV. If $C/O \lesssim 1$, the small amount of excess oxygen is first of all consumed to form SiO molecules, with similar consequences for the abundances of water and hydro-carbon molecules as used from the CO-blocking. We measure numerically that the abundances of H_2O and CH_4 intersect at $C/O = 1.00$ when $\epsilon_{Si} \rightarrow 0$, at 0.98 when $\epsilon_{Si} = 7.3$, at 0.96 when $\epsilon_{Si} = 7.51$ (solar), at 0.92 when $\epsilon_{Si} = 7.7$, and at 0.82 when $\epsilon_{Si} = 8$.

Figure 9 also shows the influence of the CO-blocking on the condensation. At 1500 K and 10 mbar, C[s] (*graphite*) can condense at $C/O \gtrsim 1.07$, SiC[s] (*silicon carbide*) at $C/O \gtrsim 0.96$ and TiC[s] (*titanium carbide*) at $C/O \gtrsim 0.93$. At smaller C/O , the oxygen-containing solids can form, here in particular $Al_2O_3[s]$ (*corundum*), $Ca_2Al_2SiO_7[s]$ (*gehlenite*), $MgAl_2O_4[s]$ (*spinel*), $Ca_2MgSi_2O_7[s]$ (*akermanite*), $CaTiO_3[s]$ (*perovskite*) and Mg_2SiO_4 (*fosterite*). Other condensates like W[s] and $ZrO_2[s]$ (not shown) and Fe[s] are not affected by C/O , because

they either do not contain oxygen or carbon, or have even higher dissociation energies than CO, like $ZrO_2[s]$.

The feedback of condensation on the molecular abundances for $C/O > 1$ is more substantial than for $C/O < 1$. The molecular concentrations of C_2H_2 and CH_4 are found to not increase any further once carbon dust is present, as any surplus carbon is simply converted into additional carbon dust, but not into additional hydro-carbon molecules in phase equilibrium. We have actually used the total (gas + dust) abundances of oxygen and carbon to create Fig. 9, because the gas-phase C/O only increases to about 1.1 even if much larger values of the total C/O are used.

At lower temperatures ($T < 850$ K at $p = 10$ mbar), however, CH_4 becomes the dominating carbon gas phase species, whereas CO becomes a trace species, and so the spell is broken. In a hydrogen-rich gas, CH_4 is thermodynamically more favourable than any solid carbon compound at low temperatures, which leads to high water concentrations independent of C/O , and no carbon dust in phase equilibrium. This confirms our earlier findings in kinetic cloud formation models for carbon-rich atmospheres (Fig. 6 in Helling et al. 2017).

5.5. Phase diagrams of the elements

Figure 10 shows some results of the equilibrium condensation model (22 elements, charges, 365 molecules and 190 condensed species, see Sect. 5.3) in the entire (p, T) -plane between 10^{-9} bar and 100 bar, and between 100 K and 2500 K, for solar abundances. The figures indicate the most abundant species that contain certain elements, after multiplication by their stoichiometric factor, hence the main reservoirs of the elements as function of pressure and temperature.

We generally see increasing complexity from high to low temperature, from ions over atoms to simple molecules, to complex molecules, and then to condensates with increasingly complex stoichiometry and phyllosilicates at the lowest temperatures. In the first approximation, all molecular transitions, as well as the type-1 and type-2 phase transitions discussed in Sect. 5.3 simply occur at lower temperatures when the pressure is decreased, roughly a factor of 2-4 in temperature for 10 decades of pressure decrease. This leads to numerous slanted and slightly curved phase transition lines, similar to phase diagrams in thermodynamics. Most prominent molecular transitions are $H \rightarrow H_2$ (at ~ 2200 K), $CO \rightarrow CH_4$ (at ~ 650 K) and $N_2 \rightarrow NH_3$ (at ~ 320 K), where the numbers in brackets refer to a pressure of $p = 10^{-4}$ bar.

The details, however, are in fact much more complicated, in particular concerning the gas \rightarrow solid transitions. The order of certain phase transitions as described in Sect. 5.3 may be different, and some condensates may be present only in certain parts of the (p, T) -plane. The main silicates can form below about 1700 K to 1100 K, and eventually the elements Mg, Si, Na, Al, K and Mn can form thermodynamically stable phyllosilicates at temperatures below about 600 K to 200 K.

We want to stress, however, that all these results depend on the chosen element abundances, as demonstrated for the case of varying C/O in Sect. 5.4. The results do also depend technically on the choice of elements, so if another element is taken into account in addition, it will interfere with all others which can potentially change the results shown in Fig. 10. Therefore, our “phase diagrams” are not as general as their name suggests. The results should not be taken out of context and be applied to e.g. the atmospheres of rocky planets. The proper way to do this is to run the GGChem-code for the expected pressures, temperatures and local element abundances in the atmosphere.




Fig. 10. Gas phase/condensed species which contain most of the elements in the (p , T)-plane. Solid species are marked by '[s]' and liquid species by '[l]'. At low temperatures, the elements Mg, Si, Na, Al and K are mostly present in form of phyllosilicates.




Fig. 10. continued: The only further element found to form phyllosilicates in phase equilibrium is manganese.

5.6. The C/O-ratio as affected by metal oxide molecules and condensation

Figure 11 (right side) shows the C/O ratio in the gas phase as affected by condensation in the (p, T) -plane. One of the most significant results of this paper again shows, namely that the solar value of about $C/O = 0.55$ only holds prior to the condensation of the main silicates. The consumption of oxygen by silicate formation is substantial, and increases the C/O ratio to about 0.71. Once the silicates become hydrated to form phyllosilicates, which happens to the most abundant silicates between about 200 K to 400 K in the equilibrium model (depending on pressure), the C/O ratio jumps to 0.83, and then gets very large once water condenses⁶. The little blue area in the central top of Fig. 11 is not a computational error – it is a small region in the (p, T) -plane where we find solid carbon C[s] (*graphite*) to condense.

To clarify the effect of condensation on gas phase C/O, we compare our results to a pure gas phase model (without conden-

sation) where we compute an “effective” C/O as

$$(C/O)_{\text{eff}} = \tilde{n}_{(C)}/\tilde{n}_{(O)} \quad (21)$$

$$\tilde{n}_{(C)} = \sum_i n_i \tilde{s}_{i,C} \quad , \quad \tilde{n}_{(O)} = \sum_i n_i \tilde{s}_{i,O} \quad ,$$

where only atoms and molecules are counted here if they are made of the elements H, C, N and O, but we set the stoichiometric coefficient $\tilde{s}_{i,C} = 0$ if a molecule i contains any other (metal) elements like Mg, Si or Fe. In a pure gas phase model, as plotted on the left side of Fig. 11, the proper $C/O \approx 0.55$ is of course constant everywhere in the (p, T) -plane by assumption, but $(C/O)_{\text{eff}}$ will be different because of the oxygen and carbon that is bound in molecules which are neglected in Eq. (21). We do this because in fast spectral retrieval models (e.g. Lavie et al. 2017), the influence of these metals on the concentration of molecules like H_2O and CO_2 is often neglected, i.e. $(C/O)_{\text{eff}} \approx C/O$ is assumed.

In Fig. 11 (left side) we see that $(C/O)_{\text{eff}}$ may become substantially larger than C/O at lower temperatures. $(C/O)_{\text{eff}}$ increases from the solar value 0.55 to about 0.58 due to the formation of SiO between about 2000 K and 3000 K, and then in-

⁶ Such strong changes in C/O, however, were not found in kinetic cloud formation models (Helling et al. 2016).




Fig. 11. The C/O ratio in the gas phase affected by the formation of metal oxide and hydroxide molecules (left) and condensation of silicates, phyllosilicates, and water ice (right). On the left side, we plot the “effective” C/O that is contained in the gas in form of molecules made of H-C-N-O only, but do not count metal oxide and hydroxide molecules. The contour lines shown are 0.58 (after SiO formation) and 0.8 (after $\text{Mg}(\text{OH})_2$ and $\text{Fe}(\text{OH})_2$ formation). On the right side, we plot the resulting gas phase abundances after the condensation of solids and liquids. The increase of $\text{C/O} > 0.7$ is due to the consumption of oxygen by the major silicates like $\text{MgSiO}_3[\text{s}]$ and $\text{Mg}_2\text{SiO}_4[\text{s}]$, and the further increase to > 0.8 is due to phyllosilicates. The sharp increase at low temperatures (red area) is due to water condensation.




Fig. 12. The concentration ($x_j = n_j/n_{\text{H}}$) of the three major magnesium-silicates in the (p, T) -plane in the equilibrium condensation model.

creases further to values above 0.8 once $\text{Mg}(\text{OH})_2$ and $\text{Fe}(\text{OH})_2$ become abundant in the gas phase, which happens at temperatures between about 600 K and 1100 K.

In both discussed cases, the large pure gas phase model and the equilibrium condensation model, the reason for the increase of the (effective) C/O ratio is the same – the consumption of oxygen atoms by the formation of metal compounds, either oxide and hydroxide molecules or condensates, which are then missing for the formation of gaseous molecules like H_2O and CO_2 . The effect is actually similar in both models, because each Mg, Si and Fe atom, on average, consumes about 1–2 oxygen atoms in both cases at low temperatures. The main difference is that in models including condensation, the establishment of large C/O ratios already occurs at higher temperatures, between about 1200 K to 1600 K. We discuss this point further in Sect. 7.

Figure 12 shows the abundance of the three most significant Mg-silicates in the (p, T) -plane. As described in Sect. 5.3, the two major reservoirs for condensed Mg and Si at relatively high temperatures are $\text{MgSiO}_3[\text{s}]$ (*enstatite*) and $\text{Mg}_2\text{SiO}_4[\text{s}]$ (*fos-*

terite). However, some more stable silicates exist, which contain e.g. Ca, Fe, Ti, Mn or Cr. These can form in smaller amounts only, due to element abundance constraints, so they “steal” some Mg and Si from enstatite and fosterite, which affects the concentrations of enstatite and fosterite in complicated ways. Fosterite generally becomes stable at slightly higher temperatures than enstatite, but once formed, enstatite can become the most abundant condensate in certain parts of the (p, T) -plane, along with $\text{Fe}[\text{s}]$, $\text{Fe}_2\text{SiO}_4[\text{s}]$, $\text{FeS}[\text{s}]$ or $\text{Fe}_2\text{O}_3[\text{s}]$.

Below 200 – 400 K (depending on pressure) both Mg-silicates are combined and hydrated to form the phyllosilicate $\text{Mg}_3\text{Si}_2\text{O}_9\text{H}_4[\text{s}]$ (*lizardite*). Interestingly, there is no feedback of water condensation on this phyllosilicate, the concentration of lizardite stays constant towards very low temperatures, showing that the formation of lizardite is thermodynamically favoured over the formation of water ice.

We conclude from Fig. 10 that Mg, Si, Na, Al, K and Mn can form thermodynamically stable phyllosilicates at low temperatures. Beside the major phyllosilicate $\text{Mg}_3\text{Si}_2\text{O}_9\text{H}_4[\text{s}]$




Fig. 13. **Left:** The nucleation rate of tungsten per hydrogen nucleus J_*/n_{H} [1/s] as function of gas pressure and temperature. **Centre:** The size of the critical cluster N_* , **Right:** The nucleation timescale τ_{nuc} [yrs] according to Eq. (25). The white dashed contour line marks saturation $S = 1$, the gas is supersaturated to the left of this line.

(lizardite), we also find $\text{NaMg}_3\text{AlSi}_3\text{O}_{12}\text{H}_2[\text{s}]$ (sodaphlogopite), $\text{CaAl}_2\text{Si}_2\text{O}_{10}\text{H}_4[\text{s}]$ (lawsonite), $\text{Mg}_3\text{Si}_4\text{O}_{12}\text{H}_2[\text{s}]$ (talc), $\text{KMg}_3\text{AlSi}_3\text{O}_{12}\text{H}_2[\text{s}]$ (phlogopite), $\text{KFe}_3\text{AlSi}_3\text{O}_{12}\text{H}_2[\text{s}]$ (annite), $\text{KMn}_3\text{AlSi}_3\text{O}_{12}\text{H}_2[\text{s}]$ (Mn-biotite) and $\text{MnAl}_2\text{SiO}_7\text{H}_2[\text{s}]$ (Mn-chloritoid) in less amounts, due to element abundance constraints.

6. The nucleation of tungsten

An interesting side-result of our models is that solid tungsten $\text{W}[\text{s}]$ is found to be the *first condensate*, i.e. metallic tungsten is the first solid material that becomes thermodynamically stable in any galactic/stellar/atmospheric cooling flow which is initially too hot to contain any condensed materials. The question arises whether tungsten could provide the first nucleation seeds required for the growth of all other, more abundant solid/liquid materials which only become stable at lower temperatures. See (Gail & Sedlmayr 1998) for a more detailed introduction to the search for the first condensate.

6.1. Is there enough tungsten?

To estimate the total number of seed particles that can be provided by tungsten, let us assume that one seed particle consists of $N_l \approx 10 - 1000$ tungsten atoms. Using the solar abundance of tungsten $\epsilon_{\text{W}} = 7.1 \times 10^{-12}$ (Asplund et al. 2009), the concentration of tungsten seed particles cannot exceed

$$\frac{n_{\text{seeds}}}{n_{\text{H}}} = \frac{\epsilon_{\text{W}}}{N_l} \approx 7 \times 10^{-15} - 7 \times 10^{-13}. \quad (22)$$

Next, we assume that the various silicates, iron and all other abundant condensates will grow on top of these seed particles later, to form a solid mantle of unique thickness. The total available volume of solid material per H nucleus is about

$$V_{\text{dust}} \approx (\epsilon_{\text{Si}} + \epsilon_{\text{Mg}} + \epsilon_{\text{Fe}}) V_1 \approx (1.2 - 2.7) \times 10^{-27} \text{ cm}^3, \quad (23)$$

where we have used solar abundances and where $V_1 \approx (1.2 - 3) \times 10^{-23} \text{ cm}^3$ is the dust volume per heavy atom (the sum of Si, Mg and Fe atoms in the condensate), estimated from the mass and solid densities of iron, MgSiO_3 and Mg_2SiO_4 . The radius of the dust particles forming this way would be given by $n_{\text{seeds}} \frac{4\pi}{3} a^3 = n_{\text{H}} V_{\text{dust}}$, i.e.

$$a = \left(\frac{3}{4\pi} V_{\text{dust}} \frac{n_{\text{H}}}{n_{\text{seeds}}} \right)^{1/3} \approx 0.075 - 0.47 \mu\text{m}, \quad (24)$$

which falls surprisingly well into the size range of observed dust particles in space. This proves nothing of course, but it shows that there is enough tungsten in a gas of solar abundances to provide the seed particles needed for astrophysical bulk condensation. In fact, from the simple estimate presented here we can conclude that the concentration of seed particles *must* not exceed about 10^{-15} to 10^{-13} per hydrogen nucleus, otherwise too many, too small dust particles would form by condensation.

In application to M-type AGB-star winds, Höfner et al. (2016) have shown that only silicate grains in a certain size range $0.1 \mu\text{m}$ to $1 \mu\text{m}$ can efficiently drive an outflow by radiation pressure, via their large scattering opacity around $\lambda \sim 1 \mu\text{m}$. In contrast, if their absorption coefficients dominate at those wavelengths (as is true when they contain some iron), the grains rather get heated in the strong IR radiation field and evaporate (Woitke 2006). In order to obtain these particular particle sizes, Höfner et al. (2016) needed to assume a certain seed particle concentration in their time-dependent hydrodynamical and dust formation models, $n_{\text{seeds}}/n_{\text{H}} \approx 3 \times 10^{-14}$ to 3×10^{-15} .

6.2. Timescales and supercooling

The nucleation rate of crystalline tungsten has been calculated according to classical nucleation theory in the formulation of Gail et al. (1984), with surface tension $\sigma = 3340 \text{ erg/cm}^2$ (Tran et al. 2016) and parameter $N_{1/2} = 10$ (the cluster size where the surface tension reduces to half its bulk value). The supersaturation ratio S and the particle density of atomic tungsten n_{W} are calculated by a pure gas-phase GGchem model without condensation. The resulting nucleation rates per hydrogen nucleus are shown in Fig. 13, from which we estimate that a supercooling of about $300 - 400 \text{ K}$ is needed to start the effective nucleation of tungsten. The time required to form these seed particles is

$$\tau_{\text{nuc}} = \frac{n_{\text{seeds}}}{J_*} = \frac{n_{\text{seeds}}}{n_{\text{H}}} \frac{n_{\text{H}}}{J_*} \approx 10^{-14} \left(\frac{J_*}{n_{\text{H}}} \right)^{-1} \quad (25)$$

From Fig. 13 we read off that the formation timescales of the tungsten seed particles would be rather long and would require high gas pressures, for example $\tau_{\text{nuc}} = 1 \text{ yr}$ at 0.1 bar and 1600 K . At lower gas pressures, the tungsten nucleation timescale increases fast, exceeding the age of the universe around 10^{-6} bar . This fast increase is mainly due to an increase of the critical cluster size N_* (central part of Fig. 13).

7. Summary and conclusions

A fast and versatile public code has been developed, called GGChem, to calculate the chemical composition of astrophysical gases in chemical equilibrium down to 100 K, with or without equilibrium condensation. We have collected and compared the thermo-chemical data for molecules and condensates from different sources, and have presented tables with temperature-fits of molecular equilibrium constants $k_p(T)$ and Gibbs free energies $\Delta G_f^\circ(T)$ which behave robustly towards low temperatures and can safely be applied down to 100 K. The data comprises 552 molecules and 257 condensates, including 38 liquids.

For pure gas-phase applications, GGChem has been thoroughly tested against the TEA-code (Blecic et al. 2016), revealing close to perfect agreement. Differences arise only when the selection of molecules differs. In GGChem, we are using a number of molecules with thermo-chemical data from Barklem & Collet (2016), which are not available in NIST-JANAF, in particular metal hydrides like CaH, TiH and FeH. GGChem is found to converge fast and robustly down to 100 K under all tested circumstances including very small and very large selections of elements and molecules, as well as for unusual element abundances.

Concerning very sparse models with just a few molecules and elements (for example H, C, N, O to quickly determine the abundances of the spectroscopically active molecules like H₂O and CH₄), we want to emphasise that the combined effect of elements like Mg, Si and Fe is substantial and might lead to an overestimation of the C/O ratio if these elements are neglected. For solar abundances (C/O = 0.55) the combined abundance of Si, Mg and Fe alone is about 10^{-4} which is larger than the nitrogen abundance ($\epsilon_N = 6.8 \times 10^{-5}$) and can be compared to the carbon and oxygen abundances ($\epsilon_C = 2.7 \times 10^{-4}$, $\epsilon_O = 4.9 \times 10^{-4}$). Since molecules like SiO, Mg(OH)₂ and Fe(OH)₂ form at low temperatures, every Si, Mg and Fe atom takes away about $x \approx 1-2$ oxygen atoms from the C-N-O system. If we consider the effective C/O ratio remaining in the C-N-O system, the result for solar element abundances is

$$(C/O)_{\text{eff}} = \frac{\epsilon_C}{\epsilon_O - x(\epsilon_{\text{Si}} + \epsilon_{\text{Mg}} + \epsilon_{\text{Fe}})} \approx \begin{cases} 0.70 & \text{for } x = 1 \\ 0.95 & \text{for } x = 2 \end{cases} \quad (26)$$

Thus, if C/O ratios are derived from observations based on sparse chemical models, where elements like Mg, Si and Fe are lacking, the results must be expected to be too large.

We have discussed the condensation sequence of the elements in phase equilibrium, generally confirming the results found earlier by Sharp & Huebner (1990) and Lodders (2003). Differences arise mostly from the different selection/availability of the thermo-chemical data for condensed species, but we did not find any evidence for systematic differences due to different numerical techniques. An important step here was to create a link to the geophysical database SUPCRTBL (Zimmer et al. 2016), from which we have extracted the thermo-chemical data for 121 condensed species, including phyllosilicates. The numerical methods rather affect the computational speed and stability of the codes.

A straightforward result from our models is that the dust/gas mass ratio in a solar composition gas is expected to be 0.0045 rather than 0.01. The latter (standard) value of 0.01 is only obtained in our models once water and ammonia ices condense at low temperatures. The ice formation increases the mass of the condensates by a factor of ~ 2.5 and the volume of the condensates by a factor of ~ 6 (from $2.7 \times 10^{-27} \text{ cm}^3$ to $1.7 \times 10^{-26} \text{ cm}^3$ per hydrogen nucleus to be precise).

Our models show that phyllosilicates (“wet silicates”) are the most abundant condensates at low temperatures in a solar composition gas in phase equilibrium. The most abundant phyllosilicate is identified to be Mg₃Si₂O₉H₄[s] (*lizardite*) which contains the majority of Mg and Si below 200 K – 400 K in phase equilibrium, depending on pressure. In a similar way, a few other phyllosilicates are found to consume most of Na, Al, K and Mn, some of which may form already at 500 K. The phyllosilicates have a notable influence on the dust/gas ratio and will increase the C/O ratio in the gas phase to about 0.83 because of the additional intake of oxygen.

Whether or not these phyllosilicates (and all other condensates) actually form in space is not discussed in this paper, as this is a paper just about chemical and phase equilibrium. A recent kinetic approach for the formation of phyllosilicates in protoplanetary discs has been submitted to A&A by Thi et al., based on a warm surface chemistry model. Thi et al. find that despite the high energy barriers involved in chemisorption and diffusion, water and hydroxyl can efficiently migrate into the bulk lattice within typical lifetimes of protoplanetary discs if the gas and dust temperatures are of order 80 K to 700 K. The maximum intake of water and hydroxyl is a parameter in these models.

As a side result, we discussed whether tungsten could be the first condensate in space and could provide the seed particles for astrophysical dust formation. We argue that solid tungsten becomes stable already at very high temperatures of order 2000 K, that it could provide just about the right numbers of seed particle per H-atom to form micron-sized silicate grains, but that a strong super-cooling of order 300 K–400 K is required and that high pressures of the order of milli-bars are necessary to make tungsten nucleation an efficient process.

Acknowledgements: We thank Dr. Wing-Fai Thi for pointing us toward the SUPCRTBL database and for his comments on the manuscript. The research leading to these results has received funding from the European Union Seventh Framework Programme FP7-2011 under grant agreement no 284405. Jasmina Blecic is supported by NASA through the NASA ROSES-2016/Exoplanets Research Program, grant NNX17AC03G. Christiane Helling highlights the hospitality of the Rijksuniversiteit Groningen and the Universiteit van Amsterdam with travel support from NWO and LKBF.

References

- Ackerman, A. S. & Marley, M. S. 2001, *ApJ*, 556, 872
- Allard, F. & Hauschildt, P. H. 1995, *ApJ*, 445, 433
- Allard, F., Hauschildt, P. H., Alexander, D. R., & Starrfield, S. 1997, *ARA&A*, 35, 137
- Allard, F., Hauschildt, P. H., Alexander, D. R., Tamanai, A., & Schweitzer, A. 2001, *ApJ*, 556, 357
- Anderson, R. S. & Anderson, S. P. 2010, *Geomorphology: The Mechanics and Chemistry of Landscapes* (Cambridge University Press)
- Asplund, M., Grevesse, N., Sauval, A. J., & Scott, P. 2009, *ARA&A*, 47, 481
- Auer, L. H. & Mihalas, D. 1968, *ApJ*, 151, 311
- Barklem, P. S. & Collet, R. 2016, *A&A*, 588, A96
- Beck, H. K. B., Gail, H.-P., Henkel, R., & Sedlmayr, E. 1992, *A&A*, 265, 626
- Beckwith, S. V. W., Sargent, A. I., Chini, R. S., & Guesten, R. 1990, *AJ*, 99, 924
- Beichman, C., Benneke, B., Knutson, H., et al. 2014, *PASP*, 126, 1134
- Benneke, B. 2015, *ArXiv e-prints* [arXiv:1504.07655]
- Berline, S. & Bricker, C. 1969, *Journal of Chemical Education*, 46, 499
- Bilger, C., Rimmer, P., & Helling, C. 2013, *MNRAS*, 435, 1888
- Birkby, J. L., de Kok, R. J., Brogi, M., Schwarz, H., & Snellen, I. A. G. 2017, *AJ*, 153, 138
- Blecic, J., Harrington, J., & Bowman, M. O. 2016, *ApJS*, 225, 4
- Booth, R. A., Clarke, C. J., Madhusudhan, N., & Ilee, J. D. 2017, *MNRAS*, 469, 3994
- Brogi, M., de Kok, R. J., Albrecht, S., et al. 2016, *ApJ*, 817, 106

- Burrows, A. 2014, arXiv preprint arXiv:1409.7320
- Burrows, A., Burgasser, A. J., Kirkpatrick, J. D., et al. 2002, *ApJ*, 573, 394
- Burrows, A. & Sharp, C. M. 1999, *ApJ*, 512, 843
- Chase, Jr., M. 1998, *Journal of Physical and Chemical Reference Data*, Monograph 9, update of 3rd Edition
- Chase, M. W. 1986, *JANAF thermochemical tables* (American Chemical Society; New York)
- Chase, M. W., Curnutt, J. L., Downey, J. R., et al. 1982, in *Journal of Physics Conference Series*, Vol. 11, *Journal of Physics Conference Series*, 695–940
- Chiang, E. I. & Goldreich, P. 1997, *ApJ*, 490, 368
- Cooper, C. S. & Showman, A. P. 2006, *ApJ*, 649, 1048
- Crossfield, I. J. M., Barman, T., Hansen, B. M. S., & Howard, A. W. 2013, *A&A*, 559, A33
- D'Alessio, P., Canto, J., Calvet, N., & Lizano, S. 1998, *ApJ*, 500, 411
- de Kok, R. J., Brogi, M., Snellen, I. A. G., et al. 2013, *A&A*, 554, A82
- Decin, L., Richards, A. M. S., Waters, L. B. F. M., et al. 2017, *ArXiv e-prints* [arXiv:1704.05237]
- Désert, J.-M., Vidal-Madjar, A., Lecavelier Des Etangs, A., et al. 2008, *A&A*, 492, 585
- Dullemond, C. P., van Zadelhoff, G. J., & Natta, A. 2002, *A&A*, 389, 464
- Eriksson, G. 1971, *Acta Chem.Scand.*, 2651
- Fraine, J., Deming, D., Benneke, B., et al. 2014, *Nature*, 513, 526
- Fulchignoni, M., Ferri, F., Angrilli, F., et al. 2005, *Nature*, 438, 785
- Gaidos, E. J. 2000, *Icarus*, 145, 637
- Gail, H.-P., Keller, R., & Sedlmayr, E. 1984, *A&A*, 133, 320
- Gail, H.-P., Scholz, M., & Pucci, A. 2016, *A&A*, 591, A17
- Gail, H.-P. & Sedlmayr, E. 1986, *A&A*, 166, 225
- Gail, H.-P. & Sedlmayr, E. 1998, *Faraday Discussions*, 109, 303
- Gail, H.-P., Wetzel, S., Pucci, A., & Tamanai, A. 2013, *A&A*, 555, A119
- Golriz, S. S., Blommaert, J. A. D. L., Vanhollebeke, E., et al. 2014, *MNRAS*, 443, 3402
- Gordon, S. & McBride, B. J. 1994, *Computer Program for Calculation of Complex Chemical Equilibrium Compositions and Applications. I. Analysis, Reference Publication RP-1311*, NASA, describes theory and numerical algorithms behind CEA computer program
- Griffith, C. A. & Yelle, R. V. 1999, *ApJ*, 519, L85
- Grillmair, C. J., Burrows, A., Charbonneau, D., et al. 2008, *Nature*, 456, 767
- Gustafsson, B. 1971, *A&A*, 10, 187
- Gustafsson, B., Edvardsson, B., Eriksson, K., et al. 2008, *A&A*, 486, 951
- Helling, C., Ackerman, A., Allard, F., et al. 2008, *MNRAS*, 391, 1854
- Helling, C. & Casewell, S. 2014, *A&A Rev.*, 22, 80
- Helling, C. & Fomins, A. 2013, *Philosophical Transactions of the Royal Society of London Series A*, 371, 20110581
- Helling, C., Jorgensen, U. G., Plez, B., & Johnson, H. R. 1996, *A&A*, 315, 194
- Helling, C., Lee, G., Dobbs-Dixon, I., et al. 2016, *MNRAS*, 460, 855
- Helling, C. & Lucas, W. 2009, *MNRAS*, 398, 985
- Helling, C., Thi, W.-F., Woitke, P., & Fridlund, M. 2006, *A&A*, 451, L9
- Helling, C., Tootill, D., Woitke, P., & Lee, G. 2017, *A&A*, 603, A123
- Heng, K. & Lyons, J. R. 2016, *ApJ*, 817, 149
- Heng, K. & Tsai, S.-M. 2016, *ApJ*, 829, 104
- Höfner, S. & Andersen, A. C. 2007, *A&A*, 465, L39
- Höfner, S., Bladh, S., Aringer, B., & Ahuja, R. 2016, *A&A*, 594, A108
- Hollenbach, D. & McKee, C. F. 1989, *ApJ*, 342, 306
- Hu, R. & Seager, S. 2014, *ApJ*, 784, 63
- Jeong, K. S., Winters, J. M., Le Bertre, T., & Sedlmayr, E. 2003, *A&A*, 407, 191
- Johnson, J. W., Oelkers, E. H., & Helgeson, H. C. 1992, *Computers and Geosciences*, 18, 899
- Juncher, D., Jørgensen, U. G., & Helling, C. 2017, *ArXiv e-prints* [arXiv:1708.06976]
- Kopparapu, R. k., Kasting, J. F., & Zahnle, K. J. 2012, *ApJ*, 745, 77
- Kreidberg, L., Bean, J. L., Désert, J.-M., et al. 2014, *ApJ*, 793, L27
- Lavie, B., Mendonça, J. M., Mordasini, C., et al. 2017, *AJ*, 154, 91
- Line, M. R., Liang, M. C., & Yung, Y. L. 2010, *ApJ*, 717, 496
- Line, M. R., Stevenson, K. B., Bean, J., et al. 2016, *AJ*, 152, 203
- Line, M. R. & Yung, Y. L. 2013, *ApJ*, 779, 3
- Lodders, K. 2003, *ApJ*, 591, 1220
- Lodders, K. & Fegley, B. 2002, *Icarus*, 155, 393
- Lunine, J. I., Hubbard, W. B., & Marley, M. S. 1986, *ApJ*, 310, 238
- Lüttke, M. 2002, PhD thesis, Thesis, Tech. Univ. Berlin
- Madhusudhan, N. 2012, *ApJ*, 758, 36
- Madhusudhan, N., Harrington, J., Stevenson, K. B., et al. 2011, *Nature*, 469, 64
- Mahapatra, G., Helling, C., & Miguel, Y. 2017, *ArXiv e-prints* [arXiv:1706.07219]
- Marley, M. S., Gelino, C., Stephens, D., Lunine, J. I., & Freedman, R. 1999, *ApJ*, 513, 879
- Marley, M. S. & Leggett, S. K. 2009, *Astrophysics and Space Science Proceedings*, 10, 101
- Marley, M. S. & Robinson, T. D. 2015, *ARA&A*, 53, 279
- Marley, M. S., Seager, S., Saumon, D., et al. 2002, *ApJ*, 568, 335
- Mbarek, R. & Kempton, E. M.-R. 2016, *ApJ*, 827, 121
- McBride, B. J. & Gordon, S. 1996, *Computer Program for Calculation of Complex Chemical Equilibrium Compositions and Applications. II. User's Manual and Program Description*, Reference Publication RP-1311-P2, NASA
- Miller-Ricci, E., Seager, S., & Sasselov, D. 2009, *ApJ*, 690, 1056
- Mollière, P., van Boekel, R., Dullemond, C., Henning, T., & Mordasini, C. 2015, *ApJ*, 813, 47
- Morley, C. V., Knutson, H., Line, M., et al. 2017, *AJ*, 153, 86
- Moses, J. I. 2014, *Philosophical Transactions of the Royal Society of London Series A*, 372, 20130073
- Moses, J. I., Visscher, C., Fortney, J. J., et al. 2011, *ApJ*, 737, 15
- Oreshenko, M., Lavie, B., Grimm, S. L., et al. 2017, *ApJ*, 847, L3
- Patzner, A. B. C. 2007, in *Astronomical Society of the Pacific Conference Series*, Vol. 378, *Why Galaxies Care About AGB Stars: Their Importance as Actors and Probes*, ed. F. Kerschbaum, C. Charbonnel, & R. F. Wing, 181
- Patzner, A. B. C., Köhler, T. M., & Sedlmayr, E. 1995, *Planet. Space Sci.*, 43, 1233
- Prydz, R. & Goodwin, R. D. 1972, *Journal of Chemical Thermodynamics*, 4, 1
- Rauer, H., Aerts, C., Cabrera, J., & PLATO Team. 2016, *Astronomische Nachrichten*, 337, 961
- Rimmer, P. B. & Helling, C. 2016, *ApJS*, 224, 9
- Sanchez-Lopez, A., Lopez-Puertas, M., Funke, B., et al. 2017, in *Highlights on Spanish Astrophysics IX*, ed. S. Arribas, A. Alonso-Herrero, F. Figueras, C. Hernández-Monteagudo, A. Sánchez-Lavega, & S. Pérez-Hoyos, 576–576
- Seager, S. & Deming, D. 2010, *Annual Review of Astronomy and Astrophysics*, 48, 631
- Sedlmayr, E. 1997, *Ap&SS*, 251, 103
- Sedlmayr, E. & Dominik, C. 1995, *Space Sci. Rev.*, 73, 211
- Semenov, D. & Wiebe, D. 2011, *ApJS*, 196, 25
- Sharp, C. M. & Huebner, W. F. 1990, *ApJS*, 72, 417
- Spang, III, H. 1962, *SIAM review*, 4, 343
- Stock, J. 2008, diploma thesis, Technische Universität Berlin, Straße des 17. Juni 135, 10623 Berlin, Germany
- Stull, D. & Prophet, H. 1971, *JANAF Thermochemical Tables*, NSRDS-NBS (U.S. Government Printing Office)
- Tran, R., Xu, Z., Radhakrishnan, B., et al. 2016, *Nature Scientific Data*, Volume 3, id. 160080 (2016), 3, 160080
- Tsuji, T. 1965, *PASJ*, 17, 152
- Tsuji, T. 1973, *A&A*, 23, 411
- Tsuji, T. 2005, *ApJ*, 621, 1033
- Tsuji, T., Ohnaka, K., Aoki, W., & Nakajima, T. 1996, *A&A*, 308, L29
- Van Eck, S., Neyskens, P., Jorissen, A., et al. 2017, *A&A*, 601, A10
- Venot, O., Hébrard, E., Agúndez, M., et al. 2012, *A&A*, 546, A43
- Visscher, C., Lodders, K., & Fegley, Jr., B. 2006, *ApJ*, 648, 1181
- Visscher, C., Lodders, K., & Fegley, Jr., B. 2010, *ApJ*, 716, 1060
- Weast, R. 1971, *CRC Handbook of Chemistry and Physics: A Ready-reference Book of Chemical and Physical Data*, CRC handbook series (Chemical Rubber Company)
- White, W. B., Johnson, S. M., & Dantzig, G. B. 1958, *J. Chem. Phys.*, 28, 751
- Witte, S., Helling, C., & Hauschildt, P. H. 2009, *A&A*, 506, 1367
- Woitke, P. 2001, in *Reviews in Modern Astronomy*, Vol. 14, *Reviews in Modern Astronomy*, ed. R. E. Schielicke, 185
- Woitke, P. 2006, *A&A*, 460, L9
- Woitke, P. & Helling, C. 2004, *A&A*, 414, 335
- Woitke, P., Helling, C., & Turner, G. 2017, *Graphical Comparison of Mineral Gibbs Free Energy Data*, Other, DOI 10.17630/3ba9b432-3564-4e6c-b350-687eb4b501a5, University of St Andrews
- Woitke, P., Min, M., Pinte, C., et al. 2016, *A&A*, 586, A103
- Worters, M., Millard, D., Hunter, G., Helling, C., & Woitke, P. 2017, *Comparison Catalogue of Gas-Equilibrium Constants*, Kp, Other, DOI 10.17630/cf49aee3-3bb3-4198-9030-c4d51f16beee, University of St Andrews
- Yaws, C. 1999, *Chemical Properties Handbook*, Chemical engineering books (McGraw-Hill Education)
- Zahnle, K., Marley, M. S., Freedman, R. S., Lodders, K., & Fortney, J. J. 2009, *ApJ*, 701, L20
- Zimmer, K., Zhang, Y., Lu, P., et al. 2016, *Computers and Geosciences*, 90, 97

Appendix A: Solving gas phase chemical equilibrium down to 100 K

Solving the system of non-linear equations (Eq. 5), i.e. finding the root vector of atomic partial pressures which satisfies the element conservation equations after elimination of all molecular partial pressures in chemical equilibrium according to Eq. (4), seems to be a simple numerical standard problem. Indeed, a simple Newton-Raphson iteration may lead to quick success at high temperatures. However, at low temperatures, the solution vector becomes more and more degenerate, with some atomic partial pressures approaching values $< 10^{-\text{several hundred}}$ dyn/cm², for example carbon, whereas others like hydrogen stay of order 1. This leads to a number of numerical problems, in particular

- (i) the conditional number of the Jacobi matrix becomes large, i.e. the resolution of the linearised equation system is not possible without substantial losses of precision, and
- (ii) the Newton-Raphson method only works satisfactory when a good initial estimate of the solution vector is provided, which becomes increasingly difficult at low temperatures.

We have developed a stable algorithm that solves these problems under all tested circumstances, including unusual element abundances and pressures, down to 100 K. The first necessary step was to switch to quadruple precision arithmetics for temperatures $T < 1000$ K, which solves problem (i).

Figure A.1 sketches how we solve problem (ii). The algorithm starts by sorting the elements according to their abundances. This hierarchy can change substantially by condensation as certain elements will be removed almost entirely from the gas phase, for example Ca and Al, whereas other elements stay in the gas phase for longer, for example S and Cl. In the example shown in Fig. A.1, we consider $K=9$ elements. There are 9 pre-iteration steps performed in this case, before a final Newton-Raphson iteration is carried out to solve the element conservation equations (Eq. 5) for all elements. The task of the pre-iterations is to provide good initial estimates of the atomic partial pressures for this final iteration, they are the key to obtain code stability.

In each pre-iteration, we only consider a subset of elements and molecules which are composed of these elements. Each pre-iteration starts with an estimate of the atomic partial pressure of a new element (short red bar) using the values of all other atomic partial pressures computed so far (long grey bar). The pre-iteration then continues by improving the atomic pressures of the last N elements (long red bars, $N=4$ in the figure) by taking into account the mutual feedbacks, however without changing the partial pressures of the other, even more abundant elements on the left (short grey bars).

For example, pre-iteration 1 (“pre-1”) only considers hydrogen with species H and H₂, from which a first guess of p_H is obtained. Next comes oxygen with additional species O, OH, O₂, HO₂, H₂O and O₃. During the first stage of pre-2, a first guess of p_O is obtained by solving the element conservation equation for O at given p_H . During the second stage of pre-2, the feedback between O on H is taken into account by solving the two coupled conservation equations for H and O by a Newton-Raphson method to improve p_H , p_O . The next element is carbon with additional species C, CH, CO, C₂, HCO, CH₂, H₂CO, CH₃, CH₄, CO₂, C₂H, C₂H₂, C₂H₄, C₂H₄O, C₂O, C₃, C₃O₂, C₄, C₅ from which p_C is first estimated and then determined, including the mutual feedback on H and O, by solving three coupled non-linear algebraic equations, and so on. The algorithm continues this way until Mg, where we stop refining p_H in stage 2 of pre-5. We are using the hierarchical order of the elements here,




Fig. A.1. Algorithm used in GGchem to iteratively solve the chemical equilibrium problem for total number of elements $K=9$, and number of elements taken into account during the pre-iterations $N=4$, see text.

i.e. whatever the outcome of p_{Mg} will be, it can only have a minor feedback on p_H since $\epsilon_{Mg} \ll \epsilon_H$. When sulphur enters the pre-iteration phase at pre-8, the atomic partial pressures of O, C and N are already frozen. The algorithm continues this way, before the full system of element conservation equations (Eq. 5) is solved with the Newton Raphson method, using the atomic pressures from the pre-iteration phase as initial estimates.

These pre-iterations in fact work so well, that the final full Newton Raphson iteration only makes minor corrections $< 5\%$ in all our benchmark tests, at most, which produces no visible changes in Fig. 3, for example, only another boost in performance. Figure A.1 sketches the algorithm for $N=4$, however we achieved best overall performance with $N=5$. Further improvements of code stability and performance are achieved by dividing the resulting atomic partial pressures by their initial guesses after each Newton-Raphson iteration, then store these ratios into the memory, and use them as correction factors next time to obtain improved initial guesses. This simple idea was the key to achieve code convergence down to 100 K.

At $T = 96$ K, however, the equilibrium constant k_p of the molecule (WO₃)₄ hits 10^{+4932} , the maximum number that can be represented in quadruple precision, and the code crashes. It might be technically possible to go deeper in temperature by eliminating a few big molecules, however these are just the molecules which dominate at those temperatures, see Fig. 3.

Appendix B: Solving phase equilibrium

We formulate the element conservation equations in the presence of condensed species in the following way

$$\epsilon_k^0 = \epsilon_k + \sum_{j=1}^N s_{j,k} c_j, \quad (\text{B.1})$$

where ϵ_k^0 are the total element abundances (or “initial” element abundances before condensation), ϵ_k are the element abundances in the gas phase and $c_j = n_j^{\text{cond}}/n_{(H)}$ is the concentration of condensed species j per hydrogen nucleus. N is the current number of present condensates, and $s_{j,k}$ are the stoichiometric factors of elements k in condensate j . Equation (B.1) is a generalised variant of (Eq. 5). Note that all equations for the gas phase in Sect. 2.1 remain valid when using ϵ_k .

An essential realisation is that Eq. (B.1) should *never* be used in computer codes to “subtract the dust”, as this would unavoidably lead to a substantial loss of numerical precision at low

temperatures. Some ϵ_k are reduced by several hundreds of orders of magnitude at low temperatures, which will soon exceed the numerical resolution required to perform this subtraction precisely, even when using quadruple precision arithmetics.

We have developed an iterative algorithm which avoids this problem by applying corrections $\delta\epsilon_k$ to ϵ_k to eventually solve

$$S_j(\epsilon_k) \leq 1 \quad \text{all condensates} \quad (\text{B.2})$$

at given temperature T and density $n_{\text{(H)}}$. Provided that we have selected the present condensates correctly (all non-selected condensates must have $S < 1$), this equation simplifies to

$$S_j(\epsilon_k) = 1 \quad \text{selected condensates.} \quad (\text{B.3})$$

The dependencies on ϵ_k are marked here because all gas phase equilibrium results only depend on $(n_{\text{(H)}}, T, \epsilon_k)$ (where $n_{\text{(H)}}$ and T are parameters to the problem), and so do the supersaturation ratios S_j . We choose N_{ind} element abundances as independent variables of the problem

$$x_i = \{\delta\epsilon_i \mid i = 1, \dots, N_{\text{ind}}\}, \quad (\text{B.4})$$

whereas the dependent variables are

$$y_j = \sum_i^{N_{\text{ind}}} A_{ji} x_i = \begin{cases} \delta c_j & \text{for } j = 1, \dots, N \\ \delta\epsilon_{k(j)} & \text{for } j = N + 1, \dots, N + D. \end{cases} \quad (\text{B.5})$$

A_{ji} is the conversion matrix and D the number of dependent elements with indices $k(j)$. One iteration step is completed by

$$\epsilon_k \rightarrow \epsilon_k + \delta\epsilon_k \quad (\text{B.6})$$

$$c_j \rightarrow c_j + \delta c_j \quad (\text{B.7})$$

The concentrations of the condensed species c_j are simply a byproduct of these computations, but they are *not used* during the iterations. The element conservation is implemented in the conversion matrix A_{ji} as explained below, and we carefully check that Eq. (B.1) remains valid after each iteration step.

To explain these definitions, let us consider the condensation of Al_2O_3 , MgSiO_3 and Mg_2SiO_4 as an example. There are 3 condensates in this case ($N = 3$) and 4 affected elements ($N_{\text{ind}} + D = 4$). In order to proceed, we must select three of them ($N_{\text{ind}} = 3, D = 1$) to obtain 3 equations (Eq. B.3) for 3 unknowns. As default rule, we pick the least abundant elements (here Al, Si and Mg) as independent variables, i.e. $i = 1$ is Al, $i = 2$ is Si and $i = 3$ is Mg. The other elements (here O, $D = 1$) become dependent elements, $k(4)$ is O. To find the conversion matrix A_{ji} we write down the conservation equations for all elements affected by condensation as

$$\begin{aligned} 2\delta c_{\text{Al}_2\text{O}_3} &+ \delta\epsilon_{\text{Al}} &= 0 \\ \delta c_{\text{MgSiO}_3} &+ \delta c_{\text{Mg}_2\text{SiO}_4} + \delta\epsilon_{\text{Si}} &= 0 \\ \delta c_{\text{MgSiO}_3} &+ 2\delta c_{\text{Mg}_2\text{SiO}_4} + \delta\epsilon_{\text{Mg}} &= 0 \\ 3\delta c_{\text{Al}_2\text{O}_3} &+ 3\delta c_{\text{MgSiO}_3} + 4\delta c_{\text{Mg}_2\text{SiO}_4} + \delta\epsilon_{\text{O}} &= 0. \end{aligned}$$

By a stepwise elimination scheme, we find the conversion matrix A in this case to be

$$\begin{pmatrix} \delta c_{\text{Al}_2\text{O}_3} \\ \delta c_{\text{MgSiO}_3} \\ \delta c_{\text{Mg}_2\text{SiO}_4} \\ \delta\epsilon_{\text{O}} \end{pmatrix} = A \cdot \begin{pmatrix} \delta\epsilon_{\text{Al}} \\ \delta\epsilon_{\text{Si}} \\ \delta\epsilon_{\text{Mg}} \end{pmatrix} = \begin{pmatrix} -\frac{1}{2} & 0 & 0 \\ 0 & -2 & 1 \\ 0 & 1 & -1 \\ \frac{3}{2} & 2 & 1 \end{pmatrix} \cdot \begin{pmatrix} \delta\epsilon_{\text{Al}} \\ \delta\epsilon_{\text{Si}} \\ \delta\epsilon_{\text{Mg}} \end{pmatrix}$$

The conversion matrix A is a $N \times (N+D)$ matrix which describes how the dependent variables change when the independent element abundances are changed. This transformation allows us to

solve Eq. (B.3) as system of $N = 3$ coupled non-linear algebraic equations for the $N = 3$ unknowns ($\delta\epsilon_{\text{Al}}, \delta\epsilon_{\text{Si}}, \delta\epsilon_{\text{Mg}}$), which succeeds easily by a Newton-Raphson iteration with several internal calls of the equilibrium chemistry to compute the Jacobi matrix at every iteration step. We have to make sure, however, to limit the Newton-Raphson step δx to prevent negative c_j or negative ϵ_k . This method is found to generally converge very quickly at all temperatures, within less than about 10 iteration steps, depending on the quality of the initial guesses of ϵ_k (here $\epsilon_{\text{Al}}, \epsilon_{\text{Si}}, \epsilon_{\text{Mg}}$).

To solve the problem of bad initial guesses for ϵ_k , we store all successfully computed results (ϵ_k, c_j) as function of $(n_{\text{(H)}}, T)$ into a database and take the initial guesses from the closest $(n_{\text{(H)}}, T)$ database point next time. This way, the program automatically becomes more stable and faster over time. But initially, we must run GGChem with small temperature steps from warm to cold.

Some practical problems may still arise, because of the somewhat unclear selection of condensed species and independent elements. The selection of condensed species is guided by monitoring the supersaturation ratios of all condensates during the iterations. If a new condensate becomes supersaturated, it will be added to the set of selected condensates, or will replace one of them. We have implemented a number and hand-crafted criteria here. For example, one condensate cannot be present simultaneously as liquid and solid. Application of the algorithm explained above would fail already in the first step, as there is only one affected element but two condensates – only one can be present. Another example is CaTiO_3 , MgTiO_3 and $\text{CaMgSi}_2\text{O}_6$, where the latter is a linear combination of the former two, in which case the elimination scheme to obtain the conversion matrix fails – only two of them can be present under any circumstances. This leads us to some quite interesting general insights into the nature of phase equilibrium:

- $N = N_{\text{ind}}$ must hold, i.e. the number of simultaneously present condensates must equal the number of independent elements. N_{ind} is limited by the number of elements affected by condensation K ($K = N_{\text{ind}} + D$ with $D \geq 0$). For any given set of condensates, this means that we can simply count the number of elements from which they are made, K , and the number of present condensates in phase equilibrium N must not exceed that number: $N \leq K$.
- Stoichiometric linear combinations of condensates must not be present simultaneously in phase equilibrium. This applies in particular to the solid/liquid phase transitions.
- The number of dependent elements D is given by the number of elements which cannot be completely converted into condensates, due to element conservation constraints, at temperature T . Toward low T , N increases and D decreases monotonically. Eventually, D becomes very small, only counting elements like O, N, C and H which are affected by condensation but need very low temperatures to completely condense⁷. All other elements eventually vanish almost entirely from the gas phase at low temperatures.

For example, N increases by one around 600 K, as soon as FeS[s] becomes stable. Since $\epsilon_{\text{Fe}}^0 > \epsilon_{\text{S}}^0$, the stoichiometry of that condensate makes it possible to completely remove sulphur from the gas phase. Without such new opportunities to lock away elements, some sudden transformations might occur between several condensates (type-2 phase transitions), but the total number of them will remain constant. Practically, we observe that condition (b)

⁷ Although eventually, O and N will condense anyway, in form of water and ammonia.

is also valid for linear combinations of condensates which conserve all elements but oxygen, the abundance of which is only marginally affected anyway by condensation.

Appendix C: The molecular k_p data

During the preparation phase for this paper, we have collected molecular k_p data from (Tsuiji 1973), (Sharp & Huebner 1990), (Lüttke 2002), (Stock 2008), and (Barklem & Collet 2016) and compared these to our own previous data collection going back to Gail & Sedlmayr (1986). The names of 1155 molecules have been homogenised and the different functional fits of $k_p(T)$ have been overplotted, similar to Fig. 1, and characterised to assess the level of agreement.

Figure C.1 shows the results of the complete comparison study (Worters et al. 2017). Most of the molecules fall into the category “data agrees” (30%) with deviations better than 0.4 dex at 200 K and better than 0.1 dex at 3000 K, disregarding the Tsuiji data. However, for the majority of the molecules, the disagreement is worse, with some cases of quite obvious disagreements.

Similar conclusions have recently been published by Barklem & Collet (2016), who have reviewed the thermochemical data of 291 diatomic and charged diatomic molecules. These deviations are mostly due to uncertainties in the dissociation energies. Barklem & Collet (2016) published their data in terms of partition functions and dissociation energies

$$k_{\text{Bark}} = \frac{p_A p_B}{p_{AB}} = \left(\frac{2\pi m_{\text{red}} kT}{h^2} \right)^{3/2} kT \frac{Q_A Q_B}{Q_{AB}} \exp\left(-\frac{D_{AB}^0}{kT}\right), \quad (\text{C.1})$$

where A and B are atoms, AB is a diatomic molecule, $m_{\text{red}} = m_A m_B / (m_A + m_B)$ is the reduced mass, m_{el} the electron mass, Q_A , Q_B and Q_{AB} are the partition functions, and D_{AB}^0 is the dissociation energy. To compute the equilibrium constants $k_p(T)$ according to our definition from this data, also for charged molecules, we need to combine Eq. (C.1) with Saha functions

$$S_+ = \frac{p_{A^+} p_{\text{el}}}{p_A} = \left(\frac{2\pi m_{\text{el}} kT}{h^2} \right)^{3/2} kT \frac{2Q_{A^+}}{Q_A} \exp\left(-\frac{\chi_A}{kT}\right) \quad (\text{C.2})$$

$$S_- = \frac{p_A p_{\text{el}}}{p_{A^-}} = \left(\frac{2\pi m_{\text{el}} kT}{h^2} \right)^{3/2} kT \frac{2Q_A}{Q_{A^-}} \exp\left(-\frac{\chi_{A^-}}{kT}\right) \quad (\text{C.3})$$

to obtain

$$k_p^{\text{AB}} = \frac{p_{AB}}{p_A p_B} = \left(k_{\text{Bark}}(\text{AB}) \right)^{-1} \quad (\text{C.4})$$

$$k_p^{\text{AB}^+} = \frac{p_{\text{AB}^+} p_{\text{el}}}{p_A p_B} = \left(k_{\text{Bark}}(\text{AB}^+) \right)^{-1} S_+ \quad (\text{C.5})$$

$$k_p^{\text{AB}^-} = \frac{p_{AB^-}}{p_A p_B p_{\text{el}}} = \left(k_{\text{Bark}}(\text{AB}^-) \right)^{-1} (S_-)^{-1}. \quad (\text{C.6})$$

χ_A is the ionisation potential of the neutral atom and χ_{A^-} is the electron affinity of the negative ion. Tables for D_{AB}^0 , χ_A , $Q_A(T)$, $Q_{A^+}(T)$, $Q_{A^-}(T)$, $Q_{AB}(T)$, $Q_{\text{AB}^+}(T)$ and $Q_{\text{AB}^-}(T)$ are given in the electronic appendices of Barklem & Collet (2016), on 42 temperatures points between 10^{-5} K and 10^4 K. The electron affinities are not included, however, so we adopt the following values from [https://en.wikipedia.org/wiki/Electron_affinity_\(data_page\)](https://en.wikipedia.org/wiki/Electron_affinity_(data_page))

$$\begin{aligned} \chi_{\text{H}^-} &= 0.7542 \text{ eV} & \chi_{\text{C}^-} &= 1.2621 \text{ eV} \\ \chi_{\text{O}^-} &= 1.4611 \text{ eV} & \chi_{\text{F}^-} &= 3.4012 \text{ eV} \\ \chi_{\text{Si}^-} &= 1.3895 \text{ eV} & \chi_{\text{S}^-} &= 2.0771 \text{ eV} \\ \chi_{\text{Cl}^-} &= 3.6127 \text{ eV} \end{aligned} \quad (\text{C.7})$$




Fig. C.1. Statistics of molecular $k_p(T)$ agreement between the functional fits provided by (Sharp & Huebner 1990), (Lüttke 2002), (Stock 2008), and (Barklem & Collet 2016).

We have fitted these data between 50 K and 6000 K with a Stock-function (Eq. 17).

Our final choice of $k_p(T)$ data for GGchem is shown in Table D.1. We have given preference to the NIST-JANAF data fitted by (Stock 2008). For molecules available in (Barklem & Collet 2016), but not available in NIST-JANAF, we use the Barklem & Collet data. The TEA benchmark test revealed a few mismatches between the (Stock 2008) data and the TEA data. For each of those molecules, we have re-fitted the NIST-JANAF data ourselves. We do not use any of the (Tsuiji 1973) or (Sharp & Huebner 1990) data directly, although some of the (Stock 2008) data we are using are based on the Tsuiji-data.

Appendix D: The condensed phase data

We have extracted condensed phase Gibbs free energy data from two sources. First, the NIST-JANAF database (Chase et al. 1982; Chase 1986)⁸ provides $\Delta G_{\text{f,NIST}}^{\circ}(T)$ -data [kJ/mol] at $p^{\circ} = 1$ bar in the 7th column of their data-files. In NIST-JANAF, these are the Gibbs free energies of formation from the respective NIST reference states of the elements, which are temperature-dependent

$$\Delta G_{\text{f,NIST}}^{\circ} = G_{\text{NIST}}^{\circ}(\text{A}_a \text{B}_b \text{C}_c [\text{cond}], T) - a G_{\text{NIST}}^{\circ}(\text{Aref}, T) - b G_{\text{NIST}}^{\circ}(\text{Bref}, T) - c G_{\text{NIST}}^{\circ}(\text{Cref}, T). \quad (\text{D.1})$$

To obtain the Gibbs free energy of formation of a condensate from free gas atoms, our $\Delta G_{\text{f}}^{\circ}$ as input for Eq. (12), we need to subtract as

$$\Delta G_{\text{f}}^{\circ} = \Delta G_{\text{f,NIST}}^{\circ}(\text{A}_a \text{B}_b \text{C}_c [\text{cond}], T) - a \Delta G_{\text{f,NIST}}^{\circ}(\text{A}, T) - b \Delta G_{\text{f,NIST}}^{\circ}(\text{B}, T) - c \Delta G_{\text{f,NIST}}^{\circ}(\text{C}, T) \quad (\text{D.2})$$

⁸ <http://kinetics.nist.gov/janaf/>




Fig. D.1. Fitting the vapour pressures of solid and liquid iron from the NIST-JANAF database. The two fitted $p^{\text{vap}}(T)$ curves intersect at 1824 K which is close to the known melting point of 1809 K according to NIST-JANAF. The model will automatically give preference to the phase with the lower vapour pressure. Solid data exist between 100 K and 2200 K, and liquid data between 298 K and 4000 K. The solid curve shoots upward for $T > 2500$ K, but this does not matter as iron condenses as a liquid at those temperatures. When designing our solid Fe vapour pressure $p^{\text{vap}}(T)$ -fit, we just have to make sure that it does not produce a second, spurious intersection point with the liquid curve.

The required input for the vapour pressure (Eq. 9) is obtained in a similar way by subtracting $\Delta G_{f,\text{NIST}}^{\circ}(A_a B_b C_c, T)$ instead.

Second, the geophysical SUPCRTBL database (Zimmer et al. 2016; Johnson et al. 1992)⁹ provides condensed phase $\Delta G_{f,\text{SU}}^{\circ}(T, p)$ data. In this database, the Gibbs free energies of formation are defined in a different way, namely with respect to the reference state of the elements at reference temperature $T_{\text{ref}} = 298.15$ K

$$\Delta G_{f,\text{SU}}^{\circ} = G_{\text{SU}}^{\circ}(A_a B_b C_c[\text{cond}], T) - a G_{\text{SU}}^{\circ}(\text{Aref}, T_{\text{ref}}) - b G_{\text{SU}}^{\circ}(\text{Bref}, T_{\text{ref}}) - c G_{\text{SU}}^{\circ}(\text{Cref}, T_{\text{ref}}). \quad (\text{D.3})$$

The database actually provides $\Delta G_{f,\text{SU}}^{\circ}(p, T)$ according to a term $\int V_i^{\circ} dP$ in Eq. (2) of (Zimmer et al. 2016). However, V° is of order of a few J/bar, and can safely be neglected at $p < \text{a few bar}$ as assumed in this paper. In order to arrive at the input for our Eq. (12), we need to subtract corrections as

$$\Delta G_f^{\circ} = \Delta G_{f,\text{SU}}^{\circ} - a \Delta G_{\text{corr}}^{\circ}(\text{A}) - b \Delta G_{\text{corr}}^{\circ}(\text{B}) - c \Delta G_{\text{corr}}^{\circ}(\text{C}), \quad (\text{D.4})$$

namely the Gibbs free energy differences between the free atoms at temperature T and the respective elements in their standard state at T_{ref} . We could not figure out a way how to consistently perform these conversions from the SUPCRTBL data, as most of the atoms are lacking. However, assuming that the standard states of the elements at T_{ref} are identical in both NIST and SUPCRTBL, we can utilise NIST-JANAF as

$$\begin{aligned} \Delta G_{\text{corr}}^{\circ}(\text{A}) &= G_{\text{NIST}}^{\circ}(\text{A}, T) - G_{\text{NIST}}^{\circ}(\text{Aref}, T_{\text{ref}}) \\ &= \Delta G_{f,\text{NIST}}^{\circ}(\text{A}, T) \\ &\quad + [H_{\text{NIST}}^{\circ}(\text{Aref}, T) - H_{\text{NIST}}^{\circ}(\text{Aref}, T_{\text{ref}})] \\ &\quad - T S_{\text{NIST}}^{\circ}(\text{Aref}, T) + T_{\text{ref}} S_{\text{NIST}}^{\circ}(\text{Aref}, T_{\text{ref}}). \end{aligned} \quad (\text{D.5}) \quad (\text{D.6})$$

where $\Delta G_{f,\text{NIST}}^{\circ}$, $[H_{\text{NIST}}^{\circ}(T) - H_{\text{NIST}}^{\circ}(T_{\text{ref}})]$ and S_{NIST}° are given in the 7th, 5th and 3rd columns of the NIST-JANAF data-files. We

have computed high-precision functional fits of these conversions and list the fit coefficients and functional form in Table D.5.

By applying these atomic corrections to the SUPCRTBL $\Delta G_{f,\text{SU}}^{\circ}$ data according to Eq. (D.4), we have converted that data to our reference states of neutral atoms needed in Eq. (12). We have extracted ΔG_f° data this way for a large number of minerals known on Earth listed in Table D.4. The table lists functional fits for ΔG_f° for 121 solid species, interestingly including some phyllosilicates which host OH or H₂O in their lattice structure. We have ordered the minerals in Table D.4, somewhat arbitrarily, by $\Delta G_{f,\text{SU}}^{\circ}(T_{\text{ref}})/N$, where N is the sum of stoichiometric factors, which gives a first expression of what could be the most thermodynamically stable solid materials (note that SUPCRTBL does not have tungsten). We did not include “aqueous” species (i.e. species solved in liquid water), and applied the following additional selection criteria:

- in case of multiple minerals with the same stoichiometric factors we only included the most stable compound at standard temperature,
- minerals with broken stoichiometric factors, or with any stoichiometric factor > 16 are ignored,
- we did not include arsenic (As) or gallium (Ga) compounds.

⁹ <http://www.indiana.edu/~hydrogeo/supcrtbl.html>

Table D.1. Temperature-fits of k_p data for 552 molecules and ions made from H, He, Li, C, N, O, F, Na, Mg, Al, Si, P, S, Cl, K, Ca, Ti, V, Cr, Mn, Fe, Ni, Zr and W (24 elements).

	molecule	source	fit	a_0	a_1	a_2	a_3	a_4	σ [kJ/mol]
1	H2	(2)	4	5.19096E+04	-1.80117E+00	8.72246E-02	2.56139E-04	-5.35403E-09	
2	Li2	(2)	4	1.23866E+04	1.56495E-02	-9.78117E+00	-3.29003E-04	2.77588E-08	
3	C2	(2)	4	7.13486E+04	-7.53302E-01	-8.82633E+00	8.40914E-05	-8.34907E-10	
4	N2	(2)	4	1.13210E+05	-1.79949E+00	-2.03355E+00	4.31303E-04	-2.48433E-08	
5	O2	(2)	4	5.95336E+04	-1.61517E+00	-3.60976E+00	4.06274E-04	-1.86946E-08	
6	F2	(2)	4	1.87418E+04	-1.49389E+00	-4.22069E+00	4.84494E-04	-3.34918E-08	
7	NA2	(2)	4	8.85584E+03	3.60393E-01	-1.15966E+01	-5.20312E-04	3.52831E-08	
8	MG2	(2)	4	2.95153E+02	-1.54214E+00	3.81836E+00	-7.93259E-04	9.60535E-08	
9	AL2	(2)	4	2.08161E+04	1.98351E-01	-1.28542E+01	-1.49552E-04	1.17923E-08	
10	Si2	(2)	4	3.73518E+04	-2.45866E-01	-1.15745E+01	1.50992E-04	-1.59304E-08	
11	P2	(2)	4	5.85956E+04	-9.74048E-01	-6.75364E+00	2.83200E-04	-3.16725E-08	
12	S2	(2)	4	5.08019E+04	-1.51882E+00	-3.18934E+00	4.61697E-04	-2.60901E-08	
13	CL2	(2)	4	2.87966E+04	-1.37354E+00	-3.87637E+00	2.85709E-04	-1.07670E-08	
14	K2	(2)	4	6.57112E+03	1.69191E-01	-9.50003E+00	-4.24701E-04	-9.32751E-09	
15	H2+	(2)	4	-1.27090E+05	6.76863E-01	-1.27744E+01	4.97800E-04	-2.04564E-08	
16	HE2+	(1)	4	-2.56952E+05	8.42165E-01	-1.31376E+01	4.52726E-04	-1.79413E-08	±0.04
17	C2+	(1)	4	-5.86384E+04	7.92144E-01	-1.67128E+01	5.48351E-04	-3.85179E-08	±0.20
18	N2+	(2)	4	-6.76237E+04	6.57659E-01	-1.52323E+01	4.51807E-04	-1.70741E-08	
19	O2+	(2)	4	-8.05914E+04	7.70714E-01	-1.69329E+01	4.25784E-04	-2.32586E-08	
20	P2+	(1)	4	-6.42167E+04	1.41044E+00	-1.81067E+01	4.70925E-04	-4.58882E-08	±0.26
21	S2+	(1)	4	-6.05511E+04	7.79066E-01	-1.59703E+01	6.14185E-04	-4.51795E-08	±0.33
22	H2-	(2)	4	2.30070E+04	-3.98549E+00	1.35395E+01	4.01297E-04	-1.74259E-08	
23	C2-	(2)	4	1.17887E+05	-4.09781E+00	1.02566E+01	3.78388E-04	-1.57139E-08	
24	LiH	(2)	4	2.81110E+04	-1.41584E+00	-7.86666E-01	4.18528E-04	-2.55524E-08	
25	CH	(2)	4	4.03796E+04	-2.06522E+00	2.73563E+00	6.03534E-04	-2.96038E-08	
26	NH	(2)	4	3.72483E+04	-1.85316E+00	1.80469E+00	3.61042E-04	-1.42478E-08	
27	OH	(2)	4	5.09586E+04	-1.86657E+00	1.28296E+00	3.27612E-04	-1.27895E-08	
28	HF	(2)	4	6.79757E+04	-2.00046E+00	1.23667E+00	3.35407E-04	-1.20171E-08	
29	NAH	(2)	4	2.38678E+04	-1.26079E+00	-1.49175E+00	3.72601E-04	-2.18874E-08	
30	MGH	(2)	4	2.32206E+04	-1.42911E+00	8.11874E-01	3.89732E-04	-2.05585E-08	
31	ALH	(2)	4	3.42887E+04	-1.51855E+00	-1.14794E+00	3.83819E-04	-1.87632E-08	
32	SiH	(2)	4	3.46400E+04	-1.54955E+00	-8.92506E-02	3.23215E-04	-1.63317E-08	
33	PH	(3)	4	3.33009E+04	-1.75432E+00	1.65582E+00	4.35721E-04	-2.40582E-08	±0.06
34	HS	(3)	4	4.23352E+04	-1.58869E+00	-1.54130E-02	2.52514E-04	-9.19992E-09	±0.13
35	HCL	(2)	4	5.13028E+04	-2.13173E+00	2.87709E+00	4.45605E-04	-1.92144E-08	
36	KH	(2)	4	2.18271E+04	-1.21500E+00	-1.44207E+00	4.13969E-04	-3.15761E-08	
37	CAH	(1)	4	2.64607E+04	-1.68708E+00	2.79074E+00	6.21435E-04	-4.76821E-08	±0.03
38	TIH	(1)	4	2.41513E+04	-1.87758E+00	2.14663E+00	1.12481E-03	-8.96218E-08	±0.46
39	CRH	(1)	4	2.23915E+04	-1.63632E+00	1.31500E+00	4.48117E-04	-3.10873E-08	±0.17
40	MNH	(1)	4	3.01855E+04	-1.68515E+00	1.97207E+00	5.43045E-04	-3.20198E-08	±0.03
41	FEH	(1)	4	1.73894E+04	-1.45574E+00	-9.26441E-01	8.18399E-04	-6.07297E-08	±0.22
42	NIH	(1)	4	2.84479E+04	-1.86698E+00	1.56057E+00	4.66092E-04	-2.32895E-08	±0.15
43	HEH+	(1)	4	-1.36386E+05	8.66741E-01	-1.41095E+01	2.66518E-04	-2.54550E-10	±0.23
44	CH+	(2)	4	-8.30023E+04	6.96425E-01	-1.45227E+01	3.80894E-04	-1.77873E-08	
45	NH+	(1)	4	-1.16665E+05	1.26447E+00	-1.55925E+01	5.59511E-05	6.26295E-09	±0.18
46	OH+	(2)	4	-1.02017E+05	6.07635E-01	-1.29560E+01	3.89881E-04	-1.60349E-08	
47	HF+	(1)	4	-1.18079E+05	7.25450E-01	-1.26387E+01	2.19710E-04	-1.57439E-09	±0.11
48	MGH+	(1)	4	-6.62374E+04	8.32839E-01	-1.25711E+01	4.57171E-04	-2.33883E-08	±0.03
49	ALH+	(1)	4	-5.05309E+04	5.24296E-01	-1.17158E+01	7.03377E-04	-4.16652E-08	±0.21
50	SiH+	(2)	4	-5.73762E+04	8.56881E-01	-1.50535E+01	3.49121E-04	-1.81757E-08	
51	PH+	(1)	4	-8.24772E+04	8.49004E-01	-1.29265E+01	3.42109E-04	-2.13700E-08	±0.07
52	SH+	(1)	4	-7.88343E+04	5.93436E-01	-1.25291E+01	3.78752E-04	-1.67501E-08	±0.11
53	HCL+	(1)	4	-9.64393E+04	8.22741E-01	-1.28902E+01	1.66628E-04	4.90184E-09	±0.21
54	CH-	(1)	4	5.46372E+04	-4.15997E+00	1.40591E+01	2.79170E-04	-5.39916E-09	±0.10
55	OH-	(2)	4	7.21674E+04	-4.40770E+00	1.44059E+01	3.38619E-04	-1.35099E-08	
56	SiH-	(1)	4	4.96635E+04	-4.73103E+00	1.83213E+01	7.93958E-04	-4.73984E-08	±0.29
57	HS-	(1)	4	6.85217E+04	-4.39583E+00	1.51490E+01	3.32732E-04	-9.75243E-09	±0.13
58	CN	(2)	4	9.02324E+04	-1.75861E+00	-1.56088E+00	4.20862E-04	-1.59741E-08	
59	CO	(2)	4	1.28998E+05	-1.75498E+00	-3.16258E+00	4.13362E-04	-2.35800E-08	
60	CF	(2)	4	6.47022E+04	-1.43350E+00	-3.23877E+00	3.54145E-04	-2.03239E-08	
61	SIC	(2)	4	5.38781E+04	4.16289E-01	-1.62625E+01	-2.85599E-04	1.44412E-08	
62	CP	(2)	4	6.98314E+04	-1.27244E+00	-4.42830E+00	3.42881E-04	-2.86410E-08	
63	CS	(2)	4	8.54423E+04	-1.49342E+00	-4.26643E+00	3.66935E-04	-2.34112E-08	
64	CCL	(2)	4	4.00861E+04	-1.21265E+00	-3.98298E+00	2.56087E-04	-1.37754E-08	
65	CN+	(2)	4	-7.34125E+04	1.01025E+00	-1.63642E+01	4.25271E-04	-2.35543E-08	
66	CO+	(1)	4	-3.35805E+04	5.48177E-01	-1.53832E+01	5.17815E-04	-2.97321E-08	±0.14
67	CN-	(2)	4	1.34538E+05	-4.23936E+00	1.17219E+01	4.10824E-04	-2.43610E-08	
68	CS-	(1)	4	8.88849E+04	-4.44100E+00	1.43923E+01	6.13513E-04	-4.02613E-08	±0.17

All k_p data for standard pressure $p^\circ = 1$ bar in cgs-units $[(\text{dyn}/\text{cm}^2)^{1-n}]$, where n is the number of atoms and electrons in the molecule. Data sources: (1) = [Barklem & Collet \(2016\)](#) converted and fitted between 50 K and 6000 K. (2) = [Stock \(2008\)](#) based on NIST-JANAF ([Chase 1986](#)). (3) = fits made by the authors to NIST-JANAF data for the full available temperature range, usually 100 K to 6000 K. The last column is the standard deviation of the fit from the data points [kJ/mol]. (4) = [Andreas Gauger \(priv.comm\)](#). (5) = ([Tsuji 1973](#)) data re-fitted by [Stock \(2008\)](#). The following formula apply, where $\theta = 5040/T$:

$$\begin{aligned} \text{fit} = 4: \quad \ln k_p(T) &= (1 - n) \ln(1 \text{ bar}) + \left(a_0/T + a_1 \ln T + a_2 + a_3 T + a_4 T^2 \right) \\ \text{fit} = 5: \quad \log_{10} k_p(T) &= -a_0 - a_1 \theta - a_2 \log_{10} \theta - a_3 (\log_{10} \theta)^2 - a_4 (\log_{10} \theta)^3 \end{aligned}$$

Table D.1. continued

	molecule	source	fit	a_0	a_1	a_2	a_3	a_4	σ [kJ/mol]
69	NO	(2)	4	7.55243E+04	-1.66485E+00	-1.61409E+00	4.13077E-04	-2.37492E-08	±0.19
70	FN	(2)	4	3.61331E+04	-1.34903E+00	-2.94868E+00	3.47267E-04	-1.96760E-08	
71	ALN	(2)	4	3.33804E+04	-9.12019E-01	-4.84637E+00	1.97911E-04	-1.11046E-08	
72	SIN	(2)	4	6.57987E+04	-1.53565E+00	-2.64502E+00	5.01934E-04	-2.62642E-08	±0.16
73	PN	(3)	4	8.18598E+04	-1.60139E+00	-2.22616E+00	5.05120E-04	-3.89487E-08	
74	SN	(2)	4	5.81195E+04	-1.39806E+00	-2.88207E+00	3.53347E-04	-2.28622E-08	
75	NCL	(1)	4	3.97098E+04	-1.56705E+00	-1.14196E+00	4.75760E-04	-2.78019E-08	±0.64
76	TIN	(1)	4	5.68328E+04	-2.24595E+00	1.36460E+00	1.01540E-03	-9.39699E-08	
77	ZRN	(2)	4	4.40761E+04	-1.46014E+00	-2.69658E+00	1.33845E-04	-2.04661E-08	
78	NO+	(2)	4	-3.20300E+04	6.58515E-01	-1.61406E+01	4.35895E-04	-2.41454E-08	±0.11
79	NS+	(1)	4	-5.87490E+04	5.80428E-01	-1.58236E+01	5.64107E-04	-3.48183E-08	
80	LIO	(2)	4	3.87653E+04	-1.01973E+00	-4.11707E+00	2.83666E-04	-2.05044E-08	
81	FO	(2)	4	2.61197E+04	-1.34033E+00	-3.79573E+00	3.45232E-04	-1.79903E-08	±0.32
82	NAO	(2)	4	3.26015E+04	-8.09503E-01	-5.04056E+00	2.15940E-04	-1.45434E-08	
83	MGO	(2)	4	4.10915E+04	4.80940E-01	-1.57611E+01	2.75858E-04	-3.48644E-08	
84	ALO	(2)	4	6.11077E+04	-1.79019E+00	-1.35971E+00	7.68615E-04	-3.93150E-08	±0.35
85	SIO	(2)	4	9.58807E+04	-1.26716E+00	-5.98429E+00	2.61720E-04	-1.67404E-08	
86	PO	(2)	4	7.05508E+04	-1.23215E+00	-4.24066E+00	3.24391E-04	-2.61888E-08	
87	SO	(2)	4	6.22989E+04	-1.59126E+00	-2.62423E+00	4.40388E-04	-2.29940E-08	±0.20
88	CLO	(2)	4	3.20486E+04	-1.33025E+00	-3.35070E+00	3.07999E-04	-1.57804E-08	
89	KO	(2)	4	3.18934E+04	-8.52327E-01	-4.49752E+00	2.92534E-04	-2.55392E-08	
90	CAO	(2)	4	4.53278E+04	-2.48368E+00	4.66213E+00	1.24840E-03	-6.67350E-08	±0.52
91	TIO	(2)	4	8.02469E+04	-8.46602E-01	-7.67141E+00	1.51196E-04	-2.20473E-08	
92	VO	(2)	4	7.61730E+04	-1.71137E+00	-2.36139E+00	3.25571E-04	-2.10832E-08	
93	CRO	(2)	4	5.49087E+04	-1.03554E+00	-4.97677E+00	2.90615E-04	-3.71885E-08	±0.24
94	MNO	(1)	4	4.30448E+04	-1.83220E+00	1.75207E-01	7.93081E-04	-6.34611E-08	
95	FEO	(2)	4	4.93909E+04	-1.54390E+00	-2.08333E+00	3.92414E-04	-2.89557E-08	
96	NIO	(1)	4	4.36215E+04	-1.96062E+00	-1.57865E+00	5.80268E-04	-3.70799E-08	±0.32
97	ZRO	(2)	4	9.57474E+04	-2.46971E+00	1.81056E+00	1.16686E-03	-8.51671E-08	
98	WO	(3)	4	8.07011E+04	-1.64832E+00	6.20649E-01	1.02786E-06	-1.00770E-08	
99	SIO+	(1)	4	-3.76151E+04	2.15564E-02	-1.15232E+01	9.39253E-04	-6.75576E-08	±0.20
100	PO+	(1)	4	-2.60911E+04	5.63925E-01	-1.43772E+01	6.86220E-04	-5.20459E-08	
101	SO+	(1)	4	-5.76275E+04	6.89404E-01	-1.52462E+01	5.60055E-04	-3.75051E-08	
102	FEO-	(1)	4	6.39751E+04	-4.66052E+00	1.40743E+01	7.84086E-04	-6.30384E-08	±0.21
103	LIF	(2)	4	6.93946E+04	-1.25530E+00	-3.61962E+00	3.57085E-04	-2.37547E-08	
104	NAF	(2)	4	5.71445E+04	-9.75105E-01	-5.09405E+00	2.57599E-04	-1.65555E-08	
105	MGF	(2)	4	5.54581E+04	-1.08881E+00	-3.35361E+00	2.76432E-04	-1.72051E-08	±0.16
106	ALF	(2)	4	8.09027E+04	-1.09464E+00	-5.96190E+00	2.36759E-04	-1.17831E-08	
107	SIF	(2)	4	6.58544E+04	-1.03152E+00	-5.51281E+00	1.84248E-04	-1.13498E-08	
108	PF	(2)	4	5.35929E+04	-1.21444E+00	-3.84164E+00	3.33851E-04	-2.21134E-08	±0.06
109	SF	(2)	4	4.10442E+04	-1.07784E+00	-5.22937E+00	2.30560E-04	-1.39484E-08	
110	KF	(2)	4	5.92978E+04	-9.99383E-01	-4.65004E+00	3.20903E-04	-2.71810E-08	
111	CAF	(2)	4	6.33281E+04	-1.22894E+00	-2.16525E+00	4.51849E-04	-4.22645E-08	±0.48
112	MNF	(1)	4	5.30891E+04	-1.61009E+00	-5.40777E-01	6.53952E-04	-5.26395E-08	
113	NIF	(1)	4	5.10347E+04	-1.96613E+00	-1.05814E+00	4.99399E-04	-3.05380E-08	
114	LINA	(1)	4	1.00750E+04	-1.04820E+00	-2.18246E+00	6.23562E-04	-5.59401E-08	±0.41
115	MGs	(2)	4	3.41340E+04	1.12247E+00	-1.92306E+01	-9.69903E-05	-1.19684E-08	
116	ALS	(2)	4	4.38521E+04	-1.73346E+00	-1.15384E+00	8.41827E-04	-4.98671E-08	
117	SIS	(2)	4	7.44656E+04	-1.05114E+00	-6.74615E+00	1.80683E-04	-1.29591E-08	±0.09
118	PS	(2)	4	5.44686E+04	-9.40310E-01	-5.56638E+00	2.21811E-04	-2.13486E-08	
119	CAS	(2)	4	3.89498E+04	-3.02245E+00	8.78829E+00	1.75934E-03	-1.12821E-07	
120	TIS	(1)	4	4.97910E+04	-1.97079E+00	5.87719E-01	7.96752E-04	-7.21988E-08	±0.21
121	CRS	(1)	4	3.93336E+04	-1.66389E+00	-2.41348E+00	6.56029E-04	-6.58245E-08	
122	LICL	(2)	4	5.69768E+04	-1.17973E+00	-3.25999E+00	3.02996E-04	-1.99484E-08	
123	NaCL	(2)	4	4.90333E+04	-1.00447E+00	-4.10949E+00	2.39064E-04	-1.46354E-08	±0.21
124	MGCL	(2)	4	3.72258E+04	-1.04853E+00	-2.79170E+00	2.35261E-04	-1.42132E-08	
125	ALCL	(2)	4	6.01726E+04	-9.82189E-01	-5.80775E+00	1.65774E-04	-6.11193E-09	
126	SICL	(2)	4	4.46257E+04	-8.90237E-01	-5.62820E+00	1.06806E-04	-5.93045E-09	±0.21
127	PCL	(2)	4	3.68530E+04	-1.02053E+00	-4.20951E+00	2.32075E-04	-1.91252E-08	
128	KCL	(2)	4	5.08136E+04	-1.05759E+00	-3.45119E+00	3.16489E-04	-2.58507E-08	
129	CaCL	(2)	4	4.82109E+04	-1.21582E+00	-1.48229E+00	4.15570E-04	-3.83871E-08	±0.09
130	MNCL	(1)	4	4.02949E+04	-1.20835E+00	-2.47233E+00	3.83768E-04	-3.43403E-08	
131	FECL	(2)	4	3.40948E+04	-1.10689E+00	-3.45623E+00	3.15583E-04	-2.79431E-08	
132	ALCLF	(2)	4	1.22290E+05	-1.60845E+00	-1.43672E+01	3.72486E-04	-1.98494E-08	±0.21
133	ALCLF2	(2)	4	1.93130E+05	-1.90100E+00	-3.00532E+01	6.68641E-04	-3.72955E-08	
134	ALOCL	(2)	4	1.25832E+05	-1.22293E+00	-2.13795E+01	4.32404E-04	-2.48564E-08	
135	ALCL2	(2)	4	1.02155E+05	-1.52215E+00	-1.47912E+01	3.11627E-04	-1.55870E-08	±0.21
136	ALCL2F	(2)	4	1.73107E+05	-1.75884E+00	-2.99858E+01	5.88482E-04	-3.19785E-08	
137	ALCL3	(2)	4	1.53343E+05	-1.63768E+00	-3.10467E+01	5.15664E-04	-2.70655E-08	
138	ALOF	(2)	4	1.48870E+05	-1.37368E+00	-2.11531E+01	5.14777E-04	-3.02703E-08	±0.21
139	ALF2	(2)	4	1.41911E+05	-1.73579E+00	-1.51383E+01	4.46882E-04	-2.48292E-08	
140	ALF2O	(2)	4	2.21740E+05	-1.82980E+00	-3.07239E+01	6.62316E-04	-3.43978E-08	
141	ALF3	(2)	4	2.13367E+05	-2.05187E+00	-3.09713E+01	7.51218E-04	-4.27263E-08	±0.21
142	NaALF4	(2)	4	3.11975E+05	-1.70831E+00	-6.29805E+01	1.04098E-03	-6.36455E-08	
143	ALOH	(2)	4	1.16643E+05	-3.49644E+00	-4.46506E+00	1.06912E-03	-5.57776E-08	
144	HALO	(2)	4	9.12736E+04	-2.61419E+00	-1.05748E+01	9.35627E-04	-5.30554E-08	±0.21
145	ALO2H	(2)	4	1.80386E+05	-3.42803E+00	-1.98423E+01	1.08737E-03	-5.85448E-08	
146	ALO2	(2)	4	1.09779E+05	-1.02797E+00	-2.18620E+01	3.81694E-04	-2.17081E-08	
147	AL2CL6	(2)	4	3.22505E+05	-1.31147E+00	-9.38002E+01	1.04696E-03	-5.50934E-08	±0.21
148	(ALF3)2	(2)	4	4.53134E+05	-2.49740E+00	-9.29287E+01	1.65542E-03	-9.43913E-08	
149	AL2O	(2)	4	1.26583E+05	-8.80611E-01	-2.31266E+01	3.35552E-04	-2.01455E-08	
150	AL2O2	(2)	4	1.86515E+05	-1.16080E+00	-3.73783E+01	4.58680E-04	-2.85416E-08	±0.21
151	ALC	(2)	4	4.27037E+04	-9.01708E-01	-6.11242E+00	1.89365E-04	-1.16993E-08	

Table D.1. continued

	molecule	source	fit	a_0	a_1	a_2	a_3	a_4	σ [kJ/mol]
152	CFCLO	(2)	4	1.90938E+05	-3.22705E+00	-2.35717E+01	1.11849E-03	-6.39039E-08	
153	CCLF3	(2)	4	2.13906E+05	-3.54875E+00	-3.96988E+01	1.43940E-03	-8.20706E-08	
154	CCLN	(2)	4	1.40543E+05	-2.11454E+00	-1.53492E+01	7.62648E-04	-4.01053E-08	
155	CCLO	(2)	4	1.37740E+05	-2.24974E+00	-1.16913E+01	5.79713E-04	-3.34800E-08	
156	CCL2	(2)	4	8.60873E+04	-2.48374E+00	-1.09431E+01	1.06378E-03	-6.44276E-08	
157	CCL2F2	(2)	4	1.92998E+05	-3.11752E+00	-4.12502E+01	1.26253E-03	-7.15634E-08	
158	CCL2O	(2)	4	1.71174E+05	-2.84335E+00	-2.58668E+01	9.57251E-04	-5.42207E-08	
159	CCL3	(2)	4	1.19890E+05	-2.26317E+00	-2.84785E+01	7.47794E-04	-4.19363E-08	
160	CCL3F	(2)	4	1.73694E+05	-2.70048E+00	-4.35096E+01	1.08834E-03	-6.11304E-08	
161	CCL4	(2)	4	1.55609E+05	-2.32138E+00	-4.65765E+01	9.26363E-04	-5.13389E-08	
162	CFN	(2)	4	1.47744E+05	-2.28893E+00	-1.48210E+01	7.78781E-04	-4.54674E-08	
163	CFO	(2)	4	1.45732E+05	-2.64677E+00	-1.06264E+01	7.52624E-04	-4.40212E-08	
164	CF2	(2)	4	1.26653E+05	-2.42892E+00	-1.27598E+01	6.63277E-04	-3.52162E-08	
165	CF2O	(2)	4	2.11342E+05	-3.64773E+00	-2.23878E+01	1.28865E-03	-7.39551E-08	
166	CF3	(2)	4	1.70744E+05	-3.29984E+00	-2.36643E+01	1.18949E-03	-6.85856E-08	
167	CF4	(2)	4	2.35878E+05	-4.02787E+00	-3.88452E+01	1.63109E-03	-9.33349E-08	
168	CF4O	(2)	4	2.45540E+05	-4.32540E+00	-4.90057E+01	1.74830E-03	-1.00809E-07	
169	CF8S	(2)	4	4.01621E+05	-5.62795E+00	-1.12184E+02	2.96591E-03	-1.71636E-07	
170	CHCL	(2)	4	8.59646E+04	-3.36069E+00	-2.98598E+00	1.30814E-03	-7.93219E-08	
171	CHCL3	(2)	4	1.67582E+05	-4.40936E+00	-2.87325E+01	1.55281E-03	-8.30020E-08	
172	CHF	(2)	4	1.05972E+05	-3.52442E+00	-2.18932E+00	9.08193E-04	-3.81824E-08	
173	CHFO	(2)	4	1.96095E+05	-4.86167E+00	-1.05290E+01	1.58514E-03	-8.66741E-08	
174	CHF3	(2)	4	2.23689E+05	-5.62269E+00	-2.32634E+01	2.04873E-03	-1.12400E-07	
175	HCN	(2)	4	1.52208E+05	-3.20440E+00	-6.26703E+00	9.84333E-04	-5.17826E-08	
176	CHNO	(2)	4	2.10425E+05	-4.30285E+00	-1.42930E+01	1.33044E-03	-7.24817E-08	
177	HCO	(2)	4	1.36277E+05	-3.47855E+00	-2.64678E+00	9.51657E-04	-5.11327E-08	
178	PCH	(2)	4	1.31799E+05	-2.71934E+00	-9.15248E+00	9.09960E-04	-5.27936E-08	
179	CH2	(2)	4	9.11697E+04	-3.73503E+00	1.06433E+00	9.79096E-04	-4.94662E-08	
180	CH2CL2	(2)	4	1.77820E+05	-6.22408E+00	-1.39323E+01	2.07847E-03	-1.08928E-07	
181	CH2CLF	(2)	4	1.92725E+05	-6.60928E+00	-1.14750E+01	2.23384E-03	-1.18076E-07	
182	CH2F2	(2)	4	2.10297E+05	-7.06755E+00	-9.97156E+00	2.41754E-03	-1.28823E-07	
183	H2CO	(2)	4	1.81076E+05	-5.95395E+00	-1.27437E+00	1.86009E-03	-9.86705E-08	
184	CH3	(2)	4	1.45821E+05	-5.79871E+00	6.20786E-01	1.63241E-03	-8.09944E-08	
185	CH3CL	(2)	4	1.87616E+05	-7.71117E+00	-2.44843E+00	2.48088E-03	-1.27804E-07	
186	CH3F	(2)	4	2.00605E+05	-8.14272E+00	-3.51338E-01	2.65388E-03	-1.37902E-07	
187	CH4	(2)	4	1.97846E+05	-8.83168E+00	5.27931E+00	2.75677E-03	-1.39667E-07	
188	KCN	(2)	4	1.43802E+05	-1.56464E+00	-1.62289E+01	6.28550E-04	-4.88291E-08	
189	NACN	(2)	4	1.44245E+05	-1.51417E+00	-1.69281E+01	5.54291E-04	-3.75330E-08	
190	CNO	(2)	4	1.53330E+05	-2.32714E+00	-1.40991E+01	8.24890E-04	-4.94008E-08	
191	CNN	(2)	4	1.29031E+05	-2.32368E+00	-1.30660E+01	7.56603E-04	-4.46197E-08	
192	NCN	(2)	4	1.42632E+05	-1.89963E+00	-1.66755E+01	7.12812E-04	-4.50701E-08	
193	COS	(2)	4	1.65625E+05	-2.33683E+00	-1.59211E+01	8.52197E-04	-4.93878E-08	
194	CO2	(2)	4	1.92841E+05	-2.77259E+00	-1.43745E+01	9.55278E-04	-5.32731E-08	
195	CS2	(2)	4	1.38365E+05	-1.96405E+00	-1.84236E+01	7.55600E-04	-4.64468E-08	
196	Si2C	(2)	4	1.29802E+05	-1.23194E+00	-2.25394E+01	2.98881E-04	-8.01112E-09	
197	C2CL2	(2)	4	1.75870E+05	-2.18122E+00	-3.11831E+01	8.66319E-04	-4.94706E-08	
198	C2CL4	(2)	4	2.31604E+05	-3.17447E+00	-5.62904E+01	1.39975E-03	-7.92920E-08	
199	C2CL6	(2)	4	2.75656E+05	-2.67884E+00	-9.31000E+01	1.62848E-03	-9.62481E-08	
200	C2F2	(2)	4	1.88465E+05	-2.61936E+00	-3.04836E+01	1.10048E-03	-6.47411E-08	
201	C2F3N	(2)	4	3.16642E+05	-4.37972E+00	-5.05423E+01	1.90213E-03	-1.10625E-07	
202	C2F4	(2)	4	2.88983E+05	-4.34112E+00	-5.12957E+01	1.89071E-03	-1.08751E-07	
203	C2F6	(2)	4	3.90525E+05	-5.02635E+00	-8.18595E+01	2.39658E-03	-1.39507E-07	
204	C2H	(2)	4	1.40482E+05	-3.02583E+00	-7.31209E+00	9.07164E-04	-4.24041E-08	
205	C2HCL	(2)	4	1.86705E+05	-3.30242E+00	-2.14329E+01	1.14526E-03	-6.18882E-08	
206	C2HF	(2)	4	1.92266E+05	-3.45036E+00	-2.10247E+01	1.21002E-03	-6.58528E-08	
207	C2H2	(2)	4	1.96460E+05	-4.51429E+00	-1.24279E+01	1.51777E-03	-7.45421E-08	
208	C2H4	(2)	4	2.68664E+05	-9.52900E+00	-5.66852E+00	3.19357E-03	-1.66353E-07	
209	C2H4O	(2)	4	3.11083E+05	-1.07569E+01	-1.47028E+01	3.83108E-03	-2.03086E-07	
210	(KCN)2	(2)	4	3.08037E+05	-2.10431E+00	-5.53302E+01	1.24792E-03	-9.71373E-08	
211	C2N	(2)	4	1.62054E+05	-1.35960E+00	-2.01803E+01	5.27257E-04	-3.56032E-08	
212	C2N2	(2)	4	2.48284E+05	-2.84562E+00	-2.77626E+01	1.12826E-03	-6.72780E-08	
213	(NACN)2	(2)	4	3.12517E+05	-2.06576E+00	-5.71243E+01	1.12359E-03	-7.59581E-08	
214	SiC2	(2)	4	1.52254E+05	-1.44598E+00	-2.06243E+01	3.96365E-04	-1.47426E-08	
215	C2O	(2)	4	1.67387E+05	-2.23683E+00	-1.51504E+01	8.99197E-04	-5.29613E-08	
216	C3	(2)	4	1.59040E+05	-3.39362E+00	-6.27788E+00	7.26049E-04	-3.93526E-08	
217	C3O2	(2)	4	3.29014E+05	-3.76315E+00	-3.89142E+01	1.61372E-03	-9.42755E-08	
218	C4	(2)	4	2.27422E+05	-3.01949E+00	-2.96110E+01	1.22636E-03	-7.50022E-08	
219	C4N2	(2)	4	3.93522E+05	-3.91501E+00	-5.34656E+01	1.83065E-03	-1.09136E-07	
220	Ni(CO)4	(2)	4	5.87870E+05	-4.40774E+00	-9.79396E+01	2.23682E-03	-1.28130E-07	
221	C5	(2)	4	3.12497E+05	-3.75947E+00	-4.24806E+01	1.65606E-03	-1.00477E-07	
222	Fe(CO)5	(2)	4	7.17355E+05	-5.75404E+00	-1.25442E+02	3.20703E-03	-1.97629E-07	
223	CACL2	(2)	4	1.06957E+05	-1.16258E+00	-1.59094E+01	5.56653E-04	-5.10835E-08	
224	CAF2	(2)	4	1.34440E+05	-1.63884E+00	-1.33684E+01	6.08477E-04	-5.57178E-08	
225	CAOH	(2)	4	1.00318E+05	-2.24867E+00	-8.95653E+00	7.99317E-04	-5.91430E-08	
226	CA(OH)2	(2)	4	2.06312E+05	-3.67249E+00	-2.69984E+01	1.34817E-03	-8.94131E-08	
227	CA2	(2)	4	8.98860E+04	-2.17354E+00	8.56959E+00	2.38871E-04	-5.44705E-08	
228	CLF	(2)	4	3.00064E+02	-8.36540E-01	-7.31349E+00	-1.31237E-04	3.44016E-08	
229	MGCLF	(3)	4	1.09651E+05	-2.12483E+00	-1.15990E+01	6.93435E-04	-4.54707E-08	±0.29
230	CLFO2S	(2)	4	1.83698E+05	-3.26458E+00	-4.10926E+01	1.34145E-03	-7.82252E-08	
231	CLFO3	(2)	4	1.15936E+05	-3.60277E+00	-4.10467E+01	1.47203E-03	-8.41633E-08	
232	PCLF2O	(2)	4	2.17848E+05	-3.27293E+00	-4.03948E+01	1.39034E-03	-8.63368E-08	
233	CLF3	(2)	4	6.19362E+04	-2.04133E+00	-3.04553E+01	7.18188E-04	-3.98530E-08	
234	CLF3SI	(2)	4	2.55495E+05	-2.28475E+00	-4.61308E+01	9.63001E-04	-5.60723E-08	

Table D.1. continued

	molecule	source	fit	a_0	a_1	a_2	a_3	a_4	σ [kJ/mol]
235	CLF5	(2)	4	9.06733E+04	-2.37569E+00	-6.35600E+01	1.21159E-03	-6.79411E-08	
236	CLF5S	(2)	4	2.20114E+05	-3.43455E+00	-7.62559E+01	1.78547E-03	-1.02728E-07	
237	CHCLF2	(2)	4	2.02940E+05	-4.98328E+00	-2.54851E+01	1.80645E-03	-9.85202E-08	
238	CHCL2F	(2)	4	1.84147E+05	-4.76213E+00	-2.60846E+01	1.70179E-03	-9.19490E-08	
239	OHCL	(2)	4	7.88945E+04	-3.23264E+00	-3.54939E+00	7.60371E-04	-3.65966E-08	
240	SIH3CL	(2)	4	1.63156E+05	-5.64837E+00	-1.53353E+01	1.97093E-03	-1.06607E-07	
241	NOCL	(2)	4	9.46165E+04	-2.30608E+00	-1.13062E+01	6.58320E-04	-3.32492E-08	
242	NO2CL	(2)	4	1.29237E+05	-3.13502E+00	-2.44274E+01	1.10857E-03	-6.39755E-08	
243	NICL	(2)	4	4.40373E+04	-1.38673E+00	-2.34884E+00	2.89903E-04	-1.61249E-09	
244	TIOCL	(2)	4	1.30707E+05	-1.13481E+00	-2.21613E+01	4.00263E-04	-4.20741E-08	
245	CLO2	(2)	4	6.14263E+04	-2.26719E+00	-1.31718E+01	6.67536E-04	-3.57808E-08	
246	CLO3	(2)	4	8.05675E+04	-2.76999E+00	-2.80064E+01	1.00255E-03	-5.81589E-08	
247	SCL	(2)	4	2.88151E+04	-9.92038E-01	-4.99727E+00	1.72279E-04	-9.79941E-09	
248	CLS2	(2)	4	7.13616E+04	-1.73435E+00	-1.40073E+01	4.23268E-04	-2.72832E-08	
249	TICL	(2)	4	5.29650E+04	-4.15122E-01	-9.18790E+00	2.51303E-04	-3.13050E-08	
250	CL2FOP	(2)	4	1.98353E+05	-2.51772E+00	-4.35669E+01	1.08964E-03	-6.86642E-08	
251	FECL2	(2)	4	9.56812E+04	-1.72272E+00	-1.43864E+01	7.07229E-04	-4.70003E-08	
252	SIH2CL2	(2)	4	1.73296E+05	-4.46958E+00	-2.46408E+01	1.58428E-03	-8.60852E-08	
253	(KCL)2	(2)	4	1.24565E+05	-1.11068E+00	-2.86471E+01	5.40359E-04	-5.16089E-08	
254	MGCL2	(2)	4	9.38138E+04	-1.01412E+00	-1.77315E+01	3.47071E-04	-2.13014E-08	
255	(NACL)2	(2)	4	1.22803E+05	-1.03283E+00	-3.08613E+01	4.04266E-04	-2.98327E-08	
256	NICL2	(2)	4	8.94776E+04	-1.27282E+00	-1.80400E+01	4.69046E-04	-2.30791E-08	
257	CLOCL	(2)	4	4.89488E+04	-1.83971E+00	-1.45206E+01	4.40040E-04	-2.40923E-08	
258	CLCLO	(2)	4	4.78510E+04	-1.86730E+00	-1.34628E+01	4.45691E-04	-2.42698E-08	
259	TIOCL2	(2)	4	1.81444E+05	-1.45251E+00	-3.30403E+01	4.85880E-04	-4.60617E-08	
260	CLO2CL	(2)	4	7.27118E+04	-1.92875E+00	-3.00779E+01	6.67081E-04	-3.78596E-08	
261	CLOCLO	(2)	4	6.75070E+04	-1.81937E+00	-2.97451E+01	6.22684E-04	-3.52097E-08	
262	CL2O2S	(2)	4	1.64512E+05	-2.96202E+00	-4.26690E+01	1.20365E-03	-6.96762E-08	
263	CL2S	(2)	4	6.41542E+04	-1.79873E+00	-1.43855E+01	4.16335E-04	-2.40025E-08	
264	SICL2	(2)	4	1.03214E+05	-1.47585E+00	-1.65100E+01	2.62899E-04	-1.47153E-08	
265	TICL2	(2)	4	1.14417E+05	-1.12113E+00	-2.07743E+01	3.93107E-04	-3.04261E-08	
266	CL3FSI	(2)	4	2.08254E+05	-1.72278E+00	-4.76916E+01	6.99317E-04	-3.95355E-08	
267	FECL3	(2)	4	1.23755E+05	-1.74468E+00	-2.86633E+01	5.51617E-04	-3.99323E-08	
268	SIHCL3	(2)	4	1.83103E+05	-3.17555E+00	-3.57354E+01	1.15601E-03	-6.32103E-08	
269	PCL3O	(2)	4	1.78669E+05	-2.19581E+00	-4.56496E+01	9.44866E-04	-5.97254E-08	
270	PCL3	(2)	4	1.16154E+05	-1.63661E+00	-3.12472E+01	5.59237E-04	-3.74063E-08	
271	PCL3S	(2)	4	1.60564E+05	-1.82383E+00	-4.74176E+01	8.15721E-04	-5.42914E-08	
272	SICL3	(2)	4	1.44470E+05	-1.63614E+00	-3.10721E+01	4.87187E-04	-2.76095E-08	
273	TICL3	(2)	4	1.65407E+05	-1.17152E+00	-3.59417E+01	4.79809E-04	-4.52503E-08	
274	FE2CL4	(2)	4	2.09744E+05	-1.98373E+00	-5.45277E+01	1.11142E-03	-7.94145E-08	
275	MG2CL4	(2)	4	2.08330E+05	-9.20014E-01	-5.90953E+01	6.54798E-04	-4.03856E-08	
276	SICL4	(2)	4	1.91821E+05	-1.78172E+00	-4.86279E+01	6.94653E-04	-3.85481E-08	
277	TICL4	(2)	4	2.06815E+05	-1.48911E+00	-4.91188E+01	6.15354E-04	-5.09373E-08	
278	CL5P	(2)	4	1.53958E+05	-1.79649E+00	-6.39289E+01	9.27158E-04	-5.66602E-08	
279	(FECL3)2	(2)	4	2.65935E+05	-1.39877E+00	-8.94974E+01	1.06886E-03	-7.78799E-08	
280	CRN	(2)	4	4.36898E+04	-1.03724E+00	-5.05499E+00	2.70519E-04	-3.48241E-08	
281	CRO2	(2)	4	1.16423E+05	-1.83062E+00	-1.57443E+01	5.74372E-04	-5.63916E-08	
282	CRO3	(2)	4	1.72567E+05	-2.17354E+00	-3.36233E+01	8.96893E-04	-7.56414E-08	
283	FEF	(2)	4	5.34439E+04	-1.46949E+00	-2.47226E+00	3.78138E-04	-3.15170E-08	
284	FHO	(2)	4	7.67186E+04	-3.33236E+00	-3.38274E+00	8.32959E-04	-4.24195E-08	
285	FHO3S	(2)	4	2.48576E+05	-4.94186E+00	-4.44159E+01	1.97377E-03	-1.10427E-07	
286	SIH3F	(2)	4	1.86320E+05	-5.90469E+00	-1.45216E+01	2.08998E-03	-1.14032E-07	
287	FNO	(2)	4	1.03702E+05	-2.41030E+00	-1.15898E+01	6.79601E-04	-3.96747E-08	
288	FNO2	(2)	4	1.38729E+05	-3.30791E+00	-2.40842E+01	1.19276E-03	-6.93140E-08	
289	FNO3	(2)	4	1.54375E+05	-3.54603E+00	-3.82963E+01	1.47776E-03	-8.64485E-08	
290	TIOF	(2)	4	1.48405E+05	-1.23406E+00	-2.24387E+01	4.66241E-04	-4.66377E-08	
291	OFO	(2)	4	2.36085E+04	-1.78598E+00	-1.64167E+01	5.37092E-04	-3.34602E-08	
292	FOO	(2)	4	6.59465E+04	-2.08776E+00	-1.32391E+01	5.72512E-04	-2.92066E-08	
293	PSF	(2)	4	1.01214E+05	-1.95266E+00	-1.30959E+01	5.81119E-04	-4.33475E-08	
294	TIF	(2)	4	7.45769E+04	-4.13878E-01	-9.95679E+00	2.70397E-04	-3.38157E-08	
295	FEF2	(2)	4	1.15879E+05	-6.55994E-01	-2.42023E+01	3.90037E-04	-3.46358E-08	
296	H2F2	(2)	4	1.39214E+05	-4.78857E+00	-6.09204E+00	1.32945E-03	-6.44300E-08	
297	SIH2F2	(2)	4	2.19730E+05	-4.96388E+00	-2.31174E+01	1.81390E-03	-1.00416E-07	
298	K2F2	(2)	4	1.44056E+05	-9.68810E-01	-3.21695E+01	5.48486E-04	-5.37565E-08	
299	MGF2	(2)	4	1.23834E+05	-1.61314E+00	-1.48912E+01	4.41170E-04	-2.82419E-08	
300	F2N	(2)	4	7.03763E+04	-2.22233E+00	-1.24082E+01	6.40864E-04	-3.77976E-08	
301	F2N2(cis)	(2)	4	1.23923E+05	-3.09351E+00	-2.43660E+01	1.14398E-03	-6.75220E-08	
302	F2N2(trans)	(2)	4	1.22423E+05	-2.95398E+00	-2.48398E+01	1.08171E-03	-6.36611E-08	
303	NA2F2	(2)	4	1.46476E+05	-1.22480E+00	-3.29465E+01	5.38240E-04	-3.92171E-08	
304	F2O	(2)	4	4.56834E+04	-1.99946E+00	-1.50778E+01	5.53675E-04	-3.20511E-08	
305	F2OS	(2)	4	1.47290E+05	-2.64335E+00	-2.76184E+01	9.66159E-04	-5.74202E-08	
306	SIOF2	(2)	4	2.18924E+05	-2.71013E+00	-2.81813E+01	9.57418E-04	-5.69004E-08	
307	TIOF2	(2)	4	2.16993E+05	-1.78398E+00	-3.40307E+01	6.69345E-04	-5.81947E-08	
308	F2O2	(2)	4	7.63441E+04	-2.01719E+00	-3.09876E+01	7.48004E-04	-4.37641E-08	
309	F2O2S	(2)	4	2.02919E+05	-3.62466E+00	-4.04633E+01	1.49805E-03	-8.77673E-08	
310	PF2	(2)	4	1.15475E+05	-1.84418E+00	-1.44554E+01	5.33554E-04	-3.79932E-08	
311	F2S	(2)	4	8.76493E+04	-2.03958E+00	-1.44050E+01	5.63785E-04	-3.41360E-08	
312	F2S2(1)	(2)	4	1.25836E+05	-1.93593E+00	-3.11273E+01	7.16160E-04	-4.52054E-08	
313	F2S2(2)	(2)	4	1.33613E+05	-2.14372E+00	-2.99840E+01	7.92571E-04	-4.95271E-08	
314	SIF2	(2)	4	1.43548E+05	-1.80429E+00	-1.60725E+01	4.45430E-04	-2.71863E-08	
315	TIF2	(2)	4	1.58647E+05	-1.14963E+00	-2.20021E+01	4.64208E-04	-3.61556E-08	
316	FEF3	(2)	4	1.76929E+05	-2.21826E+00	-2.85224E+01	8.14849E-04	-5.73470E-08	
317	SIHF3	(2)	4	2.52662E+05	-3.88364E+00	-3.36148E+01	1.48930E-03	-8.41141E-08	
318	F3H3	(2)	4	2.11700E+05	-5.74775E+00	-2.72569E+01	2.01458E-03	-1.01720E-07	

Table D.1. continued

	molecule	source	fit	a_0	a_1	a_2	a_3	a_4	σ [kJ/mol]
319	NF3	(2)	4	1.00865E+05	-2.71701E+00	-2.73662E+01	1.00793E-03	-5.86664E-08	
320	F3NO	(2)	4	1.34498E+05	-3.24725E+00	-4.14236E+01	1.36643E-03	-7.89717E-08	
321	POF3	(2)	4	2.46952E+05	-3.19634E+00	-4.21379E+01	1.38688E-03	-8.68175E-08	
322	PF3	(2)	4	1.81552E+05	-2.27768E+00	-2.98605E+01	8.79252E-04	-5.78887E-08	
323	PSF3	(2)	4	2.20930E+05	-2.69474E+00	-4.46472E+01	1.21655E-03	-7.92411E-08	
324	SF3	(2)	4	1.22080E+05	-2.07877E+00	-3.00909E+01	7.62186E-04	-4.52878E-08	
325	SIF3	(2)	4	2.12906E+05	-2.21091E+00	-2.97382E+01	7.79319E-04	-4.63991E-08	
326	TIF3	(2)	4	2.28447E+05	-1.33046E+00	-3.59961E+01	6.24937E-04	-5.58901E-08	
327	F4H4	(2)	4	2.83870E+05	-7.00233E+00	-4.50322E+01	2.68824E-03	-1.35754E-07	
328	MG2F4	(2)	4	2.80117E+05	-1.37158E+00	-6.35237E+01	9.45593E-04	-6.03845E-08	
329	F4N2	(2)	4	1.52312E+05	-3.66985E+00	-5.45367E+01	1.74012E-03	-1.01843E-07	
330	F4S	(2)	4	1.62897E+05	-2.47609E+00	-4.53144E+01	1.09253E-03	-6.41588E-08	
331	SIF4	(2)	4	2.86101E+05	-2.74301E+00	-4.56600E+01	1.15125E-03	-6.72655E-08	
332	TIF4	(2)	4	2.81532E+05	-1.76448E+00	-4.95271E+01	8.33250E-04	-6.65409E-08	
333	F5H5	(2)	4	3.56182E+05	-8.25550E+00	-6.30002E+01	3.36125E-03	-1.69746E-07	
334	PF5	(2)	4	2.77058E+05	-3.30871E+00	-5.87283E+01	1.62038E-03	-9.97622E-08	
335	SF5	(2)	4	1.89917E+05	-2.87408E+00	-6.15867E+01	1.40686E-03	-7.77492E-08	
336	F6H6	(2)	4	4.29942E+05	-9.50936E+00	-8.11142E+01	4.03458E-03	-2.03758E-07	
337	F6S	(2)	4	2.36854E+05	-3.89097E+00	-7.61409E+01	1.98028E-03	-1.14520E-07	
338	F7H7	(2)	4	5.00930E+05	-1.07630E+01	-9.93459E+01	4.70781E-03	-2.37764E-07	
339	F10S2	(2)	4	4.10025E+05	-4.93413E+00	-1.54896E+02	3.28125E-03	-1.95093E-07	
340	FE(OH)2	(2)	4	2.01370E+05	-3.53207E+00	-3.17623E+01	1.44539E-03	-9.15799E-08	
341	FES	(2)	4	3.83484E+04	-1.48336E+00	-2.02949E+00	3.73194E-04	-2.90273E-08	
342	KOH	(2)	4	9.43809E+04	-1.86933E+00	-1.18848E+01	5.97392E-04	-4.10952E-08	
343	MGOH	(2)	4	9.31677E+04	-2.13709E+00	-1.00212E+01	6.29804E-04	-3.46633E-08	
344	HNO	(2)	4	1.00185E+05	-3.46381E+00	-2.62213E+00	9.45251E-04	-5.14088E-08	
345	HNO2(trans)	(2)	4	1.51533E+05	-4.06412E+00	-1.49544E+01	1.28552E-03	-6.98861E-08	
346	HNO2(cis)	(2)	4	1.51266E+05	-4.14727E+00	-1.44178E+01	1.32602E-03	-7.23681E-08	
347	HNO3	(2)	4	1.88003E+05	-5.14751E+00	-2.56506E+01	1.85951E-03	-1.02730E-07	
348	NAOH	(2)	4	9.24108E+04	-1.78733E+00	-1.25557E+01	5.09792E-04	-2.90198E-08	
349	HO2	(2)	4	8.50031E+04	-3.52346E+00	-2.13021E+00	9.46132E-04	-4.56754E-08	
350	(KOH)2	(2)	4	2.11285E+05	-5.28939E+00	-3.16500E+01	2.06846E-03	-1.29915E-07	
351	MG(OH)2	(2)	4	1.98022E+05	-3.71156E+00	-2.82448E+01	1.20500E-03	-6.33156E-08	
352	NH2	(2)	4	8.52744E+04	-4.06091E+00	4.01425E+00	9.46257E-04	-3.84480E-08	
353	N2H2	(2)	4	1.39105E+05	-5.69139E+00	-1.43727E+00	1.74602E-03	-9.21166E-08	
354	(NAOH)2	(2)	4	2.10035E+05	-5.22073E+00	-3.29843E+01	1.93564E-03	-1.08319E-07	
355	H2O	(2)	4	1.10336E+05	-4.17836E+00	3.17447E+00	9.40647E-04	-4.04825E-08	
356	H2SO4	(2)	4	2.92715E+05	-6.31402E+00	-4.94951E+01	2.47825E-03	-1.34895E-07	
357	PH2	(3)	4	7.43018E+04	-3.79856E+00	3.13658E+00	1.11063E-03	-6.69289E-08	±0.30
358	H2S	(2)	4	8.71685E+04	-4.03150E+00	3.16992E+00	1.08827E-03	-5.46714E-08	
359	NH3	(2)	4	1.39343E+05	-6.39532E+00	4.95981E+00	1.81530E-03	-8.85794E-08	
360	PH3	(3)	4	1.12196E+05	-6.48813E+00	6.41757E+00	2.31993E-03	-1.38454E-07	±0.84
361	N2H4	(2)	4	2.05065E+05	-8.41764E+00	-9.25261E+00	2.76895E-03	-1.41252E-07	
362	SiH4	(2)	4	1.53342E+05	-6.67420E+00	-8.19009E+00	2.30166E-03	-1.23954E-07	
363	K2SO4	(2)	4	3.05748E+05	-3.06275E+00	-7.37039E+01	1.89452E-03	-1.35325E-07	
364	MGN	(2)	4	3.95834E+04	-9.35352E-01	-3.21109E+00	2.44788E-04	-1.73353E-08	
365	NO2	(2)	4	1.12149E+05	-2.82494E+00	-1.02040E+01	8.36977E-04	-4.92832E-08	
366	NO3	(2)	4	1.37605E+05	-3.09452E+00	-2.67985E+01	1.16486E-03	-6.98024E-08	
367	Si2N	(2)	4	1.17165E+05	-8.68671E-01	-2.26653E+01	2.85125E-04	-2.25874E-08	
368	VN	(2)	4	5.55604E+04	-1.50384E+00	-2.51248E+00	3.12180E-04	-2.07231E-08	
369	N2O	(2)	4	1.33220E+05	-2.55801E+00	-1.34773E+01	8.95715E-04	-5.25572E-08	
370	N2O3	(2)	4	1.92788E+05	-3.95041E+00	-3.28349E+01	1.42328E-03	-8.40108E-08	
371	N2O4	(2)	4	2.31628E+05	-4.45135E+00	-4.99813E+01	1.97018E-03	-1.15438E-07	
372	N2O5	(2)	4	2.61545E+05	-3.82145E+00	-6.84940E+01	1.68295E-03	-1.03478E-07	
373	N3	(2)	4	1.20247E+05	-2.18705E+00	-1.41755E+01	8.06863E-04	-4.95109E-08	
374	Na2SO4	(2)	4	3.02895E+05	-3.06250E+00	-7.44553E+01	1.78682E-03	-1.15155E-07	
375	NIS	(2)	4	4.18066E+04	-1.12121E+00	-4.58805E+00	2.80362E-04	-1.84292E-08	
376	S2O	(3)	4	1.02446E+05	-3.26805E+00	-6.28324E+00	1.20799E-03	-8.12174E-08	±0.68
377	PO2	(2)	4	1.35389E+05	-1.99595E+00	-1.51727E+01	6.07336E-04	-4.40557E-08	
378	SO2	(3)	4	1.27905E+05	-3.68505E+00	-5.04460E+00	1.38250E-03	-8.66752E-08	±0.71
379	SIO2	(2)	4	1.50469E+05	-1.70934E+00	-2.06909E+01	5.98301E-04	-3.73582E-08	
380	TiO2	(2)	4	1.53310E+05	-1.71501E+00	-1.83346E+01	4.42822E-04	-4.38786E-08	
381	VO2	(2)	4	1.49343E+05	-2.38363E+00	-1.34037E+01	6.09469E-04	-4.47612E-08	
382	O3	(2)	4	7.22556E+04	-2.27501E+00	-1.49294E+01	6.96886E-04	-3.98060E-08	
383	SO3	(2)	4	1.70155E+05	-3.28748E+00	-2.67274E+01	1.19646E-03	-7.14017E-08	
384	P4O6	(2)	4	5.98101E+05	-3.54488E+00	-1.32696E+02	2.65818E-03	-1.88796E-07	
385	P4O10	(2)	4	8.00437E+05	-6.82807E+00	-1.83224E+02	4.55620E-03	-2.97341E-07	
386	P4	(2)	4	1.44951E+05	-1.09878E+00	-3.81131E+01	6.02469E-04	-6.71559E-08	
387	P4S3	(2)	4	2.70471E+05	1.44929E-01	-1.02107E+02	5.51609E-04	-7.07566E-08	
388	S3	(2)	4	8.24408E+04	-2.16199E+00	-1.43603E+01	6.87235E-04	-3.11038E-08	
389	S4	(2)	4	1.15330E+05	-2.06965E+00	-3.04642E+01	8.29662E-04	-4.52606E-08	
390	S5	(2)	4	1.53154E+05	-1.79067E+00	-5.29222E+01	8.75798E-04	-5.78149E-08	
391	S6	(2)	4	1.87448E+05	-1.66239E+00	-6.86786E+01	9.93907E-04	-6.87473E-08	
392	S7	(2)	4	2.19371E+05	-1.74827E+00	-8.20709E+01	1.24279E-03	-7.81268E-08	
393	S8	(2)	4	2.54397E+05	-1.53788E+00	-1.01133E+02	1.32959E-03	-9.40614E-08	
394	Si3	(2)	4	8.58689E+04	-3.61657E-01	-2.62701E+01	3.21839E-05	-1.58542E-09	
395	AL+	(2)	4	-6.93849E+04	2.46517E+00	-1.57629E+01	1.61344E-05	-1.16897E-09	
396	AL-	(2)	4	5.13593E+03	-2.58172E+00	1.50838E+01	1.46142E-04	-1.02707E-08	
397	ALCL+	(2)	4	-4.89736E+04	1.41100E+00	-1.88107E+01	1.88111E-04	-2.10058E-09	
398	ALCLF+	(2)	4	3.10713E+04	1.31559E+00	-3.42379E+01	4.07261E-04	-2.25287E-08	
399	ALCL2+	(2)	4	1.14390E+04	1.39405E+00	-3.43748E+01	3.49853E-04	-1.84798E-08	
400	ALCL2-	(2)	4	1.25580E+05	-3.94628E+00	-1.48134E+00	2.81381E-04	-1.37037E-08	
401	ALF+	(2)	4	-3.37054E+04	8.43636E-01	-1.59807E+01	6.39666E-04	-3.27819E-08	
402	ALF2+	(2)	4	4.81723E+04	1.17093E+00	-3.51082E+01	4.87283E-04	-2.78056E-08	
403	ALF2-	(2)	4	1.66847E+05	-4.16406E+00	-1.87501E+00	4.24641E-04	-2.39215E-08	

Table D.1. continued

	molecule	source	fit	a_0	a_1	a_2	a_3	a_4	σ [kJ/mol]
404	ALF2O-	(2)	4	2.45136E+05	-4.40725E+00	-1.69695E+01	7.08766E-04	-4.12110E-08	
405	ALF4-	(2)	4	3.16357E+05	-4.40893E+00	-3.48436E+01	8.87754E-04	-5.06459E-08	
406	ALOH+	(2)	4	3.04243E+04	-6.37579E-01	-1.95500E+01	9.55290E-04	-4.98767E-08	
407	ALOH-	(2)	4	1.22012E+05	-5.51846E+00	9.67964E+00	7.33455E-04	-3.74812E-08	
408	ALO+	(2)	4	-4.92552E+04	1.64346E+00	-2.01384E+01	2.24552E-04	-1.28953E-08	
409	ALO-	(2)	4	1.01036E+05	-3.56283E+00	7.49300E+00	2.31304E-04	-1.23579E-08	
410	ALO2-	(2)	4	1.57443E+05	-3.81188E+00	-7.84722E+00	4.97451E-04	-3.00137E-08	
411	AL2O+	(2)	4	3.14094E+04	1.70485E+00	-3.68882E+01	3.09690E-04	-1.90676E-08	
412	AL2O2+	(2)	4	7.15783E+04	1.46107E+00	-5.13076E+01	4.13495E-04	-2.59657E-08	
413	C+	(2)	4	-1.30690E+05	2.50401E+00	-1.46647E+01	7.54634E-06	-2.54479E-09	
414	C-	(2)	4	1.46801E+04	-2.53563E+00	1.36692E+01	-2.17850E-06	6.43116E-09	
415	CF+	(2)	4	-4.21215E+04	1.00408E+00	-1.84405E+01	3.73618E-04	-2.14790E-08	
416	CF2+	(2)	4	-7.81347E+03	-6.99461E-02	-2.53392E+01	7.27936E-04	-4.09138E-08	
417	CF3+	(2)	4	6.42509E+04	-9.09083E-01	-3.84243E+01	1.21915E-03	-6.99523E-08	
418	HCO+	(2)	4	4.21434E+04	-6.85893E-01	-2.13756E+01	9.56708E-04	-5.08929E-08	
419	CO2-	(2)	4	1.97821E+05	-5.35293E+00	3.68660E+00	8.35497E-04	-4.94751E-08	
420	CA+	(2)	4	-7.10242E+04	2.24261E+00	-1.18612E+01	2.11708E-04	-2.32751E-08	
421	CAOH+	(2)	4	3.33347E+04	2.02710E-01	-2.35310E+01	8.47453E-04	-6.51620E-08	
422	CL+	(2)	4	-1.50477E+05	2.63854E+00	-1.48048E+01	-6.53929E-05	5.56901E-09	
423	MGCL+	(2)	4	-4.58871E+04	1.16200E+00	-1.56106E+01	3.72185E-04	-1.28915E-09	
424	SCL+	(2)	4	-7.94408E+04	1.15370E+00	-1.70532E+01	3.12184E-04	-1.38612E-08	
425	CL-	(2)	4	4.18899E+04	-2.81435E+00	1.48801E+01	8.33944E-05	-3.83220E-09	
426	SCL2+	(2)	4	-4.56365E+04	7.23668E-01	-2.80672E+01	3.96735E-04	-2.17906E-08	
427	CR+	(2)	4	-7.83915E+04	2.88638E+00	-1.69127E+01	-2.64365E-04	1.28050E-08	
428	CR-	(2)	4	7.79548E+03	-2.33232E+00	1.29129E+01	-4.98087E-05	-1.72854E-08	
429	F+	(2)	4	-2.02211E+05	2.54557E+00	-1.41415E+01	-2.07386E-05	1.47836E-09	
430	F-	(2)	4	3.94453E+04	-2.68322E+00	1.37944E+01	6.43996E-05	-3.56674E-09	
431	MGF+	(2)	4	-3.34851E+04	2.14451E+00	-2.38707E+01	6.21491E-04	-5.33979E-08	
432	FP+	(2)	4	-6.03624E+04	1.38396E+00	-1.87783E+01	2.80210E-04	-2.29725E-08	
433	FP-	(2)	4	6.63587E+04	-3.42333E+00	8.56651E+00	2.17637E-04	-1.91092E-08	
434	FS+	(2)	4	-7.60242E+04	1.10620E+00	-1.74336E+01	3.45244E-04	-1.70293E-08	
435	FS-	(2)	4	6.43041E+04	-3.62837E+00	8.51478E+00	2.33648E-04	-1.01396E-08	
436	F2K-	(2)	4	1.12418E+05	-3.34383E+00	-6.46042E+00	4.05156E-04	-3.42732E-08	
437	F2MG+	(2)	4	-3.38742E+04	1.43809E+00	-3.25976E+01	4.17191E-04	-2.62826E-08	
438	F2NA-	(2)	4	1.11599E+05	-3.39934E+00	-6.67205E+00	3.71611E-04	-2.53346E-08	
439	F2P+	(2)	4	6.88147E+02	6.49873E-01	-2.93120E+01	5.33114E-04	-3.77410E-08	
440	F2P-	(2)	4	1.32882E+05	-4.34104E+00	-9.25748E-01	5.36337E-04	-3.84804E-08	
441	F2S+	(2)	4	-3.17656E+04	4.69986E-01	-2.79870E+01	5.56605E-04	-3.34074E-08	
442	F2S-	(2)	4	9.92366E+04	-4.08806E+00	-1.84924E+00	4.00129E-04	-2.49314E-08	
443	F3S+	(2)	4	1.48526E+04	-1.95350E-01	-4.24930E+01	9.81089E-04	-5.73199E-08	
444	F3S-	(2)	4	1.54213E+05	-4.58367E+00	-1.65571E+01	7.65935E-04	-4.56783E-08	
445	F4S+	(2)	4	2.18536E+04	2.51161E-01	-5.95136E+01	1.00847E-03	-5.92881E-08	
446	F4S-	(2)	4	1.77287E+05	-4.03754E+00	-3.53696E+01	7.36157E-04	-4.24009E-08	
447	F5S+	(2)	4	6.06289E+04	-3.68076E-01	-7.66343E+01	1.40799E-03	-7.73041E-08	
448	F5S-	(2)	4	2.32662E+05	-4.52601E+00	-5.13589E+01	1.11553E-03	-6.56215E-08	
449	F6S-	(2)	4	2.50763E+05	-6.13458E+00	-6.21687E+01	1.88761E-03	-1.09317E-07	
450	FE+	(2)	4	-9.12289E+04	2.75926E+00	-1.59394E+01	5.91507E-05	-1.43108E-08	
451	FE-	(2)	4	1.99443E+05	-2.30441E+00	1.28276E+01	-3.38488E-05	-1.03653E-08	
452	H+	(2)	4	-1.57812E+05	2.49925E+00	-1.49094E+01	4.26743E-07	-2.77317E-11	
453	H-	(2)	4	8.75238E+03	-2.50077E+00	1.35336E+01	3.39359E-07	-2.30826E-11	
454	KOH+	(2)	4	7.26817E+03	6.85217E-01	-2.47635E+01	5.77413E-04	-3.99424E-08	
455	MGOH+	(2)	4	3.80425E+03	3.72841E-01	-2.49807E+01	6.28860E-04	-3.48352E-08	
456	NAOH+	(2)	4	-1.21469E+04	7.63522E-01	-2.53869E+01	4.94294E-04	-2.81809E-08	
457	H3O+	(2)	4	3.77691E+04	-3.91099E+00	-1.00755E+01	1.65917E-03	-7.69891E-08	
458	HE+	(2)	4	-2.85341E+05	2.50051E+00	-1.35313E+01	-1.69595E-07	1.04487E-11	
459	K+	(2)	4	-5.04289E+04	2.31508E+00	-1.37286E+01	1.65367E-04	-2.07326E-08	
460	K-	(2)	4	5.76506E+03	-2.68458E+00	1.47105E+01	1.65063E-04	-2.07118E-08	
461	KO-	(2)	4	5.63095E+04	-3.35727E+00	8.34499E+00	2.92793E-04	-2.56426E-08	
462	MG+	(2)	4	-8.87612E+04	2.41951E+00	-1.30050E+01	6.48336E-05	-6.96143E-09	
463	MN+	(2)	4	-8.63475E+04	2.37306E+00	-1.32278E+01	8.55175E-05	-5.76691E-09	
464	N+	(2)	4	-1.68786E+05	2.49827E+00	-1.34176E+01	1.06740E-05	-1.48024E-09	
465	N-	(2)	4	-8.59054E+02	-2.52800E+00	1.52033E+01	2.79360E-05	-3.33707E-09	
466	NO2-	(2)	4	1.39863E+05	-5.01361E+00	1.53965E+00	7.35500E-04	-4.31197E-08	
467	N2-	(2)	4	9.46450E+04	-4.15804E+00	1.27568E+01	4.20205E-04	-2.52129E-08	
468	N2O+	(2)	4	-1.64137E+04	4.33632E-01	-2.92328E+01	7.58449E-04	-4.59623E-08	
469	NA+	(2)	4	-5.96691E+04	2.40644E+00	-1.43021E+01	7.73455E-05	-8.68951E-09	
470	NA-	(2)	4	6.32960E+03	-2.59364E+00	1.41399E+01	7.72592E-05	-8.68339E-09	
471	NAO-	(2)	4	5.65147E+04	-3.31499E+00	7.81169E+00	2.19776E-04	-1.46596E-08	
472	NI+	(2)	4	-8.97596E+04	3.84622E+00	-1.82408E+01	-7.84448E-04	5.18262E-08	
473	NI-	(2)	4	1.34028E+04	-2.82102E+00	1.54025E+01	8.09394E-05	-5.05427E-09	
474	O+	(2)	4	-1.57987E+05	2.43207E+00	-1.45369E+01	2.89552E-05	-2.13814E-09	
475	O-	(2)	4	1.69699E+04	-2.52288E+00	1.39705E+01	1.54665E-05	-1.66096E-09	
476	O2-	(2)	4	6.47418E+04	-3.76154E+00	8.79338E+00	3.18913E-04	-1.78266E-08	
477	P+	(2)	4	-1.21903E+05	2.72917E+00	-1.51664E+01	-4.32460E-05	-3.13707E-09	
478	P-	(2)	4	8.57593E+03	-2.53937E+00	1.52023E+01	7.79867E-05	-1.00298E-08	
479	S+	(2)	4	-1.20204E+05	2.30891E+00	-1.35347E+01	3.49301E-05	9.99419E-11	
480	S-	(2)	4	2.41277E+04	-2.53558E+00	1.40721E+01	2.03592E-05	-3.58973E-09	
481	SI+	(2)	4	-9.45279E+04	2.70973E+00	-1.60951E+01	-1.04991E-04	3.59990E-09	
482	SI-	(2)	4	1.61666E+04	-2.72353E+00	1.49716E+01	8.90895E-05	5.32288E-09	
483	TI+	(2)	4	-7.90118E+04	3.09656E+00	-1.77870E+01	-1.55372E-04	-3.93128E-09	
484	TI-	(2)	4	9.61744E+02	-2.43366E+00	1.41116E+01	-2.07353E-05	-1.81706E-08	
485	V+	(2)	4	-7.83613E+04	1.93698E+00	-1.04833E+01	3.08091E-04	-2.17895E-08	
486	V-	(2)	4	5.91880E+03	-3.03080E+00	1.79389E+01	9.81012E-05	-1.41978E-08	

Table D.1. continued

	molecule	source	fit	a_0	a_1	a_2	a_3	a_4	σ [kJ/mol]
487	LIF4AL	(2)	4	3.19697E+05	-1.96460E+00	-6.18249E+01	1.14317E-03	-7.07626E-08	±0.50
488	CLFLI2	(2)	4	1.52729E+05	-1.98975E+00	-2.75831E+01	7.99301E-04	-5.56238E-08	
489	LIOCL	(2)	4	6.49261E+04	-2.13332E+00	-1.14020E+01	6.20542E-04	-4.05228E-08	
490	WCL	(3)	4	5.00275E+04	-1.10972E+00	-1.72470E+00	-2.74247E-04	1.37409E-08	
491	ZRCL	(2)	4	6.28220E+04	-1.73142E+00	1.23608E-01	4.06003E-04	-3.21988E-08	
492	(LICL)2	(2)	4	1.39160E+05	-1.40010E+00	-2.94460E+01	5.55678E-04	-4.10760E-08	±0.46 ±0.72
493	WO2CL2	(3)	4	2.71271E+05	-3.29372E+00	-3.51848E+01	1.14648E-03	-8.88251E-08	
494	WCL2	(3)	4	1.32700E+05	-1.21609E+00	-1.53398E+01	1.99702E-05	-7.57683E-10	
495	ZRCL2	(2)	4	1.24542E+05	-1.57847E+00	-1.60777E+01	3.00256E-04	-1.79057E-08	
496	(LICL)3	(2)	4	2.21711E+05	-2.11829E+00	-5.62529E+01	1.21088E-03	-8.31090E-08	
497	ZRCL3	(2)	4	1.79835E+05	-1.70228E+00	-2.97911E+01	5.09972E-04	-4.08965E-08	±0.10
498	WCL4O	(2)	4	2.58497E+05	-3.83453E+00	-4.89577E+01	9.60739E-04	-4.50604E-08	
499	WCL4	(2)	4	2.00071E+05	-2.93945E+00	-3.46697E+01	4.29448E-04	-1.04938E-08	
500	ZRCL4	(2)	4	2.35955E+05	-1.78407E+00	-4.55961E+01	4.61974E-04	-3.49582E-08	
501	WCL5	(2)	4	2.23864E+05	-3.04356E+00	-5.08091E+01	6.51001E-04	-2.02868E-08	
502	WCL6	(2)	4	2.48302E+05	-2.80633E+00	-7.07046E+01	7.01542E-04	-2.79814E-08	±0.33
503	WCL210	(2)	4	4.53680E+05	-3.55157E+00	-1.30172E+02	1.07060E-03	-3.86873E-08	
504	LIOF	(2)	4	6.93234E+04	-1.92822E+00	-1.32924E+01	5.79346E-04	-3.91546E-08	
505	WF	(3)	4	6.50683E+04	-1.23658E+00	-1.50575E+00	1.23019E-04	-2.94188E-08	
506	ZRF	(2)	4	7.25614E+04	-1.72651E+00	-5.33044E-01	4.69202E-04	-4.6070E-08	
507	LIF2-	(2)	4	1.22731E+05	-4.11219E+00	-2.89615E+00	6.26469E-04	-4.06383E-08	±0.22
508	LI2F2	(2)	4	1.70485E+05	-1.74505E+00	-2.96042E+01	7.43988E-04	-5.34789E-08	
509	ZRF2	(2)	4	1.59103E+05	-2.13985E+00	-1.21185E+01	3.82969E-04	-2.42775E-08	
510	F3LI3	(2)	4	2.68403E+05	-1.68551E+00	-5.90717E+01	1.13947E-03	-8.15027E-08	
511	ZRF3	(2)	4	2.34682E+05	-1.85702E+00	-3.08813E+01	6.57192E-04	-5.17709E-08	
512	WOF4	(2)	4	3.30236E+05	-3.94943E+00	-5.07060E+01	1.12586E-03	-5.78699E-08	±0.58 ±0.94 ±0.81 ±1.10 ±2.40
513	ZRF4	(2)	4	3.12537E+05	-2.00525E+00	-4.73596E+01	6.63103E-04	-4.97117E-08	
514	WF6	(2)	4	3.65791E+05	-3.70552E+00	-7.05225E+01	1.21404E-03	-6.21365E-08	
515	LIOH	(2)	4	1.03034E+05	-1.94286E+00	-1.19422E+01	5.72348E-04	-3.38212E-08	
516	LIOH+	(2)	4	-1.79956E+04	6.13541E-01	-2.51902E+01	5.56216E-04	-3.29740E-08	
517	ZRH	(2)	4	3.70538E+04	-1.82214E+00	2.20061E+00	2.49990E-04	-2.36279E-08	±0.22
518	(LIOH)2	(2)	4	2.34948E+05	-5.69699E+00	-3.10024E+01	2.12755E-03	-1.21873E-07	
519	H2WO4	(3)	4	3.81291E+05	-8.84105E+00	-2.70678E+01	5.34328E-03	-5.38002E-07	
520	Li+	(2)	4	-6.26000E+04	2.40549E+00	-1.43075E+01	8.37507E-05	-1.03116E-08	
521	Li-	(2)	4	7.14342E+03	-2.59560E+00	1.41416E+01	8.41796E-05	-1.03403E-08	
522	LIN	(2)	4	3.55149E+04	-9.47062E-01	-3.96290E+00	2.82276E-04	-2.05672E-08	±0.58 ±0.94 ±0.81 ±1.10 ±2.40
523	LINO	(2)	4	8.39059E+04	-1.86393E+00	-1.30494E+01	5.62377E-04	-3.96022E-08	
524	LINAO	(2)	4	7.41831E+04	-1.89448E+00	-1.16082E+01	6.09671E-04	-4.51642E-08	
525	LIO-	(2)	4	5.61860E+04	-3.52398E+00	8.72195E+00	2.83444E-04	-2.05889E-08	
526	LI2O	(2)	4	8.81260E+04	-1.19280E+00	-1.77261E+01	5.42980E-04	-4.26854E-08	
527	LI2O2	(2)	4	1.27278E+05	-9.05931E-01	-3.36080E+01	4.52673E-04	-3.86062E-08	±0.22
528	LI2SO4	(2)	4	3.16340E+05	-3.34050E+00	-7.20382E+01	1.90021E-03	-1.24077E-07	
529	WO2	(3)	4	1.52223E+05	-2.90088E+00	-6.10642E+00	6.14536E-04	-5.66091E-08	
530	ZRO2	(2)	4	1.67287E+05	-2.03009E+00	-1.46115E+01	3.22191E-04	-3.20231E-08	
531	WO3	(3)	4	2.26374E+05	-3.71610E+00	-2.00906E+01	1.23669E-03	-9.83304E-08	
532	(WO3)2	(3)	4	5.23332E+05	-3.64727E+00	-8.39396E+01	1.81177E-03	-1.58830E-07	±0.58 ±0.94 ±0.81 ±1.10 ±2.40
533	W3O8	(3)	4	7.50529E+05	-6.78493E+00	-1.14407E+02	4.29334E-03	-4.24109E-07	
534	(WO3)3	(3)	4	8.17838E+05	-8.05561E+00	-1.24426E+02	5.41494E-03	-5.27128E-07	
535	(WO3)4	(3)	4	1.10399E+06	-7.27410E+00	-1.95635E+02	4.47960E-03	-3.71841E-07	
536	W+	(2)	4	-9.32002E+04	1.13666E+00	-4.13254E+00	4.22006E-04	-1.64143E-08	
537	W-	(2)	4	8.70581E+03	-4.19457E+00	2.81349E+01	-7.53206E-05	1.09960E-08	±1.70 ±0.50
538	ZR+	(2)	4	-7.94818E+04	2.74809E+00	-1.56783E+01	-2.26874E-05	-1.10085E-08	
539	ZR-	(2)	4	4.72735E+03	-2.23076E+00	1.31427E+01	-3.45945E-04	7.18428E-09	
540	TIC	(4)	5	12.75293	-5.44850	-1.56672	1.56041	-0.93275	
541	Si(CH3)4	(3)	4	7.44905E+05	-1.44347E+01	-1.25573E+02	0.00000E+00	0.00000E+00	
542	SiCH3CL3	(3)	4	3.23859E+05	-9.66441E+00	-5.80646E+01	0.00000E+00	0.00000E+00	±1.70 ±0.50
543	SiH2	(5)	4	7.50441E+04	-1.11618E+00	-1.93312E+01	9.97592E-05	-7.15829E-10	
544	SiH3	(5)	4	1.16152E+05	-2.16682E+00	-2.41859E+01	3.65595E-04	-1.18831E-08	
545	TIC2	(5)	4	1.42116E+05	2.43265E+00	-4.72869E+01	-1.63153E-03	8.80077E-08	
546	C3H	(5)	4	2.21278E+05	3.37781E+00	-6.68686E+01	-2.75673E-03	1.50780E-07	
547	MNS	(5)	4	3.65977E+04	2.84321E+00	-3.24266E+01	-1.72506E-03	9.16702E-08	±1.70 ±0.50
548	CRC2	(5)	4	1.27594E+05	3.64957E+00	-5.48448E+01	-2.14973E-03	1.13743E-07	
549	Si2C2	(5)	4	2.01062E+05	1.42585E+00	-5.73237E+01	-6.10383E-04	3.03589E-08	
550	VC	(5)	4	1.41928E+05	1.38059E+00	-4.09084E+01	-1.16650E-03	6.46411E-08	
551	TIC4	(5)	4	2.92656E+05	2.79141E+00	-8.51696E+01	-1.43538E-03	7.41507E-08	
552	VC4	(5)	4	2.90307E+05	2.24789E+00	-8.25056E+01	-1.15151E-03	5.93340E-08	

Table D.2. Condensed phase data fitted and collected from different sources.

no	formula	name	ρ [g/cm ³]	fit	restriction	c_0	c_1	c_2	c_3	c_4	σ [kJ/mol]
1	Al ₂ O ₃ [s]	CORUNDUM(α)	3.97	5		3.68836E+05	-7.75003E+00	-4.32225E+01	7.53364E-03	-7.64141E-07	±0.54
2	Al ₂ O ₃ [l]		2.78	5		3.64246E+05	-1.63572E+00	-8.13311E+01	2.51029E-03	3.18884E-08	±1.80
3	MgAl ₂ O ₄ [s]	SPINEL	3.58	5		4.90917E+05	-1.01783E+01	-6.09577E+01	1.00839E-02	-9.21697E-07	±0.57
4	MgAl ₂ O ₄ [l]		2.75	5		4.70478E+05	-1.35143E+00	-1.08539E+02	2.76930E-03	-7.78980E-08	±0.86
5	TiO ₂ [s]	RUTILE	4.23	3	<4047K	-7.70443E+04	4.03144E+01	-2.59140E-03	6.02422E-07	-6.86899E-11	±0.04
6	TiO ₂ [l]		3.40	3		-7.07819E+04	3.68021E+01	-1.29214E-03	-6.87776E-08	1.89408E-11	±0.02
7	Ti ₄ O ₇ [s]	TITANIUM-OXIDE	4.19	2		1.85821E+06	-7.06188E+06	1.70429E+03	-4.67044E-02	4.71514E-06	±0.09
8	Ti ₄ O ₇ [l]		3.40	2		5.06384E+06	-6.89199E+06	1.64108E+03	-5.93652E-02	4.64893E-06	±0.43
9	Mg ₂ SiO ₄ [s]	FOSTERITE	3.21	2	<5298K	1.70117E+06	-3.93154E+06	1.04604E+03	-1.77300E-02	7.18369E-07	±0.14
10	Mg ₂ SiO ₄ [l]		2.60	2		3.26302E+06	-3.87698E+06	1.03152E+03	-2.48587E-02	1.67747E-06	±0.15
11	MgSiO ₃ [s]	ENSTATITE	3.19	2		1.06060E+06	-2.90401E+06	7.57655E+02	-1.37019E-02	1.51773E-06	±1.30
12	MgSiO ₃ [l]		2.60	2		2.94675E+06	-2.86104E+06	7.52974E+02	-2.55722E-02	2.36166E-06	±0.15
13	Fe[s]	IRON(α - δ)	7.87	3		-4.99622E+04	3.21370E+01	6.68363E-04	-1.13811E-06	2.57277E-10	±0.05
14	Fe[l]		6.98	3		-4.87741E+04	3.26355E+01	-1.38271E-03	1.09701E-07	0.00000E+00	±0.03
15	Fe ₂ SiO ₄ [s]	FAYALITE	4.39	5		4.49574E+05	-9.27141E+00	-6.61500E+01	1.04125E-02	-1.09203E-06	±0.38
16	FeS[s]	TROILITE	4.83	3		-5.69922E+04	3.86753E+01	-4.68301E-03	1.03559E-06	-8.42872E-11	±0.08
17	FeS[l]		3.90	3		-5.26135E+04	3.46138E+01	-3.55056E-03	7.59195E-07	-6.94708E-11	±0.08
18	MgTi ₂ O ₅ [s]	MG-DITITANATE	4.64	2	<5318K	2.12742E+06	-4.86574E+06	1.21637E+03	-2.34246E-02	8.39658E-07	±0.40
19	MgTi ₂ O ₅ [l]		3.70	2		5.49485E+06	-4.75803E+06	1.18602E+03	-4.08290E-02	3.08428E-06	±0.32
20	C[s]	GRAPHITE	2.27	4		3.27860E+01	-8.65139E+04	4.80395E-01	0.00000E+00	0.00000E+00	±0.06
21	TiC[s]	TITANIUM-CARBIDE	4.93	2	<6425K	1.11878E+05	-1.37612E+06	3.20666E+02	-4.63379E-03	1.85306E-07	±0.32
22	TiC[l]		3.90	2		5.34201E+05	-1.30289E+06	3.01268E+02	-6.61621E-03	5.21065E-07	±0.03
23	SiC[s]	SILICON-CARBIDE(α)	3.21	2		6.73337E+05	-1.24381E+06	3.21779E+02	-4.54405E-03	2.69711E-07	±0.07
24	SiO[s]	SI-MONOXIDE	2.18	10		-4.95200E+04	3.25200E+01				
25	SiO ₂ [s]	QUARTZ	2.65	3		-7.28086E+04	3.65312E+01	-2.56109E-04	-5.24980E-07	1.53343E-10	±0.08
26	SiO ₂ [l]		2.10	3		-7.18591E+04	3.59570E+01	-6.57765E-04	-3.78705E-08	8.78339E-12	±0.08
27	Zr[s]	ZIRCONIUM(β)	6.52	3	<3565K	-7.28816E+04	3.11094E+01	-6.94825E-04	2.42125E-07	-3.55274E-11	±0.01
28	Zr[l]		5.80	3		-7.14144E+04	3.06541E+01	-5.47166E-04	4.59756E-08	-9.40737E-13	±0.02
29	ZrO ₂ [s]	BADDELEYITE	5.68	3		-9.77238E+04	4.18463E+01	-2.22011E-03	3.20813E-07	-1.60657E-11	±0.06
30	ZrO ₂ [l]		5.00	3		-8.87286E+04	3.86685E+01	-1.94950E-03	2.38918E-07	-1.43097E-11	±0.11
31	ZrSiO ₄ [s]	ZR-SILICATE	4.56	5		4.90061E+05	-8.78711E+00	-5.52714E+01	8.61307E-03	-1.02725E-06	±0.26
32	W[s]	TUNGSTEN	19.25	3	<4431K	-1.02351E+05	3.07543E+01	-6.13643E-06	5.36528E-08	-1.19522E-11	±0.03
33	W[l]		17.60	3		-9.66863E+04	2.90491E+01	2.11260E-04	-5.93323E-08	6.15431E-12	±0.01
34	WO ₃ [s]	W-TRIOXIDE	7.16	3	<3476K	-6.63560E+04	4.06161E+01	-2.83452E-03	6.33945E-07	-7.73057E-11	±0.01
35	WO ₃ [l]		7.00	3		-5.98226E+04	3.68387E+01	-1.60492E-03	-3.33151E-07	8.09588E-11	±0.01
36	MgO[s]	PERICLASE	3.58	3	<5782K	-7.91838E+04	3.57312E+01	1.45021E-04	-8.47194E-08	4.49221E-12	±0.09
37	MgO[l]		2.90	3		-7.07437E+04	3.25683E+01	6.94583E-04	-2.71415E-07	2.24580E-11	±0.04
38	FeO[s]	FERROPERICLASE	5.99	3		-6.30018E+04	3.66364E+01	-2.42990E-03	3.18636E-07	0.00000E+00	±0.01
39	FeO[l]		5.50	3		-6.03086E+04	3.48392E+01	-2.21932E-03	2.77859E-07	-1.56828E-11	±0.01
40	Na ₂ SiO ₃ [s]	NA-METASILICATE	2.40	2	<3108K	1.31453E+06	-2.98538E+06	8.76910E+02	-2.65571E-02	1.15443E-06	±0.17
41	Na ₂ SiO ₃ [l]		1.90	2		1.72092E+06	-2.93824E+06	8.49423E+02	-3.54866E-02	3.74762E-06	±0.09
42	H ₂ O[s]	WATER-ICE	0.93	7	< 747K	611.5	23.036	-333.7	279.82		
43	H ₂ O[l]	WATER	1.00	6	> 193K	2.98605E+01	-3.15220E+03	-7.30370E+00	2.42470E-09	1.80900E-06	
44	NH ₃ [s/l]	AMONIA	0.87	8		10.53	-2161.0	-86596.0			
45	CH ₄ [s/l]	METHANE	0.66	9		3.9895	-443.028	-0.490			
46	CO[s/l]	C-MONOXIDE	1.14	6		5.18145E+01	-7.88240E+02	-2.27340E+01	5.12250E-02	4.66030E-11	
47	CO ₂ [s/l]	C-DIOXIDE	1.98	6		3.50187E+01	-1.51190E+03	-1.13350E+01	9.33830E-03	7.76260E-10	
48	H ₂ SO ₄ [s/l]	SULPHURIC-ACID	1.84	2		9.70368E+05	-2.53825E+06	9.35422E+02	-4.96224E-02	0.00000E+00	±0.35
49	Na[s]	SODIUM	0.97	3		-1.30424E+04	2.70281E+01	-1.44654E-03	-4.12467E-07	3.86734E-10	±0.00
50	Na[l]		0.93	3		-1.26494E+04	2.58560E+01	-1.36371E-03	2.63744E-07	0.00000E+00	±0.00
51	NaCl[s]	HALITE	2.17	3		-2.79146E+04	3.46023E+01	-3.11287E-03	5.30965E-07	-2.59584E-12	±0.00
52	NaCl[l]		1.30	3		-2.48880E+04	3.18494E+01	-3.08748E-03	4.84990E-07	-2.60359E-11	±0.13
53	KCl[s]	SYLVITE	1.99	3		-2.69250E+04	3.39574E+01	-2.04903E-03	-2.83957E-07	1.82974E-10	±0.01
54	KCl[l]		1.60	3		-2.50293E+04	3.39453E+01	-4.61815E-03	7.36857E-07	0.00000E+00	±0.04
55	S[s]	SULPHUR	2.05	3		-3.32020E+04	2.90980E+01	3.44461E-03	-4.54179E-06	1.85126E-09	±0.00
56	S[l]		1.70	3		-3.32601E+04	3.07134E+01	-2.01083E-03	4.47359E-07	0.00000E+00	±0.04
57	MgS[s]	MG-SULPHIDE	2.68	2		-3.08737E+04	-7.70053E+05	2.69294E+02	-5.60302E-03	3.38303E-07	±0.11
58	LiCl[s]	LI-CHLORIDE	2.07	3		-2.57733E+04	3.36349E+01	-2.43056E-03	2.35233E-07	0.00000E+00	±0.08
59	LiCl[l]		1.70	3		-2.38196E+04	3.23179E+01	-4.07306E-03	1.00591E-06	-9.65764E-11	±0.01
60	SiS ₂ [s]	SI-DISULFIDE	1.85	2	<2571K	-1.96413E+05	-1.21744E+06	4.29645E+02	-1.32885E-02	9.27515E-07	±0.01
61	SiS ₂ [l]		1.50	2		5.45975E+05	-1.21556E+06	4.34344E+02	-1.99379E-02	2.31933E-06	±0.02

Solid compounds are marked with '[s]' and liquids with '[l]', whereas '[s/l]' denote condensates treated as a single combined species. The standard pressure is $p^\circ = 1$ bar unless otherwise noted. $T_C = T - 273.15$ is the temperature in °C. Restriction: this fit should not be applied beyond the indicated temperature as it will create a secondary spurious intersection point with the other phase. In GGchem, we deal with this by artificially increasing p^{vap} above/below this temperature for solid/liquid species, respectively, which is far away from the melting point anyway. σ is the standard deviation of the fitted values from the data points [kJ/mol].

$$\begin{aligned}
\text{fit = 1:} \quad \Delta G_f^\circ [\text{cal/mol}] &= c_0/T + c_1 + c_2 T + c_3 T^2 + c_4 T^3, \\
\text{fit = 2:} \quad \Delta G_f^\circ [\text{J/mol}] &= c_0/T + c_1 + c_2 T + c_3 T^2 + c_4 T^3, \\
\text{fit = 3:} \quad \ln p^{\text{vap}} [\text{dyn/cm}^2] &= c_0/T + c_1 + c_2 T + c_3 T^2 + c_4 T^3 \\
\text{fit = 4:} \quad \ln p^{\text{vap}} [\text{dyn/cm}^2] &= c_0 + c_1/(T + c_2) \\
\text{fit = 5:} \quad -\Delta G_f^\circ/RT &= c_0/T + c_1 \ln(T) + c_2 + c_3 T + c_4 T^2 \\
\text{fit = 6:} \quad \log_{10} p^{\text{vap}} [\text{mmHg}] &= c_0 + c_1/T + c_2 \log_{10} T + c_3 T + c_4 T^2 \\
\text{fit = 7:} \quad p^{\text{vap}} [\text{dyn/cm}^2] &= c_0 \exp[(c_1 T_C + T_C^2/c_2)/(T_C + c_3)] \\
\text{fit = 8:} \quad \ln p^{\text{vap}} [\text{bar}] &= c_0 + c_1/T + c_2/T^2 \\
\text{fit = 9:} \quad \log_{10} p^{\text{vap}} [\text{bar}] &= c_0 + c_1/(T + c_2) \\
\text{fit = 10:} \quad \ln p^{\text{vap}} [\text{dyn/cm}^2] &= c_0/T + c_1
\end{aligned}$$

data from [Sharp & Huebner \(1990\)](#), note that $p^\circ = 1$ atm fitted to NIST-JANAF data ([Chase 1986](#)) by the authors fitted to NIST-JANAF data ([Chase 1986](#)) by the authors fitted to NIST-JANAF data ([Chase 1986](#)) by the authors fitted to NIST-JANAF data ([Chase 1986](#)) by the authors data from [Yaws \(1999\)](#) data from [Ackerman & Marley \(2001\)](#) data from [Weast \(1971\)](#) data from [Prydz & Goodwin \(1972\)](#) data from [Gail et al. \(2013\)](#)

Table D.2. continued

no	formula	name	ρ [g/cm ³]	fit	restriction	c_0	c_1	c_2	c_3	c_4	σ [kJ/mol]
62	AlCl ₃ [s]	AL-TRICHLORIDE	2.48	2	<1998K	4.17188E+05	-1.40425E+06	5.68472E+02	-1.79150E-02	-4.81449E-06	±0.08
63	AlCl ₃ [l]		2.00	2		-1.51097E+06	-1.36116E+06	4.88316E+02	-2.96109E-02	3.95859E-06	±0.01
64	CaO[s]	LIME	3.35	2		-4.94275E+04	-1.06229E+06	2.81700E+02	-7.00317E-03	4.60182E-07	±0.20
65	CaO[l]		2.70	2		7.88379E+05	-9.87984E+05	2.61388E+02	-8.44845E-03	6.19197E-07	±0.15
66	CaCl ₂ [s]	CA-DICHLORIDE	2.15	3		-3.91597E+04	3.71795E+01	-1.93815E-03	1.18087E-07	0.00000E+00	±0.09
67	CaCl ₂ [l]		1.70	3		-3.68100E+04	3.61822E+01	-3.53647E-03	5.03350E-07	-3.05198E-11	±0.03
68	LiH[s]	LI-HYDRIDE	0.78	3		-2.76405E+04	3.07600E+01	3.06028E-03	-3.26876E-06	7.70757E-10	±0.04
69	LiH[l]		0.62	3		-2.63155E+04	3.30329E+01	-3.77190E-03	6.41033E-07	0.00000E+00	±0.04
70	MgTiO ₃ [s]	GEIKIELITE	3.88	2		8.18358E+05	-2.94809E+06	7.57824E+02	-1.56613E-02	1.01510E-06	±0.20
71	MgTiO ₃ [l]		3.10	2		2.95447E+06	-2.88536E+06	7.40614E+02	-2.52836E-02	1.86167E-06	±0.19
72	K ₂ SiO ₃ [s]	K-SILICATE	2.40	2		1.08754E+06	-2.93466E+06	8.61167E+02	-3.66511E-02	3.06399E-06	±0.09
73	K ₂ SiO ₃ [l]		1.90	2		-2.09568E+06	-2.87130E+06	8.04917E+02	-3.03325E-02	2.75492E-06	±0.34
74	Ti[s]	TITANIUM(beta)	4.51	2	<4302K	1.31154E+05	-4.67808E+05	1.45369E+02	-3.49413E-03	0.00000E+00	±0.02
75	Ti[l]		4.14	2		1.72969E+06	-4.68384E+05	1.54389E+02	-9.64953E-03	7.21551E-07	±0.18
76	TiO[s]	TI-MONOXIDE(β)	4.95	3		-7.14013E+04	3.80934E+01	-1.49104E-03	6.06190E-08	0.00000E+00	±0.01
77	TiO[l]		4.00	3		-6.61504E+04	3.58743E+01	-1.97510E-03	2.33017E-07	-1.21467E-11	±0.01
78	LiOH[s]	LI-HYDROXIDE	1.46	3		-3.01062E+04	3.37718E+01	9.66853E-04	-2.02320E-06	5.34064E-10	±0.04
79	LiOH[l]		1.20	3		-2.87677E+04	3.42848E+01	-3.81580E-03	5.39401E-07	0.00000E+00	±0.07
80	VO[s]	V-MONOXIDE	5.76	3	<3404K	-6.74603E+04	3.82717E+01	-2.78551E-03	5.72078E-07	-7.41840E-11	±0.02
81	VO[l]		5.20	3		-6.01063E+04	3.45413E+01	-2.64854E-03	4.70724E-07	-3.85068E-11	±0.01
82	V ₂ O ₃ [s]	KARELIANITE	4.95	5		3.59474E+05	-4.09857E+00	-6.48556E+01	4.73778E-03	-3.36133E-07	±0.32
83	V ₂ O ₄ [s]	PARAMONTROSEITE	4.09	5		4.14455E+05	-2.22575E+00	-9.36637E+01	3.73856E-03	-2.64367E-07	±0.53
84	V ₂ O ₅ [s]	SHCHERBINAIT	3.37	5		4.58651E+05	-6.83064E+00	-8.22840E+01	6.02078E-03	0.00000E+00	±0.51
85	CaS[s]	CALCIUM-SULFIDE	2.59	2		-8.11371E+04	-9.28021E+05	2.69568E+02	-7.11550E-03	5.21453E-07	±0.13
86	FeS ₂ [s]	PYRITE	4.90	2		5.30142E+05	-1.14524E+06	4.72929E+02	-4.33681E-03	-1.49521E-06	±0.05
87	Na ₂ S[s]	NA-SULFIDE	1.86	2		1.99053E+06	-8.70027E+05	4.06721E+02	-2.78298E-02	0.00000E+00	±0.30
88	Mn[s]	MANGANESE($\alpha - \delta$)	7.43	3		-3.41316E+04	3.12732E+01	-5.59461E-04	-4.63498E-07	1.01677E-10	±0.01
89	Mn[l]		5.95	3		-3.22207E+04	3.01880E+01	-1.14796E-03	-3.02939E-08	2.15830E-11	±0.00
90	MnS[s]	ALABANDITE	4.08	1		1.12482E+05	-1.81938E+05	5.87107E+01	8.89360E-05	-4.20876E-09	
91	Ni[s]	NICKEL	8.91	3	<2915K	-5.17726E+04	3.25355E+01	-8.54276E-04	1.93715E-07	-3.72769E-11	±0.03
92	Ni[l]		7.81	3		-4.96579E+04	3.09204E+01	-2.80251E-04	-1.00993E-07	1.63913E-11	±0.00
93	Cr[s]	CHROMIUM	7.19	3	<2981K	-4.78455E+04	3.22423E+01	-5.28710E-04	-6.17347E-08	2.88469E-12	±0.00
94	Cr[l]		6.30	3		-4.47712E+04	3.09753E+01	-7.84094E-04	-5.92580E-10	1.25866E-11	±0.01
95	CrN[s]	CARLSBERGITE	6.09	3		-7.51246E+04	3.88434E+01	-3.48286E-03	1.12997E-06	-1.66901E-10	±0.06
96	CaSiO ₃ [s]	WOLLASTONITE	2.91	5		3.60350E+05	-6.69046E+00	-4.57724E+01	7.14526E-03	-8.10884E-07	±0.26
97	CaTiO ₃ [s]	PEROVSKITE	3.98	1		1.19107E+04	-7.30327E+05	1.75930E+02	-2.84630E-03	1.10392E-07	
98	NiS[s]	MILLERITE	5.37	1		-4.32421E+04	-1.89653E+05	7.16749E+01	-2.50684E-03	0.00000E+00	
99	NiS ₂ [s]	VAESITE	4.45	1		0.00000E+00	-2.66982E+05	1.08176E+02	-1.96812E-03	0.00000E+00	
100	Ni ₃ S ₂ [s]	HEAZLEWOODITE	5.87	1		3.12877E+05	-5.01379E+05	2.14522E+02	-3.99302E-02	7.88559E-06	

Table D.3. Overview of pairs of solid/liquid data and melting points.

condensate	intersection point [K]	melting point [K]	ref.
W	3702	3680	(1)
TiC	3292	3290	(1)
CaO	3200	3200	(1)
MgO	3105	3105	(1)
ZrO ₂	2952	2950	(1)
MgAl ₂ O ₄	2408	2408	(1)
Al ₂ O ₃	2327	2327	(1)
Mg ₂ SiO ₄	2171	2171	(1)
TiO ₂	2129	2130	(1)
Cr	2137	2130	(1)
VO	2063	2063	(1)
TiO	2025	2023	(1)
MgTi ₂ O ₅	1960	1963	(1)
MgTiO ₃	1949	1953	(1)
Ti ₄ O ₇	1947	1950	(1)
Ti	1919	1939	(1)
MgSiO ₃	1852	1850	(1)
Fe	1822	1809	(1)
Ni	1725	1728	(1)
FeS	1490	1463	(1)
Mn	1516	1519	(1)
SiO ₂	1698	1696	(1)
Zr	2116	2125	(1)
WO ₃	1748	1745	(1)
FeO	1651	1650	(1)
SiS ₂	1364	1363	(1)
Na ₂ SiO ₃	1362	1362	(1)
K ₂ SiO ₃	1245	1249	(1)
NaCl	1078	1074	(1)
KCl	1054	1044	(1)
CaCl ₂	1035	1045	(1)
LiH	966	962	(1)
LiCl	877	883	(1)
LiOH	733	744	(1)
AlCl ₃	466	466	(1)
S	385	388	(1)
Na	368	371	(1)
H ₂ O	273	273	(2)

(1) data from NIST-JANAF ([Chase 1986](#)), fitted by the authors(2) fits from ([Ackerman & Marley 2001](#)) and ([Yaws 1999](#))

Table D.4. Condensed phase Gibbs free energy data extracted from the SUPCRTBL database (Zimmer et al. 2016).

no	formula	name	$\Delta G_{f,\text{SU}}^*(T_{\text{ref}})/N$	b_0	b_1	b_2	b_3	b_4	σ [kJ/mol]
1	ZrO ₂	BADDELEYITE	-347.6	2.648723e+05	-3.340550e+00	-3.414969e+01	3.635220e-03	-4.584324e-07	±0.14
2	ZrSiO ₄	ZIRCON	-320.1	4.900611e+05	-8.787114e+00	-5.527139e+01	8.613074e-03	-1.027249e-06	±0.26
3	Al ₂ O ₃	CORUNDUM	-316.4	3.686315e+05	-8.641215e+00	-3.798714e+01	8.874690e-03	-1.024209e-06	±0.42
4	Ca ₂ Al ₂ SiO ₇	GEHLENITE	-315.9	8.625975e+05	-1.533457e+01	-1.143058e+02	1.632128e-02	-1.836678e-06	±0.53
5	Ca ₃ Al ₂ Si ₃ O ₁₂	GROSSULAR	-314.1	1.457200e+06	-3.049843e+01	-1.691883e+02	3.164365e-02	-3.657780e-06	±1.28
6	Ca ₂ SiO ₄	LARNITE	-313.3	4.925717e+05	-7.561371e+00	-7.350215e+01	8.818589e-03	-9.618190e-07	±0.23
7	CaAl ₂ SiO ₆	Ca-TSCHERMAKS	-313.3	7.292797e+05	-1.488941e+01	-8.659854e+01	1.582366e-02	-1.810913e-06	±0.57
8	Ca ₃ Si ₂ O ₇	RANKINITE	-312.1	8.533219e+05	-1.389306e+01	-1.213161e+02	1.564223e-02	-1.814195e-06	±0.60
9	MgAl ₂ O ₄	SPINEL	-311.0	4.912197e+05	-1.043835e+01	-6.070647e+01	1.107213e-02	-1.213724e-06	±0.42
10	CaSiO ₃	WOLLASTONITE	-309.8	3.603498e+05	-6.690462e+00	-4.577239e+01	7.145265e-03	-8.108841e-07	±0.26
11	Ca ₅ P ₃ O ₁₂ F	FLUORAPATITE	-309.1	1.411299e+06	-2.348482e+01	-2.134680e+02	2.699044e-02	-3.118360e-06	±0.84
12	Ca ₃ MgSi ₂ O ₈	MERWINITE	-308.5	9.728542e+05	-1.618741e+01	-1.392443e+02	1.804333e-02	-2.020680e-06	±0.68
13	CaAl ₂ Si ₂ O ₈	ANORTHITE	-308.3	9.529318e+05	-2.009489e+01	-1.046593e+02	2.000063e-02	-2.277610e-06	±0.84
14	CaTiSiO ₅	SPHENE	-307.4	5.922714e+05	-1.238208e+01	-6.590011e+01	1.323755e-02	-1.584733e-06	±0.61
15	Ca ₂ MgSi ₂ O ₇	AKERMANITE	-305.8	8.405318e+05	-1.269676e+01	-1.275788e+02	1.371402e-02	-1.450183e-06	±0.37
16	Al ₂ SiO ₅	KYANITE	-305.4	5.917326e+05	-0.473362e+01	-5.773181e+01	1.438186e-02	-1.670007e-06	±0.55
17	CaMgSiO ₄	MONTECELLITE	-304.9	4.818448e+05	-8.787430e+00	-6.715236e+01	9.912023e-03	-1.160460e-06	±0.34
18	CaMgSi ₂ O ₆	DIOPSIDE	-302.8	7.087080e+05	-1.484631e+01	-8.419222e+01	1.555722e-02	-1.793114e-06	±0.62
19	MgAl ₂ SiO ₆	Mg-TSCHERMAKS	-302.0	7.118491e+05	-1.538535e+01	-8.324974e+01	1.591182e-02	-1.724839e-06	±0.46
20	CaO	LIME	-301.6	1.273800e+05	-1.455968e+00	-2.479564e+01	2.33874e-03	-2.664018e-07	±0.11
21	KMg ₃ AlSi ₃ O ₁₀ F ₂	FLUORPHLOGOPITE	-300.9	1.342665e+06	-2.662439e+01	-1.797165e+02	2.659617e-02	-2.835390e-06	±1.12
22	Mg ₃ Al ₂ Si ₃ O ₁₂	PYROPE	-296.7	1.402656e+06	-3.072607e+01	-1.635848e+02	3.136489e-02	-3.582534e-06	±1.23
23	TiO ₂	RUTILE	-296.4	2.296961e+05	-3.392573e+00	-3.338416e+01	3.426975e-03	-3.572185e-07	±0.07
24	MgTiO ₃	GEIKIELITE	-296.1	3.517143e+05	-6.698669e+00	-4.773530e+01	7.802853e-03	-9.366533e-07	±0.34
25	Ca ₂ Al ₃ Si ₃ O ₁₃ H	CLINOZOISITE	-295.6	1.560747e+06	-3.510366e+01	-1.680363e+02	3.352398e-02	-3.806883e-06	±1.37
26	CaSi ₂ O ₅	CaSi-TITANITE	-293.7	5.759983e+05	-1.513590e+01	-4.985255e+01	1.548385e-02	-1.893205e-06	±0.83
27	Mg ₂ SiO ₄	FORSTERITE	-293.4	4.68652e+05	-9.176566e+00	-6.544643e+01	9.627992e-03	-1.038920e-06	±0.30
28	MgSiO ₃	ENSTATITE	-291.6	3.458775e+05	-7.257035e+00	-4.345839e+01	7.516612e-03	-8.200336e-07	±0.22
29	Ca ₅ Si ₂ CO ₁₁	SPURRITE	-291.4	1.329424e+06	-2.132758e+01	-1.950763e+02	2.422856e-02	-2.735895e-06	±0.83
30	KAlSi ₃ O ₈	MICROCLINE	-288.5	9.258438e+05	-2.002663e+01	-1.044972e+02	1.928080e-02	-2.158177e-06	±0.60
31	Ca ₅ P ₃ O ₁₃ H	HYDROXYAPATITE	-288.1	1.440061e+06	-2.948136e+01	-1.882099e+02	3.109138e-02	-3.217938e-06	±1.61
32	KAlSiO ₄	KALSILITE	-286.9	4.774370e+05	-9.688874e+00	-6.070594e+01	1.032885e-02	-1.168394e-06	±0.35
33	KAlSi ₂ O ₆	LEUCITE	-286.7	6.994399e+05	-1.448596e+01	-8.441743e+01	1.342159e-02	-1.489176e-06	±0.33
34	NaAlSi ₃ O ₈	ALBITE	-285.6	9.232283e+05	-2.012507e+01	-1.039484e+02	1.953868e-02	-2.217366e-06	±0.63
35	SiO ₂	QUARTZ	-285.5	2.222506e+05	-5.478967e+00	-1.969170e+01	4.907829e-03	-5.612476e-07	±0.18
36	NaAlSi ₂ O ₆	JADEITE	-284.7	7.011353e+05	-1.413739e+01	-9.120226e+01	1.397186e-02	-1.513554e-06	±0.45
37	MgO	PERICLASE	-284.6	1.195366e+05	-2.104910e+00	-2.122681e+01	2.564436e-03	-2.957752e-07	±0.11
38	NaAlSiO ₄	NEPHELINE	-282.9	4.759180e+05	-1.098517e+01	-5.306336e+01	1.100831e-02	-1.241451e-06	±0.32
39	Ca ₂ MnAl ₂ Si ₃ O ₁₃ H	PIEMONTE(ORDERED)	-280.0	1.513754e+06	-3.077556e+01	-1.907176e+02	2.793700e-02	-3.063564e-06	±1.09
40	CaAl ₄ Si ₂ O ₁₂ H ₂	MARGARITE	-279.0	1.443099e+06	-3.373819e+01	-1.571716e+02	3.303015e-02	-3.705003e-06	±1.07
41	Ca ₂ Al ₂ Si ₃ O ₁₂ H ₂	PREHNITE	-277.4	1.434914e+06	-3.196401e+01	-1.641458e+02	3.145388e-02	-3.528334e-06	±1.01
42	Ca ₂ FeAl ₂ Si ₃ O ₁₃ H	EPIDOTE(ORDERED)	-276.6	1.521106e+06	-3.241005e+01	-1.857802e+02	3.245728e-02	-3.668356e-06	±1.36
43	Ca ₅ Si ₂ C ₂ O ₁₃	TILLEYITE	-273.3	1.537055e+06	-2.652043e+01	-2.135833e+02	2.876369e-02	-3.172582e-06	±0.87
44	Ca ₃ Fe ₂ Si ₃ O ₁₂	ANDRADITE	-271.4	1.373546e+06	-2.766753e+01	-1.840431e+02	3.011350e-02	-3.455637e-06	±1.16
45	KMg ₂ Al ₃ Si ₂ O ₁₂ H ₂	EASTONITE	-270.2	1.348934e+06	-3.232163e+01	-1.763878e+02	3.221755e-02	-3.756024e-06	±1.09
46	Mn ₃ Al ₂ Si ₃ O ₁₂	SPESSARTINE	-268.3	1.381612e+06	-2.857798e+01	-1.787062e+02	3.043611e-02	-3.453339e-06	±1.15
47	CaFeSi ₂ O ₆	HEDENBERGITE	-268.0	6.981350e+05	-1.348390e+01	-9.302132e+01	1.450960e-02	-1.588910e-06	±0.45
48	Mg ₃ Cr ₂ Si ₃ O ₁₂	KNORRINGITE	-267.8	1.348934e+06	-2.497408e+01	-1.966352e+02	2.701188e-02	-3.090497e-06	±1.02
49	K ₂ Si ₄ O ₉	Si-WADEITE	-267.5	1.016675e+06	-2.009772e+01	-1.385171e+02	1.940090e-02	-2.058947e-06	±0.50
50	Mg ₂ Al ₂ Si ₃ O ₁₂ H ₂	TSCHERMAK-TALC	-267.2	1.400979e+06	-3.625955e+01	-1.386779e+02	3.417640e-02	-3.826802e-06	±1.51
51	KAl ₃ Si ₃ O ₁₂ H ₂	MUSCOVITE	-266.5	1.414776e+06	-3.449122e+01	-1.503871e+02	3.394598e-02	-3.828482e-06	±1.10
52	KMg ₃ Al ₃ Si ₃ O ₁₂ H ₂	PHLOGOPITE	-265.2	1.417930e+06	-3.144794e+01	-1.792887e+02	3.133804e-02	-3.671162e-06	±1.07
53	NaAl ₃ Si ₃ O ₁₂ H ₂	PARAGONITE	-264.9	1.413363e+06	-3.295879e+01	-1.610931e+02	3.236865e-02	-3.608740e-06	±0.90
54	AlSi ₂ O ₆ H	PYROPHYLLITE	-263.4	6.892108e+05	-1.659504e+01	-7.266395e+01	1.542338e-02	-1.735233e-06	±0.40
55	NaMg ₃ Al ₃ Si ₃ O ₁₂ H ₂	SODAPHLOGOPITE	-263.3	1.414930e+06	-3.155196e+01	-1.788363e+02	3.159871e-02	-3.730675e-06	±1.11
56	FeAl ₂ O ₄	HERCYNITE	-262.9	4.816286e+05	-1.048218e+01	-6.061770e+01	1.156339e-02	-1.306461e-06	±0.48
57	Mg ₃ Si ₄ O ₁₂ H ₂	TALC	-262.8	1.382604e+06	-3.420699e+01	-1.502025e+02	3.251454e-02	-3.772182e-06	±1.35
58	KMgAlSi ₄ O ₁₂ H ₂	CELADONITE	-259.8	1.390437e+06	-3.361618e+01	-1.544995e+02	3.306604e-02	-3.743581e-06	±1.09
59	NaCrSi ₂ O ₆	KOSMOCHLOR	-257.5	6.765279e+05	-1.125471e+01	-1.086861e+02	1.179478e-02	-1.267462e-06	±0.34
60	Ca ₂ FeAl ₃ Si ₃ O ₁₂ H ₂	FERRI-PREHNITE	-257.1	1.393161e+06	-3.104842e+01	-1.689573e+02	3.137888e-02	-3.535514e-06	±1.00
61	MnTiO ₃	PYROPHANITE	-256.7	3.435012e+05	-5.536905e+00	-5.436775e+01	7.170529e-03	-8.519687e-07	±0.32
62	Ca ₂ Fe ₂ Al ₃ Si ₃ O ₁₃ H	Fe-EPIDOTE	-256.5	1.478087e+06	-3.103944e+01	-1.946633e+02	3.139057e-02	-3.529827e-06	±1.35
63	MgAl ₂ SiO ₇ H ₂	Mg-CHLORITOID	-254.5	8.346090e+05	-2.320869e+01	-7.849664e+01	2.128252e-02	-2.422756e-06	±0.80
64	MnSiO ₃	PYROXMANGITE	-249.2	3.355832e+05	-6.945405e+00	-4.433766e+01	7.426667e-03	-8.613014e-07	±0.34
65	CaAl ₂ Si ₄ O ₁₄ H ₄	WAIRAKITE	-248.8	1.633745e+06	-3.834118e+01	-1.809888e+02	3.572679e-02	-4.035846e-06	±1.15
66	KAlSi ₃ O ₉ H ₂	K-CYMRITE	-247.4	1.038223e+06	-2.399976e+01	-1.183881e+02	2.236149e-02	-2.482468e-06	±0.68
67	Fe ₃ Al ₂ Si ₃ O ₁₂	ALMANDINE	-246.9	1.376944e+06	-3.016241e+01	-1.707578e+02	3.187560e-02	-3.563550e-06	±1.11
68	Al ₂ SiO ₆ H ₂	HYDROXY-TOPAZ	-244.2	7.100468e+05	-1.892466e+01	-7.372280e+01	1.753940e-02	-1.941487e-06	±0.55
69	KFeAlSi ₄ O ₁₂ H ₂	FERROCELADONITE	-243.0	1.378812e+06	-3.315759e+01	-1.566005e+02	3.278410e-02	-3.675750e-06	±1.02
70	NaFeSi ₂ O ₆	ACMITE	-241.7	6.583105e+05	-1.441096e+01	-8.729880e+01	1.476475e-02	-1.542019e-06	±0.48
71	MnAl ₂ SiO ₇ H ₂	Mn-CHLORITOID	-238.6	8.259787e+05	-2.2111654e+01	-9.285036e+01	2.038330e-02	-2.293290e-06	±0.66
72	NaAlSi ₂ O ₇ H ₂	ANALCITE	-237.6	8.176590e+05	-1.519133e+01	-1.214758e+02	1.562853e-02	-1.269436e-06	±0.38
73	CaAl ₂ Si ₂ O ₁₀ H ₄	LAWSONITE	-237.6	1.191594e+06	-3.176561e+01	-1.226891e+02	2.966229e-02	-3.109709e-06	±0.77
74	MgCr ₂ O ₄	PICROCHROMITE	-236.0	4.431007e+05	-8.938663e+00	-6.835976e+01	1.066679e-02	-1.296484e-06	±0.52
75	Mn ₂ SiO ₄	TEPHROITE	-233.4	4.489460e+05	-7.633874e+00	-7.402296e+01	8.919088e-03	-1.011800e-06	±0.33
76	KMn ₃ Al ₃ Si ₃ O ₁₂ H ₂	Mn-BIOTITE	-233.1	1.379102e+06	-2.873641e+01	-1.929652e+02	3.004383e-02	-3.637965e-06	±1.01

The condensates are sorted according to $\Delta G_{f,\text{SU}}^*(T_{\text{ref}})/N$, i.e. SUPCRTBL's Gibbs free energy of formation [kJ/mol] from the elements at reference pressure $p^* = 1$ bar and reference temperature $T_{\text{ref}} = 298.15$ K, divided by N , the sum of stoichiometric factors. ΔG_f^* are the Gibbs free energies of formation of the condensates from free atoms at $p^* = 1$ bar and temperature T , needed for Eq. (12), which are fitted according to $-\Delta G_f^*/(RT) = b_0/T + b_1 \ln(T) + b_2 + b_3 T + b_4 T^2$. The last column is the standard deviation between the generated SUPCRTBL data points and the functional fit between 100 K and 2500 K in kJ/mol.

Table D.4. continued

no	formula	name	$\Delta G_{f,SU}^{\circ}(T_{ref})/N$	b_0	b_1	b_2	b_3	b_4	σ [kJ/mol]
77	MgAl ₂ Si ₂ O ₁₀ H ₄	MAGNESIOCARPHOLITE	-232.4	1.176370e+06	-3.070094e+01	-1.290393e+02	2.805212e-02	-3.037731e-06	±0.76
78	FeTiO ₃	ILMENITE	-230.9	3.435788e+05	-5.480155e+00	-5.476246e+01	6.285519e-03	-7.119639e-07	±0.24
79	Al ₂ O ₃	DIASPORE	-230.3	2.443624e+05	-7.279776e+00	-2.289286e+01	6.441950e-03	-5.931350e-07	±0.12
80	FeAl ₂ SiO ₇ H ₂	Fe-CHLORITOID	-228.9	8.258670e+05	-2.327161e+01	-7.938651e+01	2.116909e-02	-2.404658e-06	±0.79
81	Mg ₇ Si ₂ O ₁₄ H ₆	PHASEA	-228.0	1.656847e+06	-4.230253e+01	-2.158169e+02	4.247591e-02	-4.797719e-06	±1.52
82	CaCO ₃	CALCITE	-226.0	3.411928e+05	-6.591620e+00	-4.368617e+01	6.352508e-03	-6.931829e-07	±0.23
83	Mg ₃ Si ₂ O ₉ H ₄	LIZARDITE	-224.4	1.054418e+06	-2.910595e+01	-1.166171e+02	2.654429e-02	-3.029452e-06	±0.88
84	Al ₂ Si ₂ O ₉ H ₄	KAOLINITE	-223.7	1.050606e+06	-2.819000e+01	-1.056585e+02	1.966983e-02	-2.504474e-06	±0.88
85	FeSiO ₃	FERROSILITE	-223.5	3.361566e+05	-6.749696e+00	-4.728394e+01	7.277447e-03	-7.573112e-07	±0.14
86	CaSO ₄	ANHYDRITE	-220.4	3.449608e+05	-9.113763e+00	-4.690825e+01	8.600984e-03	-6.567852e-07	±0.24
87	KFe ₃ AlSi ₃ O ₁₂ H ₂	ANNITE	-218.0	1.386502e+06	-3.005579e+01	-1.891851e+02	3.068660e-02	-3.467367e-06	±0.85
88	CaMgC ₂ O ₆	DOLOMITE	-216.4	6.676592e+05	-1.445077e+01	-8.264391e+01	1.429423e-02	-1.546931e-06	±0.40
89	CaAl ₂ Si ₄ O ₁₆ H ₈	LAUMONTITE	-216.3	1.868803e+06	-4.628609e+01	-2.147229e+02	4.189556e-02	-4.685634e-06	±1.32
90	FeAl ₂ Si ₂ O ₁₀ H ₄	FERROCARPHOLITE	-214.1	1.165522e+06	-3.037176e+01	-1.316984e+02	2.831269e-02	-3.060630e-06	±0.76
91	Fe ₃ Si ₄ O ₁₂ H ₂	MINNESOTAITE	-213.0	1.349960e+06	-3.399941e+01	-1.519206e+02	3.265094e-02	-3.538871e-06	±1.27
92	Cr ₂ O ₃	ESKOLAITE	-211.3	3.216934e+05	-2.887880e+00	-7.289829e+01	4.520548e-03	-5.320269e-07	±0.21
93	MgCO ₃	MAGNESITE	-205.5	3.255978e+05	-7.798617e+00	-3.888094e+01	7.661015e-03	-8.339379e-07	±0.22
94	KCl	SYLVITE	-204.5	7.801979e+04	4.147547e-01	-3.263640e+01	1.037051e-03	6.931110e-09	±0.00
95	Fe ₂ TiO ₄	ULVOSPINEL	-199.7	4.536963e+05	-9.821801e+00	-5.985428e+01	1.017629e-02	-6.533094e-07	±0.73
96	Fe ₂ SiO ₄	FAYALITE	-197.0	4.495737e+05	-9.271411e+00	-6.615000e+01	1.041250e-02	-1.092031e-06	±0.38
97	NaCl	HALITE	-192.3	7.717022e+04	3.607723e-01	-3.269074e+01	9.594929e-04	1.834797e-08	±0.00
98	MgFe ₂ O ₄	MAGNESIOFERRITE	-190.1	4.092716e+05	-8.152176e+00	-7.527872e+01	1.095039e-02	-1.255442e-06	±0.37
99	CaFeC ₂ O ₆	ANKERITE	-182.0	6.574137e+05	-1.363162e+01	-8.757014e+01	1.322217e-02	-1.409341e-06	±0.35
100	MnO	MANGANOSITE	-181.7	1.103123e+05	-6.866758e-01	-2.930136e+01	1.564511e-03	-1.440846e-07	±0.04
101	NaAlCO ₅ H ₂	DAWSONITE	-178.3	5.687719e+05	-2.621938e+01	1.013648e+01	7.713633e-04	-7.242316e-08	±0.01
102	Mn ₂ O ₃	BIXBYITE	-176.5	2.728569e+05	-2.903180e+00	-6.914515e+01	4.215259e-03	-2.567819e-07	±0.03
103	MgO ₂ H ₂	BRUCITE	-167.0	2.396506e+05	-7.315243e+00	-3.115064e+01	7.048891e-03	-8.256913e-07	±0.28
104	Fe ₃ Si ₂ O ₉ H ₄	GREENALITE	-166.8	1.022771e+06	-2.804547e+01	-1.240261e+02	2.772103e-02	-3.109114e-06	±1.01
105	AlO ₃ H ₃	GIBBSITE	-165.0	3.602374e+05	-1.611001e+01	-1.217224e+01	1.567462e-02	-4.410396e-07	±0.10
106	MnCO ₃	RHODOCHROSITE	-163.8	3.159405e+05	-6.661948e+00	-4.500950e+01	6.514539e-03	-6.894651e-07	±0.18
107	Fe ₂ O ₃	HEMATITE	-148.8	2.876529e+05	-6.810027e+00	-4.976293e+01	8.724692e-03	-1.038569e-06	±0.40
108	Fe ₃ O ₄	MAGNETITE	-144.6	4.019320e+05	-8.960632e+00	-7.101785e+01	1.191424e-02	-1.404312e-06	±0.46
109	FeCO ₃	SIDERITE	-137.8	3.161250e+05	-6.980649e+00	-4.434616e+01	6.585898e-03	-6.958439e-07	±0.17
110	FeO	FERROPERICLASE	-125.8	1.121860e+05	-2.152325e+00	-2.083896e+01	2.956682e-03	-3.223296e-07	±0.15
111	FeO ₂ H	GOETHITE	-122.8	2.026749e+05	-4.655123e+00	-3.861734e+01	5.315705e-03	-5.712016e-07	±0.15
112	NiO	NICKEL	-105.7	1.103349e+05	-8.363217e-01	-3.178273e+01	1.348578e-03	-1.056011e-07	±0.06
113	CuO	TENORITE	-64.2	8.900429e+04	-1.587221e+00	-2.477537e+01	2.420378e-03	-2.394143e-07	±0.14
114	FeS ₂	PYRITE	-53.5	1.364603e+05	-3.370114e+00	-3.495254e+01	3.621202e-03	-2.916809e-07	±0.19
115	FeS	TROILITE	-50.7	9.450871e+04	-2.495865e+00	-1.829773e+01	3.174381e-03	-3.069607e-07	±0.13
116	Cu ₂ O	CUPRITE	-49.4	1.317125e+05	-6.268504e-01	-4.560941e+01	2.920620e-03	-3.213970e-07	±0.08
117	Cu	COPPER	-0.1	4.065093e+04	3.278646e-02	-1.652274e+01	5.549892e-04	-2.618359e-08	±0.03
118	S	SULPHUR	-0.1	3.327240e+04	-4.494524e-01	-1.393588e+01	1.047391e-03	-9.902057e-08	±0.02
119	Ni	NICKEL	-0.1	5.178431e+04	-5.284788e-02	-1.837439e+01	6.292585e-04	-3.003321e-08	±0.03
120	Fe	IRON	-0.0	4.989867e+04	-6.889399e-01	-1.461748e+01	1.161476e-03	-5.259868e-08	±0.04
121	C	GRAPHITE	-0.0	8.563846e+04	-2.345056e+00	-3.634772e+00	1.828533e-03	-1.986659e-07	±0.03

Table D.5. Fit coefficients for the atomic corrections $\Delta G_{\text{corr}}^{\circ}(T)$ to convert the SUPCRTBL data to our reference states of neutral atoms, see Eq. (D.6), derived from the NIST-JANAF database.

atom	a_0	a_1	a_2	a_3	a_4	error [kJ/mol]
H	-2.78172E+4	2.49885	-2.93973E+0	1.14321E-6	-1.34658E-10	±0.001
Li	-1.94521E+4	2.52691	-2.27778E-1	-2.14827E-5	2.96822E-09	±0.039
C	-8.56514E+4	2.52074	2.14377E+0	-2.28858E-5	3.90851E-09	±0.002
N	-5.95408E+4	2.50073	1.69028E+0	-2.07541E-6	4.79627E-10	±0.004
O	-3.28428E+4	2.71152	1.26593E+0	-1.35177E-4	1.26727E-08	±0.027
F	-1.23701E+4	2.76996	6.19482E-1	-1.28288E-4	9.91217E-09	±0.013
Na	-1.40060E+4	2.48726	1.80092E+0	2.38611E-5	-3.17527E-09	±0.058
Mg	-1.81146E+4	2.51860	1.00708E+0	-8.33362E-6	3.74599E-10	±0.033
Al	-3.98785E+4	2.65608	2.03015E+0	-9.87400E-5	8.59574E-09	±0.032
Si	-5.39603E+4	2.82697	1.34719E+0	-2.23157E-4	2.56696E-08	±0.062
P	-3.87736E+4	2.54304	2.61689E+0	-4.90659E-5	8.96721E-09	±0.045
S	-3.36266E+4	2.82008	1.33086E+0	-7.74986E-5	0.00000E+00	±0.17
Cl	-1.78181E+4	2.58828	2.50684E+0	9.72935E-5	-1.51727E-08	±0.073
K	-1.22688E+4	2.55355	2.20391E+0	-5.36966E-5	8.24159E-09	±0.050
Ca	-2.21218E+4	2.55750	1.53540E+0	-7.09263E-5	1.21389E-08	±0.067
Ti	-5.71391E+4	3.20557	4.48870E-1	-5.18014E-4	7.00646E-08	±0.10
Cr	-4.78986E+4	2.56936	3.82072E+0	-1.27308E-4	3.48156E-08	±0.041
Mn	-3.44654E+4	2.52906	3.96202E+0	-2.51476E-5	3.59496E-09	±0.046
Fe	-5.00408E+4	3.17358	6.57213E-1	-2.96787E-4	3.12523E-08	±0.29
Ni	-5.19591E+4	2.80126	3.06725E+0	1.48777E-4	-2.21945E-08	±0.084
Cu	-4.10420E+4	2.54198	3.01655E+0	-5.98838E-5	1.27558E-08	±0.062
Zr	-7.39272E+4	2.79313	3.25202E+0	1.26147E-4	1.15173E-08	±0.072

$$-\Delta G_{\text{corr}}^{\circ}/(RT) = a_0/T + a_1 \ln(T) + a_2 + a_3 T + a_4 T^2.$$