Dragon Engineering: A Speculative Biology Thought Experiment

Alexi G. Choueiri
Independent Researcher
alexichoueiri@gmail.com

November 7, 2026

Abstract. A flying, six-limbed, fire-breathing dragon breaks real biology in three places, and flight is the most quantifiable of them. Scaling the best-studied giant flier, the azhdarchid pterosaur *Quetzalcoatlus* (∼10 m span, ∼250 kg), shows that required flight power grows as $m^{7/6}$ while muscle power grows only as about $m^{5/6}$. Under stated assumptions, sustained flapping caps out between roughly 16 and 90 kg; short anaerobic bursts can lift a horse-sized animal (∼500 kg) only with at least 25% of body mass in flight muscle, after which it must soar. An elephant-sized dragon (∼5,000 kg) has about half the burst power it needs and cannot fly. A sensitivity analysis shows that the 5,000 kg result holds under every combination of assumptions tested, while the 500 kg result depends on how much burst power muscle can deliver. The model is released as open Python code. The other two problems resist numbers. A third limb pair would require a new limb field in the embryo, which no vertebrate has evolved in roughly 370 million years of tetrapod history. Fire is chemically plausible through known biological reactions, but needs dedicated tissue protection and an ignition source no animal possesses. This paper is a thought experiment: it proposes no organism and no laboratory work.

Keywords: speculative biology; dragon; wyvern; flight scaling; allometry; pterosaur; *Quetzalcoatlus*; flight power; limb development; developmental genetics; Hox genes; reproducible model; ancient DNA; bombardier beetle; archosaurs; thought experiment

1. Introduction and design specification

The question is not "can we build a dragon" but "which parts of a dragon does known biology forbid, and by how much." Treating the dragon as an engineering specification turns a myth into a set of testable constraints. The working specification is:

1. **Powered flight** from a standing start, without cliffs or headwinds.
2. **Six limbs:** four legs plus a separate pair of wings.
3. **Fire:** a directed, ignited stream from the mouth.
4. **Large size:** at least horse-sized (∼500 kg), and ideally larger.
5. **Scaled skin** and a reptile-like appearance.

Each section asks what real organism comes closest to a trait, what physics or development limits it, and how far the specification overshoots that limit. Numbers come from published fossil and flight studies; where the paper adds its own estimates, the assumptions are stated.

Scope. This is a speculative-biology exercise. It discusses genes and mechanisms only at the level of published, general science, and contains no protocols or designs for modifying real organisms.

2. Flight scaling

Flight is where a dragon fails by numbers. With 25% of body mass in flight muscle, burst power drops below what flapping needs somewhere between 500 and 1,000 kg.




Figure 1. Artist's impression of the proposed wyvern in its quadrupedal stance, wings folded. AI-generated image; not a scientific reconstruction.




Figure 2. Three body plans compared (side view, not to scale): the classic six-limbed dragon, the four-limbed wyvern, and the real azhdarchid pterosaur *Quetzalcoatlus*. The wyvern keeps the pterosaur's four-limb plan, which is why later sections treat it as the plausible design.

2.1 Real benchmarks

Table 1 lists the largest fliers for which flight has been modeled. Mass estimates for *Quetzalcoatlus* are disputed: Henderson (2010) proposed 544 kg and flightlessness, which Witton and Habib (2010) attribute to an oversized torso model. This paper uses their 200–250 kg range.

Table 1. Benchmark fliers.

Animal	Mass	Span	How it gets airborne
Quetzalcoatlus northropi (pterosaur)	200–250 kg	10–11 m	Proposed quadrupedal vault from a standing start, then soaring (Habib 2008; Witton & Habib 2010)
Argentavis magnificens (extinct bird)	∼70 kg	∼7 m	Likely needed slopes, wind or a run; soared on updrafts (Chatterjee et al. 2007)
Kori bustard (living bird)	∼18 kg	∼2.5 m	Runs to take off (Ledford 2007)
Wandering albatross (living bird)	∼8 kg	∼3.5 m	Long run into wind; dynamic soaring (Witton & Habib 2010)

2.2 The scaling argument

Scale an animal up with the same shape (isometric scaling). Wing area grows as $m^{2/3}$, so wing loading and flight speed rise. The power needed to fly rises faster than mass:

$$P_{\text{req}} \propto m\, g\, V \propto m^{7/6}. \tag{1}$$

Muscle power is muscle mass times work per wingbeat times wingbeat frequency f. Frequency falls as animals grow, roughly as $m^{-1/6}$ under isometry:

$$P_{\text{av}} = \phi\, m\, Q\, f \propto m^{5/6}, \tag{2}$$

where ϕ is the flight-muscle fraction and Q is work per kilogram of muscle per beat. The margin $P_{\text{av}}/P_{\text{req}}$ therefore falls as $m^{-1/3}$: **every eightfold increase in mass halves the power margin** (Figure 3).




Figure 3. Ratio of muscle power available to power needed for flapping flight, for 25% flight muscle, scaled isometrically from *Quetzalcoatlus*. Burst power falls below the requirement between 500 and 1,000 kg. Author's model; methods in Section 2.3.

2.3 Methods

The model scales a flier isometrically from the *Quetzalcoatlus* planform of Witton (2008), as tabulated by Witton and Habib (2010), so that wingspan $b \propto m^{1/3}$ and wing area $S \propto m^{2/3}$. All parameters are listed in Table 2.

Power required. Following the approach of Pennycuick (2008), the mechanical power for level flight at speed V is the sum of induced and parasite (body) power, with profile power added as a fixed multiple X of that sum:

$$P_{\mathrm{req}}(V) = (1+X)\left[\frac{k\,(mg)^2}{2\rho V S_d} + \tfrac{1}{2}\rho V^3 S_b C_{Db}\right], \tag{3}$$

where $S_d = \pi b^2/4$ is the actuator-disc area and $S_b = 0.00813\,m^{0.666}$ is the body frontal area (m^2, with m in kg). P_{req} is taken at the speed that minimizes it.

Power available. Muscle power is $P_{\mathrm{av}} = \phi\, m\, Q\, f$, where the wingbeat frequency follows Pennycuick's empirical relation

$$f = m^{3/8}\, g^{1/2}\, b^{-23/24}\, S^{-1/3}\, \rho^{-3/8}. \tag{4}$$

Calibration. The work per kilogram of muscle per beat, Q, is not well known for large fliers. It is set so that an animal with a bird-like 17% flight-muscle fraction reaches its sustained-flight limit at 16 kg, close to the heaviest living fliers. This gives $Q = 33.2\,\mathrm{J\,kg^{-1}}$ for sustained (aerobic) output. Burst (anaerobic) output is taken as 2.5 times sustained output.

Mass limit. For each muscle fraction, the mass limit is the largest m at which $P_{\mathrm{av}} \geq P_{\mathrm{req}}$, found by bisection. The model is implemented in `dragon_flight_model.py` (see Code availability), which reproduces every number in this section.

Table 2. Model parameters.

Parameter	Value	Source or basis
Reference mass, span, wing area	259 kg, 9.64 m, 11.36 m^2	*Quetzalcoatlus*; Witton & Habib (2010), Table 1
Air density ρ	1.225 $\mathrm{kg\,m^{-3}}$	Sea level
Induced-drag factor k	1.2	Typical value in Pennycuick-style models
Body drag coefficient C_{Db}	0.1	Assumed
Profile power ratio X	1.2	Assumed
Maximum lift coefficient	2.2	Membrane wing; Witton & Habib (2010)
Sustained-limit anchor	16 kg at 17% muscle	Heaviest living fliers
Burst factor	2.5	Assumed; tested in Section 2.5

2.4 Results

Table 3 gives the resulting mass limits for three flight-muscle fractions, and Table 4 shows what a dragon of each size would look like.

Table 3. Estimated mass limits for flapping flight by flight-muscle fraction.

Flight muscle	Sustained flight limit	Burst flight limit
17% (typical bird)	∼16 kg	∼250 kg
25%	∼50 kg	∼800 kg
30%	∼90 kg	∼1,400 kg

Table 4. Dragon geometry at three sizes, scaled isometrically from *Quetzalcoatlus*. Stall speed assumes a maximum lift coefficient of 2.2.

Mass	Wingspan	Wing area	Wing loading	Stall speed	Wingbeats
500 kg	12 m	17.6 m^2	279 N/m^2	14 m/s	1.1 s^{-1}
1,000 kg	15 m	28 m^2	351 N/m^2	16 m/s	0.9 s^{-1}
5,000 kg	26 m	82 m^2	600 N/m^2	21 m/s	0.7 s^{-1}

2.5 Sensitivity analysis

The two least certain inputs are the burst factor and the sustained-flight anchor used to calibrate Q. Because the power margin falls as $m^{-1/3}$, the mass limit scales as the *cube* of any multiplier on muscle output: raising the burst factor from 2 to 3 multiplies the limit by $1.5^3 \approx 3.4$. Table 5 shows the burst mass limit across these inputs.

Table 5. Sensitivity of the burst flight mass limit (kg). Left: 25% flight muscle, varying the calibration anchor and burst factor. Right: 16 kg anchor, varying flight-muscle fraction and burst factor.

Anchor	×2.0	×2.5	×3.0
12 kg	306	597	1,033
16 kg	408	797	1,377
22 kg	561	1,095	1,893

Muscle	×2.0	×2.5	×3.0
17%	128	250	433
25%	408	797	1,377
30%	705	1,377	2,380
35%	1,119	2,187	3,780

Two conclusions hold across every combination tested. First, no case lets a 5,000 kg dragon fly: the most generous one (35% muscle, burst factor 3) stops at about 3,800 kg. Second, sustained flapping never reaches horse size. The 500 kg result is less secure: it holds in most cases with at least 25% muscle, but fails if burst output is only twice sustained output and the anchor is 16 kg or lower.

2.6 Implications for the design

A horse-sized dragon is plausible only as a burst-launch soarer. It would flap hard for about a minute, then need thermals or ridge lift within 1–2 km, as Witton and Habib (2010) propose for giant pterosaurs (Figure 4). An elephant-sized dragon has about half the burst power it needs.

Six limbs make this worse. Pterosaurs likely launched by vaulting on their wings, using the flight muscles as legs (Habib 2008). A dragon's wings are a separate pair, so its legs cannot borrow flight muscle, and its four legs are dead weight in the air.

3. Six limbs

The six-limbed body plan is the dragon's least biological trait. Every land vertebrate descends from four-limbed ancestors, and no lineage has added a third pair in roughly 370 million years of tetrapod history.

3.1 Why vertebrates stop at four

- **Limb fields are set early.** In the embryo, Hox genes mark positions along the body. Only two regions become competent to grow limbs: one for the forelimbs, one for the hindlimbs.
- **Each pair has its own identity genes.** *Tbx5* is associated with forelimb identity, and *Tbx4* with *Pitx1* with hindlimb identity. A third pair would need a third identity, or a duplicate of one.
- **Limbs are wired into everything.** Each pair needs its own girdle of bone, muscles, nerves from the spinal cord, and blood supply. A new pair is not one change but dozens.




Figure 4. Artist's impression of the soaring phase after a burst launch, with a person below for scale. AI-generated image; not a scientific reconstruction.

3.2 Hints that it is not impossible

Classic chick-embryo experiments showed that a local growth-factor signal in the flank can trigger an extra limb (Cohn et al. 1995): flank tissue has latent limb-forming ability. Extra limbs (polymelia) also occur as rare developmental defects in many vertebrates, and in frogs can be caused by the trematode parasite *Ribeiroia* (Johnson et al. 1999). These extra limbs are malformed and non-functional. Nothing in them suggests a path to a working third pair with its own girdle and flight muscles.

3.3 Developmental genetics of dragon-like traits

Several dragon traits have counterparts whose developmental genetics has been studied in real animals (Table 6). A recurring theme is that new structures evolve by changing when and where existing signals act, not by inventing new genes. Bat wings, a third independent origin of vertebrate flight, show this clearly: the wing membrane persists because signals that normally remove tissue between the digits are blocked (Weatherbee et al. 2006), and the long wing digits reflect increased Bmp2 signaling in the forelimb (Sears et al. 2006). The table is a summary of published descriptive biology, not a design.

3.4 Design conclusion

The biologically honest dragon is a **wyvern**: four limbs, with the forelimbs as wings. That is the pterosaur plan, and Section 2 shows it is also better for launching. If six limbs are kept, they are the one trait in the specification with no known evolutionary or developmental route.

4. Fire

Fire-breathing is chemically less absurd than six limbs, but no known animal ignites anything. The trait splits into three problems: fuel, ignition, and survival.

4.1 Fuel: animals already make flammable gas

Methane is made by gut microbes in cattle and many other animals; a dragon could store it in a gut chamber, as some fish store gas in a swim bladder. *Hydrogen* is also a by-product of gut fermentation. Dickinson (1979) proposed a different source: acid dissolving bone releases hydrogen, which the dragon stores for buoyancy and vents as fire. The buoyancy idea fails on numbers. A cubic meter of hydrogen lifts only about 1.1 kg at sea level, so a 500 kg dragon would need around 450 m^3 of gas, a sac larger than a house.

Table 6. Published developmental genetics relevant to dragon-like traits.

Trait	Known genetic basis in real animals	Source
Forelimb formation	*Tbx5* is required for forelimb bud formation and outgrowth (mouse, chick)	Rallis et al. (2003)
Hindlimb identity	*Pitx1* acts upstream of *Tbx4*; misexpressing *Pitx1* in chick wing buds gives partial leg-like features	Logan & Tabin (1999)
Extra limb	FGF signaling in the flank can induce an additional limb in chick embryos	Cohn et al. (1995)
Wing membrane	Interdigital webbing retained in bat wings by Gremlin and FGF signals that block cell death	Weatherbee et al. (2006)
Elongated wing digits	Higher Bmp2 expression and signaling in bat forelimb digits	Sears et al. (2006)
Scales	Reptile scales develop from placodes, sharing ancestry with feathers and hair	Di-Poï & Milinkovitch (2016)
Air sacs and pneumatic bone	Genetic basis poorly characterized	–
Fire organs	No known genetic basis	–

4.2 Delivery: the bombardier beetle

The bombardier beetle mixes two stored chemicals with enzymes in a hardened chamber. The reaction is exothermic and sprays a near-boiling, pulsed jet from its abdomen (Arndt et al. 2015). It shows that animals can run a hot reaction safely in a dedicated organ and aim the output. It is a hot spray, not a flame.

4.3 Ignition: the missing piece

No animal produces a flame or a spark. The options are speculative: an electric discharge (one electric eel species, *Electrophorus voltai*, discharges up to 860 V; de Santana et al. 2019), friction against mineralized tissue, or a catalyst that triggers combustion. Each would be a new organ with no clear evolutionary precursor. This paper leaves ignition as an open question rather than proposing a mechanism.

4.4 Survival: not burning yourself

- The mouth and throat would need a heat-resistant lining, perhaps thick keratin or mineralized tissue.
- Flame would have to start outside the mouth, so the gas ignites in open air rather than in the airway.
- Cells would rely on heat-shock proteins, which all animals use to repair heat damage.

Verdict: fuel and delivery have real biological analogs; ignition has none.

5. Phylogenetic placement and physiology

A dragon belongs among the archosaurs, the group containing crocodilians, birds, and extinct dinosaurs and pterosaurs. It is the only lineage that has produced both giant fliers and large armored reptiles.

5.1 Why archosaurs

- **Scales:** crocodilians have them all over, and birds keep them on their feet.
- **Flight:** two archosaur groups evolved flight independently, pterosaurs and birds, and pterosaurs reached the largest flying sizes known.
- **Lungs:** birds breathe with a flow-through lung and air sacs, which extracts oxygen more efficiently than mammal lungs. Alligators also show one-way airflow in their lungs (Farmer & Sanders 2010), so the trait may be ancestral to the group.

- **Light bones:** birds and pterosaurs have air-filled (pneumatic) bones, connected to that respiratory system.

5.2 Physiology the flight budget demands

- **Warm-blooded metabolism.** A one-minute burst at the power levels in Section 2 requires a high resting metabolism and fast recovery. Pterosaurs and birds were both likely warm-blooded.
- **A large heart and high blood pressure** to supply flight muscle that makes up a quarter or more of body mass.
- **Insulation.** Pterosaurs had hair-like filaments called pycnofibers. A scaled dragon would lose heat fast in flight unless scales are limited to the body and legs.

The closest real template is therefore a large azhdarchid pterosaur with a crocodilian-style scaled hide over the body.

6. Parts list: dragon traits mapped to real biology

Three of eight traits have close real analogs; two have none (Table 7). The gaps are the paper's main result.

Table 7. Dragon traits, their closest real analogs, and whether a known biological path exists.

Trait	Closest real analog	Biology involved	Known path?
Scaled hide	Crocodilians; bird feet	Keratin scale development	Yes
Flow-through lungs	Birds; alligators	Air sacs, one-way airflow	Yes
Pneumatic bones	Birds; pterosaurs	Air sacs invading bone	Yes
Giant wings	*Quetzalcoatlus*	Membrane or feather wing on forelimb	Partly: roughly 250–800 kg limit (Section 2)
Flammable gas	Gut fermentation in cattle	Methane- and hydrogen-producing microbes	Partly: storage and release are new
Hot chemical jet	Bombardier beetle	Enzyme-driven exothermic reaction	Partly: hot spray, not flame
Ignition	Electric eel (power source only)	None known	No
Third limb pair	Rare malformations only	Limb-field patterning (Hox, Tbx genes)	No

7. Resurrection versus design

An alternative to designing a dragon is recovering one: finding preserved DNA from its closest real relatives, the dinosaurs, perhaps frozen in ice or trapped in amber. This route fails for a physical reason, not a technical one.

7.1 DNA does not last

Allentoft et al. (2012) measured DNA decay in 158 radiocarbon-dated bones of moa, extinct giant birds of New Zealand, and estimated a half-life of about 521 years for the DNA backbone in bone. Extrapolating to an ideal $-5\,^{\circ}\mathrm{C}$, they estimated that DNA fragments would become too short to read after roughly 1.5 million years, and that essentially all backbone bonds would be broken by about 6.8 million years. Non-avian dinosaurs died out about 66 million years ago, more than 40 times beyond the readable limit. Freezing slows decay but cannot stop it over that span, and the same arithmetic rules out dinosaur DNA from blood meals of insects preserved in amber.

7.2 What has actually been recovered

The oldest genetic material known comes from cold, frozen settings and is far younger than any dinosaur. Genome-wide data have been recovered from mammoth teeth more than a million years old preserved in Siberian permafrost (van der Valk et al. 2021). Environmental DNA about two million years old, recovered from sediments of the Kap København Formation in northern Greenland, records an ancient ecosystem including mastodon, reindeer, hares, and birch and poplar trees (Kjær et al. 2022). These are the practical limits of the field: ice-age animals, not Mesozoic reptiles.

7.3 Reverse evolution in living dinosaurs

Birds descend from theropod dinosaurs, and some ancestral developmental programs remain latent in their embryos. Bhullar et al. (2015) used small-molecule inhibitors to block signaling from a facial zone unique to birds in chicken embryos. The developing beak and palate shifted toward a snout-like configuration resembling small non-avian dinosaurs such as *Velociraptor* and *Archaeopteryx*.

This connects to the question that motivated this paper: whether a genome can be worked out from an animal's appearance. It cannot, because a photograph or a fossil does not contain the information. What can sometimes be done is to uncover ancestral body shapes that living relatives still encode. That approach reshapes existing structures; it does not add organs, so it offers no route to a third limb pair or a fire gland.

7.4 Verdict

Resurrection is ruled out for dinosaurs by the chemistry of DNA decay. Any dragon, even as a thought experiment, has to be designed from the biology of living animals, which is the approach taken in this paper.

8. Limitations

- **The flight model is crude.** It uses isometric scaling, a single wing shape, and an assumed burst factor of 2.5. Real giants scale allometrically, and pterosaur wings differed from bird wings. The limits should be read as order-of-magnitude.
- **Fossil inputs are uncertain.** *Quetzalcoatlus* is known mostly from wing fragments, and published mass estimates span from about 70 kg to 544 kg.
- **Muscle performance is extrapolated.** Work per wingbeat was calibrated to living birds and may not hold for a much larger animal.
- **Fire and limbs are qualitative.** Those sections identify constraints but do not model them.

9. Conclusion

A dragon that fits known biology is a wyvern: a four-limbed archosaur about the size of a horse, with a 12 m wingspan, at least a quarter of its mass in flight muscle, and a flight style of short powered bursts followed by soaring (Figure 5). It could plausibly carry a scaled hide and store flammable gut gas. It could not breathe fire without an ignition organ that no animal has, and it could not keep six limbs. The traits that make a dragon iconic are exactly the ones biology rules out.

Code and data availability

All calculations in Section 2 are reproduced by the Python script `dragon_flight_model.py` that accompanies this paper. It requires only Python 3.8 or later (matplotlib is optional, for Figure 3). Running `python3 dragon_flight_model.py` prints Tables 3, 4, and 5 and the data behind Figure 3. All other data are taken from the cited literature.




Figure 5. The proposed design, summarizing the paper's findings (not to scale).

AI use disclosure

AI tools assisted with drafting, literature search, calculations, figures, and typesetting. Figures 1 and 4 are AI-generated artist's impressions.

Disclaimer

This work is provided "as is" for educational and discussion purposes, without warranty of any kind. The author accepts no liability for errors, omissions, or any use made of this material. Nothing here is a protocol, design, or recommendation for modifying any organism.

References

Allentoft, M. E., Collins, M., Harker, D., Haile, J., Oskam, C. L., Hale, M. L., Campos, P. F., Samaniego, J. A., Gilbert, M. T. P., Willerslev, E., Zhang, G., Scofield, R. P., Holdaway, R. N., & Bunce, M. (2012). The half-life of DNA in bone: Measuring decay kinetics in 158 dated fossils. *Proceedings of the Royal Society B*, 279(1748), 4724–4733. https://doi.org/10.1098/rspb.2012.1745

Arndt, E. M., Moore, W., Lee, W.-K., & Ortiz, C. (2015). Mechanistic origins of bombardier beetle (Brachinini) explosion-induced defensive spray pulsation. *Science*, 348(6234), 563–567. https://doi.org/10.1126/science.1261166

Bhullar, B.-A. S., et al. (2015). A molecular mechanism for the origin of a key evolutionary innovation, the bird beak and palate, revealed by an integrative approach to major transitions in vertebrate history. *Evolution*. https://doi.org/10.1111/evo.12684

Chatterjee, S., Templin, R. J., & Campbell, K. E. (2007). The aerodynamics of *Argentavis*, the world's largest flying bird from the Miocene of Argentina. *Proceedings of the National Academy of Sciences*, 104(30), 12398–12403. https://doi.org/10.1073/pnas.0702040104

Cohn, M. J., Izpisúa-Belmonte, J.-C., Abud, H., Heath, J. K., & Tickle, C. (1995). Fibroblast growth factors induce additional limb development from the flank of chick embryos. *Cell*, 80, 739–746.

de Santana, C. D., et al. (2019). Description of three electric eel species, including *Electrophorus voltai*. *Nature Communications*. Reported at https://phys.org/news/2019-09-species-electric-eel-highest-voltage.html

Di-Poï, N., & Milinkovitch, M. C. (2016). The anatomical placode in reptile scale morphogenesis indicates shared ancestry among skin appendages in amniotes. *Science Advances*, 2(6), e1600708. https://doi.org/10.1126/sciadv.1600708

Dickinson, P. (1979). *The Flight of Dragons* (W. Anderson, Illus.). Pierrot Publishing.

Farmer, C. G., & Sanders, K. (2010). Unidirectional airflow in the lungs of alligators. *Science*, 327(5963), 338–340. https://doi.org/10.1126/science.1180219

Habib, M. B. (2008). Comparative evidence for quadrupedal launch in pterosaurs. *Zitteliana*, B28, 159–166.

Henderson, D. M. (2010). Pterosaur body mass estimates from three-dimensional mathematical slicing. *Journal of Vertebrate Paleontology*, 30(3), 768–785. https://doi.org/10.1080/02724631003758334

Johnson, P. T. J., Lunde, K. B., Ritchie, E. G., & Launer, A. E. (1999). The effect of trematode infection on amphibian limb development and survivorship. *Science*, 284(5415), 802–804. https://doi.org/10.1126/science.284.5415.802

Kjær, K. H., et al. (2022). A 2-million-year-old ecosystem in Greenland uncovered by environmental DNA. *Nature*. https://doi.org/10.1038/s41586-022-05453-y

Ledford, H. (2007). Giant bird was a glider. *Nature News*. https://doi.org/10.1038/news070702-3

Logan, M., & Tabin, C. J. (1999). Role of *Pitx1* upstream of *Tbx4* in specification of hindlimb identity. *Science*, 283(5408), 1736–1739. https://doi.org/10.1126/science.283.5408.1736

Pennycuick, C. J. (2008). *Modelling the Flying Bird*. Academic Press (Elsevier). ISBN 978-0-12-374299-5.

Rallis, C., et al. (2003). *Tbx5* is required for forelimb bud formation and continued outgrowth. *Development*, 130(12), 2741–2751. https://doi.org/10.1242/dev.00473

Sears, K. E., Behringer, R. R., Rasweiler, J. J., & Niswander, L. A. (2006). Development of bat flight: Morphologic and molecular evolution of bat wing digits. *Proceedings of the National Academy of Sciences*, 103(17), 6581–6586. https://doi.org/10.1073/pnas.0509716103

van der Valk, T., Pečnerová, P., Díez-del-Molino, D., et al. (2021). Million-year-old DNA sheds light on the genomic history of mammoths. *Nature*, 591(7849), 265–269. https://doi.org/10.1038/s41586-021-03224-9

Weatherbee, S. D., Behringer, R. R., Rasweiler, J. J., & Niswander, L. A. (2006). Interdigital webbing retention in bat wings illustrates genetic changes underlying amniote limb diversification. *Proceedings of the National Academy of Sciences*, 103(41), 15103–15107. https://doi.org/10.1073/pnas.0604934103

Witton, M. P. (2008). A new approach to determining pterosaur body mass and its implications for pterosaur flight. *Zitteliana*, B28, 143–158.

Witton, M. P., & Habib, M. B. (2010). On the size and flight diversity of giant pterosaurs, the use of birds as pterosaur analogues and comments on pterosaur flightlessness. *PLoS ONE*, 5(11), e13982. https://doi.org/10.1371/journal.pone.0013982