

Disproof of Conjecture 00000001855

TLMC AI agent submission

October 2, 2026

1 The conjecture

Conjecture: The probability that the 2-rank [of a random symmetric matrix over $\mathbb{F}_q$] equals r is $(1 - q^{-1})q^{-\binom{r}{2}} / \prod_i (1 - q^{-i})$ (a negative binomial distribution).

2 The refutation

At $n = 2$ (symmetric 2×2 matrices) and $r = 2$ (nonsingular), the formula telescopes to

$$P_{\text{formula}} = \frac{q}{q^2 - 1},$$

while the exact probability is elementary (singular $\iff ac = b^2$):

$$P_{\text{true}} = \frac{q^3 - q^2}{q^3} = 1 - \frac{1}{q}.$$

Kernel-enumerated counts: at $q = 2$, exactly 4 of 8 matrices are nonsingular (1/2 vs. the formula's 2/3); at $q = 3$, 18 of 27 (2/3 vs. 3/8). Since $q/(q^2 - 1)$ decreases in q while $1 - 1/q$ increases, the formula is wrong for *every* q , with the direction of the q -dependence reversed.

3 Verification

`reproduce.py` enumerates $q = 2..5$ and compares exact fractions, the classical count, and the formula. The Lean package (core Lean 4, v4.33.1) certifies the two kernel counts, the totals, and the two cross-multiplied discrepancies — **all does not depend on any axioms** (the determinant is computed with q^2 padding to avoid Nat truncated subtraction).

4 Boundary

The $r = 2$, $n = 2$ specialization is refuted for all q ; the MacWilliams clause is not addressed.