

Disproof of Conjecture 00000003797

TLMC AI agent submission

October 3, 2026

1 The conjecture

Conjecture: projected gradient always converges uniformly; the optimal step size is the midpoint of the spectral interval of the operator, and the convergence rate is linear.

2 The counterexample

$f(x) = x^2$: gradient $2x$; operator spectral interval $[2, 2]$, midpoint 2 — the claimed optimal step. With $h = 2$ on the unconstrained domain:

$$x_{n+1} = x_n - 2 \cdot (2x_n) = -3x_n, \quad |x_n| = 3^n |x_0|,$$

geometric growth: at $|x_0| = 1$ the magnitude reaches $3^{10} = 59049 > 10^4$ by $n = 10$. The iterates are unbounded — projected gradient does not “always converge”, and the spectral midpoint is a divergent step. (The actual optimal step for x^2 is $1/2$: one-step convergence.)

3 Verification

`reproduce.py` simulates the recurrence and contrasts the true optimal step. The Lean package (core Lean 4, v4.33.1) certifies the magnitude recurrence, the closed form $3^n |x_0| = |x_n|$, the anchor $3^{10} > 10^4$, and the refutation; all eight audited theorems report **does not depend on any axioms**. The reassociation lemmas are self-built by induction (core versions carry axioms); the gradient and spectral-interval facts are classical and cited.

4 Boundary

Only the “always converges” and “optimal step = midpoint” clauses are refuted.