

Disproof of Conjecture 3490: chromatic polynomials determine chromatic numbers

Local verified working submission
October 3, 2026

Claim. Conjecture 3490 asserts the existence of graph families with identical chromatic polynomial and spectral radius but different chromatic numbers, with minimum order ten. We prove that even a single pair of finite graphs with identical chromatic polynomial and different chromatic numbers cannot exist. Thus the extra spectral-radius condition and the stated minimum order cannot repair the assertion.

Definitions. For a finite graph G and a nonnegative integer k , a proper k -coloring is a function $V(G) \rightarrow \{0, \dots, k-1\}$ assigning different colors to adjacent vertices. The chromatic polynomial P_G is characterized by

$$P_G(k) = \#\{\text{proper } k\text{-colorings of } G\} \quad (k \in \mathbb{N}).$$

The chromatic number $\chi(G)$ is the least k admitting a proper k -coloring. These conventions also cover the empty graph, whose chromatic number is zero.

Proof. For every $k \in \mathbb{N}$,

$$G \text{ is } k\text{-colorable} \iff P_G(k) > 0.$$

Consequently $P_G = P_H$ implies that G and H admit proper k -colorings for exactly the same values of k . Their least such values agree, so $\chi(G) = \chi(H)$. Equivalently, if $\chi(G) < \chi(H)$, evaluation at $k = \chi(G)$ would give $P_G(k) > 0 = P_H(k)$, a contradiction; the reverse inequality is excluded in the same way. No pair claimed in the conjecture exists.

Lean verification and semantic correspondence. The self-contained file `Main.lean` imports only `Std`. Vertices are `Fin n`; the edge relation is arbitrary and decidable, so the result applies in particular to every finite simple graph. The list `words k n` enumerates the length- n lists with entries below k . Theorems `mem_words` and `words_nodup` establish completeness and absence of duplicates. The predicate `Proper` checks the edge constraints, and `chromaticEvaluation` counts the proper assignments. The theorem `evaluation_pos_iff_colorable` proves that this count is positive exactly when a proper coloring exists.

`ChromaticNumberIs` uses the actual existence and minimality definition. The theorem `equal_evaluations_equal_chromatic_numbers` proves equality of chromatic numbers for any two graphs with identical natural coloring counts. Integer polynomials are represented explicitly by coefficient lists and evaluated by Horner's rule in `polynomialEval`. The predicate `IsChromaticPolynomial` is precisely the counting characterization above. The final theorem `not_polynomial_separating_pair_claim` refutes the existence of two graphs sharing an actual integer polynomial with that characterization and having distinct chromatic numbers. Thus the polynomial-to-counting bridge is included in the formal statement, and not left as an assumed numerical coincidence.

Reproduction and trust. Run `lean Main.lean` using Lean 4.19.0. The command exits successfully. The principal results depend only on the standard Lean axioms `propext`, `Classical.choice`, and `Quot.sound`. There are no added axioms, placeholders, or uses of `native_decide`. The proof is uniform in the numbers of vertices and colors and is not limited to a bounded search.